INFORMATION TO USERS

This material was produced from a microfilm copy of the original document. While the most advanced technological means to photograph and reproduce this document have been used, the quality is heavily dependent upon the quality of the original submitted.

The following explanation of techniques is provided to help you understand markings or patterns which may appear on this reproduction.

1. The sign or "target" for pages apparently lacking from the document photographed is "Missing Page(s)". If it was possible to obtain the missing page(s) or section, they are spliced into the film along with adjacent pages. This may have necessitated cutting thru an image and duplicating adjacent pages to insure you complete continuity.
2. When an image on the film is obliterated with a large round black mark, it is an indication that the photographer suspected that the copy may have moved during exposure and thus cause a blurred image. You will find a good image of the page in the adjacent frame.
3. When a map, drawing or chart, etc., was part of the material being photographed the photographer followed a definite method in "sectioning" the material. It is customary to begin photoing at the upper left hand corner of a large sheet and to continue photoing from left to right in equal sections with a small overlap. If necessary, sectioning is continued again — beginning below the first row and continuing on until complete.
4. The majority of users indicate that the textual content is of greatest value, however, a somewhat higher quality reproduction could be made from "photographs" if essential to the understanding of the dissertation. Silver prints of "photographs" may be ordered at additional charge by writing the Order Department, giving the catalog number, title, author and specific pages you wish reproduced.
5. PLEASE NOTE: Some pages may have indistinct print. Filmed as received.

Xerox University Microfilms

300 North Zeeb Road
Ann Arbor, Michigan 48106

76-24,385

GRATZ, Ronald Karl, 1946-
ENERGETICS FOR ACTIVITY IN THE DIAMOND-
BACK WATER SNAKE, NATRIX RHOMBIFERA.

The University of Oklahoma, Ph.D., 1976
Zoology

Xerox University Microfilms, Ann Arbor, Michigan 48106

THE UNIVERSITY OF OKLAHOMA

GRADUATE COLLEGE

ENERGETICS FOR ACTIVITY IN THE DIAMOND-BACK WATER SNAKE,

NATRIX RHOMBIFERA

A DISSERTATION

SUBMITTED TO THE GRADUATE FACULTY

in partial fulfillment of the requirements for the

degree of

DOCTOR OF PHILOSOPHY

BY

RONALD KARL GRATZ

Norman, Oklahoma

1976

ENERGETICS FOR ACTIVITY IN THE DIAMOND-BACK WATER SNAKE,
NATRIX RHOMBIFERA

APPROVED BY

Victor A. Battison
Charles E. Haines
Howard Haines
Frank G. Lottner

DISSERTATION COMMITTEE

ACKNOWLEDGEMENTS

I would like to thank all members of my dissertation committee for their aid and encouragement throughout the various stages of this work: Dr. Charles C. Carpenter, Dr. Howard B. Haines, Dr. Victor H. Hutchison (chairman), and Dr. Frank J. Sonleitner. I am deeply indebted to Mildred Fung and Karla Zachary for technical assistance and to Neil Ford, James Gillingham and Pat Manville for aid in collecting animals. Further technical assistance of many kinds was provided by Dennis Andress, Michael Fair, Ladon Detherage and Jerry Black and to these and all others who contributed in any way, I am thankful.

This study was conducted while the author held a graduate teaching assistantship in the Department of Zoology, University of Oklahoma and was supported in part by NSF Grant BMS 73-06725 (to Victor Hutchison). This manuscript has been prepared in the style for submission to The Journal of Experimental Biology.

TABLE OF CONTENTS

	Page
List of Tables and Appendices	v
List of Illustrations	vi
Summary	1
Introduction	3
Materials and Methods	5
Results	10
Diurnal cycles	10
Levels of metabolic rate	10
Energy production during activity	11
Recovery	12
The respiratory exchange ratio	13
Discussion	14
Diurnal cycles	14
Levels of metabolic rate	14
Energy production during activity	15
Relative contributions of aerobiosis and anaerobiosis	16
Oxygen debt	18
Total energy expenditure for activity	20
Metabolite compartmentalization	20
Respiratory exchange ratio	22
Ecological implications	23
References	25
Tables	32
Figure Legends	35
Figures	36
Appendices	40

LIST OF TABLES AND APPENDICES

Table		Page
1	Metabolic rate and routine metabolic scope of the diamond-back water snake, <u>Natrix rhombifera</u>	32
2	Energy budget for 10 minutes forced activity in the diamond-back water snake, <u>Natrix rhombifera</u>	33
3	Tissue metabolites in <u>Natrix rhombifera</u> : controls and immediately after 10 minutes forced activity at 25°C .	34
Appendix		
I	Diurnal variation in oxygen consumption in the diamond-back water snake, <u>Natrix rhombifera</u>	40
II	Diurnal variation in routine activity in the diamond-back water snake, <u>Natrix rhombifera</u>	45
III	Oxygen consumption volumes and maximum sustained rates during activity and recovery from 10 minutes forced activity in the diamond-back water snake, <u>Natrix rhombifera</u>	50
IV	Metabolite concentrations before, immediately after and during recovery from 10 minutes forced activity at 25°C, 1400 hrs in the diamond-back water snake, <u>Natrix rhombifera</u>	51
V	Results of the regression analysis of "standard" metabolic rate vs acclimation temperature in <u>Natrix rhombifera</u> .	58

LIST OF ILLUSTRATIONS

Figure		Page
1	Diurnal cycle of oxygen consumption and routine activity in the diamond-back water snake, <u>Natrix rhombifera</u>	36
2	The effect of temperature and hour of measurement on metabolic scope for activity in <u>N. rhombifera</u>	37
3	Tissue metabolite levels in <u>N. rhombifera</u> during recovery from 10 minutes forced activity at 25°C, 1400 hrs	38
4	Total energy cost of 10 minutes forced activity in <u>N. rhombifera</u>	39

ENERGETICS FOR ACTIVITY IN THE DIAMOND-BACK WATER SNAKE,

NATRIX RHOMBIFERA

SUMMARY

The effect of routine and forced activity on total energetics of the diamond-back water snake, Natrix rhombifera, was studied at 15°C, 25°C and 35°C, all under a photoperiod of LD 12:12 with the onset of photophase at 0600 CDT.

This snake exhibited significant daily cycles of oxygen consumption at 15°C and 35°C but not at 25°C. At all temperatures, oxygen consumption increased at or just before the onset of photophase and at 25°C and 35°C is maximal at this time ($\dot{V}_{O_2} = 211.1 \mu\text{l g}^{-1} \text{ hr}^{-1}$ and $485 \mu\text{l g}^{-1} \text{ hr}^{-1}$ respectively). At 15°C, oxygen consumption is maximal midway thru photophase ($161.9 \mu\text{l g}^{-1} \text{ hr}^{-1}$). Significant cycles of routine activity exist at all temperatures and followed the oxygen consumption cycles at 15°C and 35°C. At 25°C activity was greatest in late photophase and early scotophase.

The snakes were stimulated electrically to maximum activity at 1400 hrs CDT at all temperatures and at 2200 hrs CDT at 25°C. Oxygen consumption and total body lactate production were measured and indicated a high dependence on anaerobiosis at all temperatures. Yields of ATP from aerobiosis and anaerobiosis respectively were: 1.4 and 16.2 $\mu\text{Moles g}^{-1}$ at 15°C, 1400 hrs; 5.4 and 30.2 $\mu\text{Moles g}^{-1}$ at 25°C, 1400 hrs;

4.8 and 17.7 $\mu\text{Moles g}^{-1}$ at 25°C, 2200 hrs; and 5.3 and 28.9 $\mu\text{Moles g}^{-1}$ at 35°C, 1400 hrs. The oxygen debt was 100.9 $\mu\text{l g}^{-1}$, 175.8 $\mu\text{l g}^{-1}$, 182.3 $\mu\text{l g}^{-1}$, and 108.0 $\mu\text{l g}^{-1}$ at these respective temperatures and times and could account for the oxidation of only 10 to 15% of the lactate formed. Tissue glycogen and blood glucose measurements indicate that most of the remaining lactate undergoes gluconeogenesis.

Tissue metabolites were measured immediately after and during recovery from activity at 25°C, 1400 hrs. Muscle lactate rose from 1.36 to 3.03 mg g^{-1} during activity while muscle glycogen dropped from 10.46 to 3.93 mg g^{-1} . Both returned to resting levels by 3 hr after activity. Blood and liver lactate increased from 0.09 to 1.08 mg ml^{-1} and from 0.30 to 0.93 mg g^{-1} respectively during activity. They continued to rise to a maximum at 1 hr of recovery and then declined to resting levels by 5 to 7 hrs of recovery. Blood glucose rose from 0.15 to 0.70 mg ml^{-1} during activity and continued to rise to a maximum at 5 hrs of recovery. There was no significant decline by 7 hrs of recovery.

The respiratory exchange ratio rose to well above 1.0 during the first hr of recovery, a result of blood lactate being buffered by the bicarbonate buffer system. The lack of an increased R during activity, a time when blood lactate has become elevated, indicates other buffers to be important during this time.

The strong dependence on anaerobiosis during activity is taken as an adaptation to a semi-aquatic life style in which only a short but powerful burst of activity is needed to escape a terrestrial predator and in which the snake would have little or no oxygen available for struggles with an aquatic predator.

INTRODUCTION

While lizards have been studied extensively, there are few studies of energetics during activity in snakes and only Gratz and Hutchison (1976) examined the contributions of both aerobic and anaerobic metabolism. They found that while the coachwhip snake, Masticophis flagellum, meets most of its increased energy demands during activity by anaerobiosis, lactate levels were generally indicative of a lesser reliance on anaerobiosis than is true in lizards (Moberly, 1968; Bennett and Dawson, 1972; Bennett and Licht, 1972). The coachwhip, however, is a fast moving animal that may often sustain rapid movement for long periods of time, a situation not compatible with high lactate production and the resultant acidotic condition.

The diamond-back water snake, Natrix rhombifera, a resident of most bodies of water in the south-central United States and northern Mexico, rarely ventures far from the water's edge (Preston, 1970). As such it must only make short dashes to reach the safety of the water and avoid terrestrial predators. Aquatic predators would find the snake in an essentially anaerobic environment (for the snake). Either situation would favor anaerobiosis over aerobiosis for energy production.

Many reptiles exhibit maximum energy production at or near their preferred body temperature (Dawson, 1975) but also show a degree

of of temperature independence of various aspects of metabolism within their normal body temperature range. Apparently no study of the affect of temperature on total energetics during activity of snakes has been made.

Strategies of recovery have been little investigated in reptiles. Moberly (1968) presented data on blood lactate during recovery in Iguana iguana and Bennett and Licht (1972) followed muscle, liver and blood lactate thru the first 90 min of recovery in Anolis carolinensis. Bennett (1972a) presented data on the extent and repayment time of the oxygen debts of the racehorse goanna, Varanus gouldii, and the spiny chuchwalla, Sauromalus hispidus. The mechanism of lactate removal and glycogen replenishment have not been investigated for any reptile.

I undertook this study to determine (1) the level of routine oxygen consumption in the diamond-back water snake, Natrix rhombifera, and its variation with a diurnal light-dark cycle, acclimation temperature and routine activity; (2) the relative contributions of aerobic and anaerobic metabolism to increased energy production during forced activity and, by including the oxygen debt, the total cost of the activity; (3) the mechanism of recovery from the activity.

MATERIALS AND METHODS

Natrix rhombifera from 60g to 1400g were collected over a two year period from areas within 170 km of Norman, Oklahoma. Since no sexual differences were found in the results, the data from both sexes were combined. One to four snakes were housed together in plastic containers and acclimatized at 15°C, 25°C or 35°C and a photoperiod of LD 12:12 for at least one week before use. Snakes scheduled for experimentation within one month after capture were not fed. Others were fed live or frozen fish at irregular intervals. As water dishes kept in the containers tended to be spilled almost immediately with the resultant danger of blister disease (Frye, 1973) no water was left in the containers. Snakes were soaked in a bucket of water for 30 min to 1 hr every 2 - 4 days and before introduction into an experimental chamber. All measurements were made at the acclimation temperatures.

The following definitions were used:

Aerobic scope. This is the metabolic scope of Fry (1947) and represents the difference between the minimum and maximum sustained oxygen consumption rates under given conditions.

Index of metabolic expansibility. This is the ratio of the maximum sustained metabolic rate during activity to the minimum rate at the same time and temperature.

Aerobic capacity. This is the amount of oxygen consumed above resting during the activity period.

Anaerobic capacity. According to Bennett and Licht (1972) this is the total amount of lactate produced above the resting level during the activity period. This is energetically equivalent to the oxygen deficit as defined by Knuttgen (1971).

Oxygen debt. According to Knuttgen (1971) this is the amount of oxygen consumed above resting during the recovery period.

Gas exchange was measured by open system respirometry. The snakes were placed in individual Plexiglas chambers (2400 - 4000 cc) 4 to 8 hr prior to the start of measurements. Air was drawn through the chamber at 300 to 600 cc min⁻¹, depending upon the size of the animal and the temperature used, through a tube of dessicant and into the oxygen analyzer (Applied Electrochemistry Model S-3A). When respiratory exchange ratio (R) measurements were to be made a carbon dioxide analyzer (Beckman Model 15A) was connected in series between the oxygen analyzer and the pump. The volume of the tubing from the animal chamber to the oxygen analyzer was 50 to 60 cc and produced a negligible lag in readings. Gas concentrations could be read directly from the analyzer displays or recorded continuously on a Linear Instruments Model 485 dual channel recorder. Oxygen consumption was calculated from the Depocas and Hart (1957) formula for open systems. All gas volumes are presented at STPD. Ten individuals were used at each acclimation temperature for determination of the 24-hr cycles of oxygen consumption.

Maximum activity was induced by electrical stimulation through safety pin electrodes attached through the skin of the tail and leading

from a Grass SM6 stimulator set at 100 vAC with 5 pulses s^{-1} of 1 msec duration. Non-stimulated animals at corresponding times and acclimation temperatures were examined as controls (N = 10 individuals for $\dot{V}_{O_2}$ and 5 - 14 individuals for biochemical data). Oxygen consumption during activity and recovery was measured on 7 - 10 individuals to yield N = 10 at each acclimation temperature. Measurements were made by integration of the area under the oxygen pen tracing of the recorder with a polar planimeter and performing the necessary calculations to yield ml O_2 . Recovery oxygen consumption was measured from the end of activity until oxygen consumption either returned to the resting rate for that individual or, after a significant decrease, levelled out above the rest resting rate and remained constant for at least 60 min. This latter phenomenon is ascribed to an increased resting metabolism and has been described for mammals (Knuttgén, 1971) and snakes (Gratz and Hutchison, 1976). The oxygen debt was determined by subtracting the resting volume from the total volume of oxygen consumed during recovery.

Measurement of total body lactate required maceration of the entire animal (Bennett and Licht, 1972). The snakes (N = 5 for each condition) were decapitated and chopped into approximately 8 cm segments before homogenization in 4x their weight of perchloric acid. Fifty ml of the solution was further homogenized in 60 ml perchloric acid and then stored on ice. For tissue metabolites, the snakes (N = 5 - 7 for each condition) were decapitated and 1 ml of blood was taken from the severed trunk for analyses of blood glucose and blood lactate. Since dissection of the trunk produced strong squirming due to spinal reflexes, the trunk was spinal pithed before dissection. Liver and muscle samples

were removed and stored on ice until weighed. The longissimus, the articularis and the spinalis series of muscles lateral to the vertebrae were used; all are of major importance in snake locomotion (Gans, 1962). Samples for lactic acid analysis were homogenized in 10x their weight of perchloric acid. All lactate homogenates were then centrifuged for 10 min at 3000 rpm and the supernatant recentrifuged at 10,000 rpm for 10 min, all at 0°C. Duplicate 100 ul samples of the final supernatant and of the processed blood plasma were analyzed with enzymatic kits (Lactate-UV method, Boehringer-Mannheim Corp.).

Tissue samples for glycogen analyses were homogenized in deproteinizing solution (50g trichloroacetic acid and 1g Ag_2SO_4 in 1 liter water) to provide a 1:100 dilution (liver) or 1:20 dilution (muscle). The tube containing the homogenate was placed in boiling water for 15 min, cooled and centrifuged at 3000 rpm for 10 min. One ml of supernatant was drawn and analyzed by the method of Kemp and Van Heijningen (1954). All lactate and glycogen determinations were made on a Varian Techtron Model 635 spectrophotometer. Blood glucose was determined by the "Glucostat" method (Worthington Biochemical Corp.) on a Beckman Spinco 151 spectrophotometer. All metabolite data are presented as mean $\pm$ 1 standard error mg g^{-1} tissue weight or mg ml^{-1} blood plasma.

Energy production is presented in $\mu\text{Moles ATP}$ formed per gram body weight. Caloric values are also presented. Bennett and Licht (1972) provide the following conversions for determining ATP yields from aerobiosis and anaerobiosis:

$$1.0 \text{ mg lactate formed} = 0.0167 \text{ millimoles ATP}$$

$$1.0 \text{ ml } O_2 \text{ (STPD) consumed} = 0.290 \text{ millimoles ATP}$$

This assumes glycogen to be the only substrate catabolized and the latter figure also assumes a P/O ratio of 3 with a net yield of 39 moles ATP formed per mole of glycogen utilized. This latter figure is still in doubt for skeletal muscle due to the uncertainty of the fate of glycolytic NADH (Greville, 1969; Newsholme and Start, 1973). Caloric equivalents are taken from Hochachka and Somero (1973):

$$1.0 \text{ mg lactate formed} = -0.2609 \text{ cal}$$

$$1.0 \text{ ml } O_2 \text{ (STPD) consumed} = -5.104 \text{ cal}$$

Both of these are based on glucose as the substrate utilized.

The 24-hr activity cycles were determined on 5 - 6 individuals at each acclimation temperature. Measurements were made with mercury switches taped horizontally on the side of the snake at approximately the level of the heart. The animals were placed in individual plastic boxes inside an environmental chamber 4 - 8 hr prior to the start of an experiment. The mercury switch leads were connected to a Rustrak Model 292-8 recorder and the number of pen movements per hour was used as an indicator of activity. Activity was not recorded simultaneously with, nor necessarily on, the same animals as the $\dot{V}O_2$ measurements.

An analysis of variance was performed on the data to determine if significant variation in metabolism or activity occurred over the 24-hr period. Because activity was recorded as a percentage, an arcsin transformation was made on these data. Duncan's multiple range test (Duncan, 1955) was used to identify which hours were significantly different in the 24-hr cycle. Metabolite levels were compared by the Student t-test or the Duncan multiple range test. Significance was judged at the $P < 0.05$ level.

RESULTS

Diurnal cycles

There was a significant ($P < 0.05$) daily variation of oxygen consumption at 15°C and 35°C and of routine activity at all temperatures (Fig. 1). Oxygen consumption increased at or shortly before the onset of photophase at all temperatures, and at 25°C and 35°C it was maximal at this time and then declined with only random fluctuations during the remainder of the 24-hr cycle. Routine activity followed this pattern closely at 35°C but not at 25°C . At 15°C the cycles were essentially diurnal, activity and oxygen consumption being significantly depressed during most of the scotophase.

Levels of metabolic rate

The grand mean $\dot{V}_{\text{O}_2}$ for 24 hrs at each acclimation temperature is presented in Table 1 along with the corresponding $\dot{V}_{\text{CO}_2}$, R and Q_{10} values. Hutchison and Kohl (1971) suggested that the difference between the metabolic rates of the low hours of the daily cycles and the high hours ("routine" metabolic scope) might be a better indicator of the animal's energetics under natural conditions than the metabolic scope of forced maximal activity. The values for the lowest two or three hour period and the highest hour of the daily cycle are also presented in Table 1 along with the routine metabolic scope.

The mean $\dot{V}_{O_2}$ of the low hours of the daily cycle was taken as the "standard" metabolic rate at each temperature and this value was regressed on acclimation temperature to yield the equation:

$$\dot{V}_{O_2} = 12.11 T_a - 166.55$$

Where $\dot{V}_{O_2}$ is presented in $\mu\text{l g}^{-1} \text{ hr}^{-1}$ STPD and T_a is the acclimation temperature in degrees celsius. An analysis of variance showed significant linear regression ($P < 0.05$) and insignificant deviation from linearity ($P > 0.25$).

Energy production during activity

Snakes were stimulated to maximum activity for 10 min at 1400 hrs at all acclimation temperatures. Snakes at 25°C were also stimulated at 2200 hrs to determine if a diurnal pattern of maximum energy expenditure exists. Both are times of stable oxygen consumption intermediate between the high and low rates of the diurnal cycle at the respective temperatures. The snakes responded to stimulation with an initial burst of violent thrashing similar to that described for Dipsosaurus (Bennett and Licht, 1972; Bennett and Dawson, 1972), but absent in Masticophis (Gratz and Hutchison, 1976). This lasted for only 1 - 2 min after which movement became slow and labored. During recovery the snakes lay quietly with little movement.

Oxygen consumption increased only slightly during the initial 3 - 5 min of activity but thereafter showed a marked increase to a level roughly double the resting rate. Within the first 30 min following activity oxygen consumption increased slightly above this rate. The maximum sustained oxygen consumption rates attained during activity and recovery are shown in Fig. 2 along with the corresponding resting rate

aerobic scope for activity and index of metabolic expansibility.

Mean values for oxygen consumed and total body lactate produced during activity are presented in Table 2. At 15°C the total energy expenditure during activity is lowest (measured as total $\mu\text{M ATP g}^{-1}$), as is the relative contribution of aerobiosis. The greatest energy production is at 25°C, 1400 hrs but the values obtained are only slightly above those at 35°C. At 25°C, 2200 hrs aerobic energy production is only slightly below that of 1400 hrs ($P > 0.50$) but lactate production and therefore total energy expenditure are far below those at 1400 hrs.

Tissue metabolites were measured immediately after activity at 25°C, 1400 hrs and 2200 hrs, and were compared with controls for the same hour (Table 3). All tissue lactate levels and blood glucose increased significantly ($P < 0.001$) during activity. Muscle glycogen decreased significantly ($P < 0.01$) at 1400 hrs but not at 2200 hrs ($P > 0.10$). Liver glycogen showed a slight decrease at both times but neither change was significant.

Recovery

Oxygen consumption returned to or to near resting levels in approximately 2 - 2.5 hrs at 15°C and 25°C and in 1 hr at 35°C. The oxygen debt (Table 2) was lowest at 15°C and 35°C and highest at 25°C.

Total body lactate and tissue metabolites were monitored for 7 hours after activity at 25°C, 1400 hrs (Fig. 3). Total body lactate and muscle lactate were maximum immediately after activity and then declined steadily. Muscle lactate required 3 hrs to reach pre-activity levels while total body lactate required 5 hrs. Muscle glycogen was minimum immediately after activity, rose to a value not significantly different

from the pre-activity level by the first hour of recovery and continued to rise until 5 hrs after activity. Blood and liver lactate required 7 hrs to return to control levels while blood glucose remained elevated throughout the recovery period. Liver glycogen showed no significant change throughout recovery.

The respiratory exchange ratio

The respiratory exchange ratio (R) was measured before, during and during recovery in some animals (N = 2 - 7) under all conditions. At 15°C, 1400 hrs, 25°C, 1400 hrs and 25°C, 2200 hrs R decreased from mean pre-activity levels of 0.88, 0.79 and 0.78 to 0.76, 0.66 and 0.57, respectively, during the activity period. At 35°C, 1400 hrs, R increased from 0.78 before activity to 0.85 during activity. During the initial hour of recovery R increased rapidly to 0.98, 1.31, 1.23 and 1.03, respectively, under the above conditions. Thereafter it gradually decreased to resting levels.

DISCUSSION

Diurnal cycles

The diamond-back water snake is essentially a diurnal animal during the cool weather of early spring but divides its activity equally between day and night during warmer weather (Preston, 1970). Personal field observations and reports of fish hatchery personnel find that the animal may be active and feeding anytime from before sunrise to past midnight during mid-summer. The activity and oxygen consumption patterns at 15°C and 35°C suggest that photoperiod alone may regulate the snakes' activity at extreme temperatures. At 25°C some activity is present at almost every hour and examination of individual records indicates that different animals are active at various hours but with major concentrations of activity at mid-afternoon and early evening. The general lack of agreement of activity and oxygen consumption cycles at 25°C is puzzling but not unprecedented. Hutchison and Kohl (1971) found $\dot{V}O_2$ to be independent of activity in some individual Bufo marinus and theorized the latter to be circadian while the former is not.

Levels of metabolic rate

The routine metabolic scope (Table 1) roughly involves a doubling of the minimum metabolic rate. Hutchison and Kohl (1971) found similar results in the marine toad, Bufo marinus, as did Gratz and Hutchison (1976) in the coachwhip. By comparison, the forced metabolic

scopes of Natrix rhombifera at 15°C and 25°C are approximately double the respective routine metabolic scopes while at 35°C the two are roughly equivalent (Fig. 2). This is most likely due to the greater intensity of activity at 35°C. Although there is not close agreement, the general correlation of major peaks of activity and oxygen consumption suggest that routine activity employs aerobiosis as its major energy source.

The mean low oxygen consumption values reported here at 25°C and 35°C are much higher than those reported for this species by Jacobson and Whitford (1970) or for other species (see Dmi'el, 1971, for a review), but the values reported here at 15°C are below those otherwise reported. The linear relationship of mean low $\dot{V}O_2$ to acclimation temperature found in this study is in contrast to curvilinear relationships found for this species and the garter snake, Thamnophis proximus (Jacobson and Whitford, 1970), for the gopher snake, Pituophis catenifer, (Greenwald, 1971), and for the desert snake, Spalerosophis cliffordi, (Dmi'el and Borut, 1972). In all these studies, however, a larger range of temperatures was examined than was used in this study, and the animals were not acclimated to each temperature used. The differences between the metabolic rates of acclimated animals and unacclimated animals suggests a clockwise rotation of metabolic rate with acclimation (type III, Prosser, 1973).

Energy production during activity

The initial burst of violent activity is the time of the greatest lactate production in lizards (Bennett and Licht, 1972) and presumably other reptiles. Where this initial violent activity is absent (e.g. Masticophis, Gratz and Hutchison, 1976) lactate production is low. In Natrix rhombifera the initial burst of activity is pronounced and the

lactate levels achieved at 25°C, 1400 hrs and 35°C, 1400 hrs are among the highest reported in reptiles (Bennett and Dawson, 1972; Bennett and Licht, 1972).

The maximum aerobic scope of N. rhombifera is $179.5 \mu\text{l O}_2 \text{ g}^{-1} \text{ hr}^{-1}$ at 35°C, 1400 hrs, considerably less than the maximum values reported for Pituophis ($420 \mu\text{l O}_2 \text{ g}^{-1} \text{ hr}^{-1}$, Greenwald, 1971) or Spalerosophis ($265 \mu\text{l O}_2 \text{ g}^{-1} \text{ hr}^{-1}$, Dmi'el and Borut, 1972) but higher than that of Masticophis ($159.8 \mu\text{l O}_2 \text{ g}^{-1} \text{ hr}^{-1}$, Gratz and Hutchison, 1976). As this rate was usually achieved during the latter 5 min of activity, a time when movements had become slow and labored and when some lactate oxidation may have been occurring, the aerobic scope during maximum activity is considerably less than this value.

Relative contributions of aerobiosis and anaerobiosis

Under all conditions studied, anaerobiosis was the primary source of energy for activity, providing from 78.7 to 91.9% of the total ATP generated above standard energy production (Table 2). Even these values may be somewhat conservative as the high liver lactate values following activity suggest that some of the increased oxygen consumption may be due to lactate oxidation by the liver and thus would not contribute ATP to the muscle. The greatest lactate production (1.81 mg g^{-1}) occurred at 25°C, 1400 hrs and is identical to the maximum lactate production of Dipsosaurus dorsalis (1.82 mg g^{-1} at 40°C; Bennett and Dawson, 1972) which has the highest lactate production of any lizard studied. The highest aerobic capacity ($18.79 \mu\text{l O}_2 \text{ g}^{-1}$) of Natrix also occurs at 25°C, 1400 hrs but this is much lower than that of Dipsosaurus (Bennett and Dawson, 1972) or the marine iguana, Amblyrhynchus cristatus

(Bennett, Dawson and Bartholomew, 1975) and is roughly the same as recorded for Masticophis (Gratz and Hutchison, 1976). Moreover, the values for lizards are for two minutes while those for snakes are for ten minutes of activity. The smaller size of Dipsosaurus contributes in part to this difference, but the specimens of Amblyrhynchus used in the above study were of the same general size as the snakes used in this study. The lower aerobic capacity of snakes may be due to their lung structure which for most species, including Natrix, is composed of a small alveolar section approximately one-third of the total lung volume, and a large, poorly vascularized air sac with little or no gas exchange capacity (George and Shah, 1956, 1965; Brongersma, 1957; Kardong, 1972a & b; McDonald, 1959). This reduced aerobic capacity is partially offset by the greater efficiency of snake locomotion, less than half the energy is required by a garter snake to move a given distance than by a monitor lizard of the same size and moving at the same speed (Chodrow and Taylor, 1973).

Aerobic and anaerobic capacity and aerobic scope all have a high thermal dependance between 15°C and 25°C ($Q_{10} = 3.81, 1.87$ and 2.27 , respectively) but not between 25°C and 35°C ($Q_{10} = 0.99, 0.95$ and 1.10 , respectively). The mean body temperature of Natrix rhombifera measured in the field is 27.46°C overall and 28.51°C for active individuals (Preston, 1970). The range of all body temperatures is 19°C to 32°C. Several lizards show a degree of thermal independence of various aspects of metabolism at body temperatures near their activity range (Dawson, 1967, 1975; Bennett, 1972a; Moberly, 1968). Dmi'el and Borut (1972) reported only a slight difference in the aerobic scope of Spalerosophis

cliffordi from 25° to 40°C and Hoskins and Aleksuk (1973b) reported Q_{10} s approaching 1.0 in the metabolic rate of tissues of the garter snake, Thamnophis sirtalis parietalis, measured within the animal's preferred body temperature range. Studies on the effect of temperature on the kinetics of two metabolic enzymes of the garter snake indicate that temperature dependant isoenzymes and positive thermal modulation of enzyme activity may be the basis of the low Q_{10} s in the animal's activity range (Aleksiuk, 1971; Hoskins and Aleksuk, 1973a). This phenomenon may allow the animal to generate maximum energy for activity over a wide range of environmental temperatures and thus obtain a degree of metabolic homeostasis within its normal activity range (Dawson and Bartholomew, 1956; Bennett, 1972a; Hoskins and Aleksuk, 1973a & b).

At 25°C, 2200 hrs there is a significant ($P < 0.01$) decrease in total body lactate production but no significant change in aerobic energy production ($P < 0.50$). Although diurnal variation in routine metabolic rate (Roberts, 1968; Hutchison and Kohl, 1971; Turney and Hutchison, 1974; Gratz and Hutchison, 1976), aerobic scope (Turney and Hutchison, 1974) and routine blood lactate levels (Hutchison and Turney, 1975) have been reported in ectotherms, this is the first indication of diurnal variation in anaerobic energy production in an ectothermic vertebrate.

Oxygen debt

The oxygen debt can be divided into an alactacid debt, attributable to resynthesis of depleted high energy phosphates and reoxygenation of respiratory pigments, and a lactacid debt, attributable to the oxygen needed to provide energy for gluconeogenesis. In mammals

roughly 80% of the lactate formed during activity is oxidized and 20% is reconverted to glucose (Minaire, 1973). Oxidation may take place in many organs, but the liver and kidney are the primary sites of gluconeogenesis. Liver and possibly kidney oxidation of lactate is accompanied by an increased oxygen uptake while oxidation by other tissues is not, the lactate merely substituting for other substrates (Knuttgén, 1971; Kramer, Driessen and Brechtelsbauer, 1971; Minaire, 1973).

Six moles of ATP are required to convert 2 moles of lactate to 1 mole of glucose via gluconeogenesis and 36 moles of ATP are produced by the oxidation of two moles of lactate. Thus the oxidation of 14.3% of the lactate formed would provide enough ATP to reconvert the remaining 85.7% to glucose ($0.143 \times 36 = 0.857 \times 6$). The measured oxygen debts would account for the oxidation of 13.8%, 12.6%, 22.8% and 8.2% of the lactate formed at 15°C, 1400 hrs, 25°C, 1400 hrs, 25°C, 2200 hrs, and 35°C, 1400 hrs respectively, assuming no alactacid debt. A significant alactacid debt would reduce this figure somewhat but, since some lactate is most likely oxidized in tissues not contributing to the oxygen debt, lactate oxidation could account for most or all of the energy needed for gluconeogenesis. It appears that only a minimum of the lactate is oxidized and that the primary route of lactate removal is by gluconeogenesis. Data on metabolite compartmentalization lend further support to this hypothesis (see below). The low oxygen debt at 35°C may reflect a greater use of lactate as substrate for normal energetics and a consequent lower level of gluconeogenesis.

Total energy expenditure for activity

The total energy cost of activity must include both the energy expended during activity and any energy utilized to restore metabolic homeostasis. The former parameter has been calculated or estimated for a number of reptiles (Moberly, 1968; Bennett, 1972a; Bennett and Dawson, 1972; Bennett and Licht, 1972; Gatten, 1974; Bennett, Dawson and Bartholomew, 1975; Gratz and Hutchison, 1976), but the latter has gone largely ignored. The cost of relieving the metabolic acidosis incurred by the high lactate build-up and for restoring the depleted muscle glycogen stores is a significant added energy cost to the animal and must be accounted for in determinations of animal energy budgets. From the evidence provided above, it appears that most of the oxygen debt is due to lactate oxidation which occurs primarily in the liver and kidney (Knuttgen, 1971; Kramer, Driessen and Brechtelsbauer, 1971; Minaire, 1973). The oxidation of a mole of lactate produces 18 moles ATP and has a free energy change of -326 kcal (Hoch, 1971). These are equivalent to 0.268 millimoles ATP and -4.85 cal per ml O_2 STPD of the oxygen debt (Table 2). The sum of the values for the oxygen debt and for energy utilization during activity represents the total energy cost of the activity. This is presented for Natrix rhombifera in Fig. 4.

Metabolite compartmentalization

The decrease in muscle glycogen (Table 3) is more than adequate to account for all of the muscle lactate formed at 25°C. Moberly (1968) noted the same to be true for Iguana. In Iguana, however, blood glucose did not change after activity while in Natrix blood glucose rose

considerably. The duration of activity may explain the difference, as Moberly stimulated Iguana for 2 - 3 min while Natrix was stimulated for 10 minutes. Hutchison and Turney (1975) provide convincing evidence that the duration of activity influences blood glucose levels. There are indications at both hours of stimulation of a decrease in liver glycogen during the period of stimulation, although the differences noted are not significant due to the high variability of this metabolite. This initial drop, if real, may be the source of the initial rise in blood glucose. Blood and liver lactate levels have risen considerably during activity as perfusion of the muscle has carried some lactate to the liver for oxidation or gluconeogenesis.

After activity, the first priority is that of restoring the muscle to resting conditions. A large amount of lactate must be removed and the depleted muscle glycogen must be restored. All of this is accomplished within three hours after activity, approximately the time for oxygen consumption to return to normal. It appears that lactate is removed from the muscle and carried to the liver. As has been shown, just enough lactate appears to be oxidized to provide the energy for gluconeogenesis of the remainder, and the greatest degree of oxidation occurs within the first 60 to 90 minutes of recovery. The initial strategy appears to be both oxidation and gluconeogenesis to remove lactate as quickly as possible. The glucose manufactured at this time is returned to the blood, as evidenced by the continued rise in blood glucose, and this is in turn carried to the muscle to replenish the glycogen stores. The lack of any change in liver glycogen after activity indicates that this organ does not contribute glucose for restoration of muscle glycogen during recovery, as it does in some fishes (Black,

Robertson and Parker, 1961). After three hours all the excess lactate has been removed from the muscle and glycogen stores have been replenished. The excess lactate is now stored in the blood, liver and perhaps other organs. This lactate may be used as substrate by many organs but the continued rise in blood glucose and muscle glycogen indicates continued gluconeogenesis in the liver and subsequent glycogen synthesis in the muscle. By 7 hr after activity all excess lactate has disappeared and both blood glucose and muscle glycogen levels have begun to decline.

This system is then quite different from that in fish where liver glycogen contributes large amounts of glucose for muscle glycogen restoration (Black, Robertson and Parker, 1961), the liver glycogen being replenished at the next feeding. The long periods between feedings in snakes may have necessitated this unique strategy which places a high priority on gluconeogenesis to restore muscle glycogen and thus spare liver glycogen for other needs.

Respiratory exchange ratio

Lactic acid entering the blood of mammals is buffered primarily by the bicarbonate buffer system and results in an excess release of CO_2 in the lungs (Masoro and Siegel, 1971). The increased R during the first hour of recovery is undoubtedly due to this release of CO_2 . However, several questions arise which will require further investigation.

The decreased R during activity is due to a proportionately greater increase in $\dot{V}_{\text{O}_2}$ than in $\dot{V}_{\text{CO}_2}$. Carbon dioxide release does not begin to increase appreciably until approximately 10 minutes after

activity, a time when blood lactate usually reaches its peak (Moberly, 1968; Bennett, 1973). This is puzzling, since blood lactate has risen considerably by the end of the activity period. Bennett (1973) provided evidence that buffers other than bicarbonate and phosphate may be important in regulating blood pH during metabolic acidosis in reptiles. This may be particularly true in Natrix rhombifera whose blood bicarbonate level is low compared to other reptiles (7 mM l^{-1} ; Dessauer, 1970). The changes in R observed here suggest that the bicarbonate buffer system of Natrix may only be used when blood lactate becomes extremely high.

Ecological implications

The initial burst of violent twisting and thrashing during stimulation is the time of greatest lactate production. The activity of phosphofructokinase, the rate limiting enzyme of glycolysis, is much greater in lizards than in the laboratory rat (Bennett, 1972b). While interpolation to all reptiles and wild mammals is hazardous, if valid it would mean that a reptile can generate large amounts of energy anaerobically at a more rapid rate than a mammal over a short period of time. This power output may give the reptile a decided advantage in short-lived encounters, and this could mean the difference between life and death (Coulson and Hernandez, 1974). For any prolonged battle, the low aerobic scope of most reptiles would put them at a decided disadvantage.

Predator avoidance would seem to be the most likely situation requiring maximum energy expenditure in the water snake. While a burst of speed and a small struggle may also be necessary in subduing prey,

Anderson (1975) and personal observations on wild and captive specimens feeding on live fish did not show any intense activity. The most likely predators on Natrix, at least in Oklahoma, are racoons, hawks, coyotes, snapping turtles, predatory fish, other snakes, and, of course, man. While no study of the responses of Natrix to a natural predator has been made, personal observations made while capturing animals for this study reveal the following:

When a snake is grasped, the response is an immediate, violent twisting and thrashing of the entire body accompanied by vicious striking and a copious anal discharge. If the snake were being held in the mouth of a predator, this behavior would result in the musk being discharged directly into the eyes, nose and mouth of the predator. This might well result in a momentary shock during which the violent thrashing may be sufficient to break the snake free. Were this encounter to take place under water, the animal would have to rely entirely on anaerobiosis and a high anaerobic capacity would be essential for survival.

REFERENCES

- Aleksiuk, M. (1971). An isoenzymic basis for instantaneous cold compensation in reptiles: lactate dehydrogenase kinetics in Thamnophis sirtalis. Comp. Biochem. Physiol. 40B, 671 - 681.
- Anderson, K. (1975). Hunting behavior in captive Natrix erythrogaster transversa and Natrix rhombifera rhombifera. Unpublished manuscript deposited in the library of the University of Oklahoma Biological Station, Willis, Oklahoma.
- Bennett, A. (1972a). The effect of activity on oxygen consumption, oxygen debt and heart rate in the lizards Varanus gouldii and Sauromalus hispidus. J. Comp. Physiol. 79, 259 - 280.
- Bennett, A. (1972b). A comparison of activities of metabolic enzymes in lizards and rats. Comp. Biochem. Physiol. 42B, 637 - 647.
- Bennett, A. (1973). Blood physiology and oxygen transport during activity in two lizards, Varanus gouldii and Sauromalus hispidus. Comp. Biochem. Physiol. 46A, 673 - 690.
- Bennett, A. and P. Licht. (1972). Anaerobic metabolism during activity in lizards. J. Comp. Physiol. 81, 277 - 288.
- Bennett, A. and W. Dawson. (1972). Aerobic and anaerobic metabolism

during activity in the lizard Dipsosaurus dorsalis. J. Comp. Physiol. 81, 289 - 299.

Bennett, A., W. Dawson and G. Bartholomew. (1975). Effects of activity and temperature on aerobic and anaerobic metabolism in the Galapagos marine iguana. J. Comp. Physiol. 100, 317 - 329.

Black, E., A. Robertson and R. Parker. (1961). Some aspects of Carbohydrate metabolism in fish. In Comparative Physiology of Carbohydrate Metabolism in Heterothermic Animals (ed. A. Martin). Seattle: Univ. Washington Press.

Brongersma, L. (1957). Notes upon the trachea, the lungs and the pulmonary artery in snakes. I, II, III. Proc. Kon. Ned. Akad. Wetensch., Ser. C 60, 299 - 308, 309 - 313, 451 - 457.

Chodrow, R. and C. Taylor. (1973). Energetic cost of limbless locomotion in snakes. Fed. Proc. 32, 422.

Coulson, R. and T. Hernandez. (1974). Intermediary metabolism of Reptiles. In Chemical Zoology: Volume IX Amphibia and Reptilia. New York and London: Academic Press.

Dawson, W. (1967). Interspecific variation in physiological responses of lizards to temperature. In Lizard Ecology: A Symposium (ed. W. W. Milstead). Columbia: Univ. Missouri Press.

Dawson, W. (1975). On the physiological significance of the preferred body temperatures of reptiles. In Perspectives of Biophysical Ecology (ed. D. Gates and R. Schmerl). New York: Springer-Verlag.

- Dawson, W. and G. Bartholomew. (1956). Relation of oxygen consumption to body weight, temperature and temperature acclimation in lizards Uta stansburiana and Sceloporus occidentalis. *Physiol. Zool.* 29, 40 - 51.
- Depocas, F. and J. Hart. (1957). Use of the Pauling oxygen analyzer for measurement of O_2 consumption of animals in open circuit systems and in short-lag, closed circuit apparatus. *J. Appl. Physiol.* 10, 388 - 392.
- Dessauer, H. (1970). Blood chemistry of reptiles: physiological and evolutionary aspects. In Biology of the Reptilia vol. 3 (ed. C. Gans and T. Parsons). London and New York: Academic Press.
- Dmi'el, R. (1972). Relation of metabolism to body weight in snakes. *Copeia* 1972, 179 - 181.
- Dmi'el, R. and A. Borut. (1972). Thermal behavior, heat exchange and metabolism in the desert snake, Spalerosophis cliffordi. *Physiol. Zool.* 45, 78 - 94.
- Duncan, D. (1955). Multiple range and Multiple F-tests. *Biometrics* 11, 1 - 6.
- Fry, F. (1947). Effects of the environment on animal activity. *Pub. Ontario Fish Res. Lab.* 68, 1 - 62.
- Frye, F. (1973). Husbandry, Medicine and Surgery in Captive Reptiles. Bonner Springs: VM Publishing Co.

Gans, C. (1962). Terrestrial locomotion without limbs. Amer. Zool.
2, 167 - 182.

Gatten, R. (1974). Effects of temperature and activity on aerobic and
anaerobic metabolism and heart rate in the turtles Pseudemys
scripta and Terrapene ornata. Comp. Biochem. Physiol. 48A,
619 - 648.

George, J. and R. Shah. (1956). Comparative morphology of the lung in
snakes with remarks on the evolution of the lung in reptiles.
J. Anim. Morph. Physiol. 3, 1 - 7.

George, J. and R. Shah. (1965). Evolution of the air sacs in Sauropsida.
J. Anim. Morph. Physiol. 12, 255 - 263.

Gratz, R. and V. Hutchison. (1976). Metabolic scope for activity in the
coachwhip snake, Masticophis flagellum. J. Exp. Zool. (manuscript
submitted).

Greenwald, O. (1971). The effect of body temperature on oxygen
consumption and heart rate in the Sonora Gopher snake, Pituophis
catenifer affinis Hallowell. Copeia 1971, 98 - 106.

Greville, G. (1969). Intracellular compartmentalization and the citric
acid cycle. In Citric Acid Cycle: Control and Compartmentalization
(ed. J. Lowenstein). New York and London: Marcel Dekker.

Hoch, F. (1971). Energy Transformations in Mammals: Regulatory
Mechanisms. Philadelphia: W.B. Saunders Co.

- Hochachka, P. and G. Somero. (1973). Strategies of Biochemical Adaptation. Philadelphia: W. B. Saunders Co.
- Hoskins, M. and M. Aleksuk. (1973a). Effects of temperature on the kinetics of malate dehydrogenase from a cold climate reptile, Thamnophis sirtalis parietalis. Comp. Biochem. Physiol. 45B, 343 - 353.
- Hoskins, M. and M. Aleksuk. (1973b). Effects of temperature, photoperiod and season on in vitro metabolic rates of tissues from Thamnophis sirtalis parietalis, a cold climate reptile. Comp. Biochem. Physiol. 45A, 737 - 756.
- Hutchison, V. and M. Kohl. (1971). The effect of photoperiod on daily rhythms of oxygen consumption in the tropical toad, Bufo marinus. Z. Vergl. Physiol. 75, 367 - 382.
- Hutchison, V. and L. Turney. (1975). Glucose and lactate concentrations during activity in the leopard frog, Rana pipiens. J. Comp. Physiol. 99, 287 - 295.
- Jacobson, E. and W. Whitford. (1970). The effects of acclimation on physiological responses to temperature in the snakes, Thamnophis proximus and Natrix rhombifera. Comp. Biochem. Physiol. 35, 439 - 449.
- Kardong, K. (1972a). Morphology of the respiratory system and its musculature in different snake genera (part I) Crotalus and Elaphe. Gegenbaur's Morph. Jahrb. 117, 285 - 302.

- Kardong, K. (1972b). Morphology of the respiratory system and its musculature in different snake genera (part II) Charina bottae. Gegenbauer's Morph. Jahrb. 117, 364 - 376.
- Kemp, A. and A. Van Heijningen. (1954). A colorimetric micro-method for the determination of glycogen in tissues. Biochem. J. 56, 646 - 648.
- Knuttgen, H. (1971). Lactic acid and oxygen debt: an introduction. In Muscle Metabolism during Exercise. (ed. B. Pernow and B. Saltin). New York and London: Plenum Press.
- Kramer, K., G. Driessen and H. Brechtelsbauer. (1971). Lactate elimination and O_2 consumption of the liver in narcotized dogs. Pflügers Arch. 330, 195 - 205.
- Masoro, E. and P. Siegel. (1971). Acid-Base Regulation: its Physiology and Pathophysiology. Philadelphia: W. B. Saunders Co.
- McDonald, H. (1959). Respiratory functions of the ophidian air sac. Herpetologica 15: 193 - 198.
- Minaire, Y. (1973). Origine et destinée du lactate plasmatique. J. Physiol., Paris. 66, 229 -257.
- Moberly, W. (1968). The metabolic responses of the common iguana, Iguana iguana, to activity under restraint. Comp. Biochem. Physiol. 27, 1 - 20.
- Newsholme, E. and C. Start. (1973). Regulation in Metabolism. London: John Wiley & Sons.

- Preston, W. (1970). The comparative ecology of two water snakes, Natrix rhombifera and Natrix erythrogaster, in Oklahoma. Ph.D. Dissertation deposited in the library of the University of Oklahoma, Norman, Oklahoma.
- Prosser, C. (1973). Temperature. In Comparative Animal Physiology. (ed. C. Prosser). W.B. Saunders Co.
- Roberts, L. (1968). Oxygen consumption in the lizard Uta stansburiana. Ecology 49, 809 - 819.
- Turney, L. and V. Hutchison. (1974). Metabolic scope, oxygen debt, and diurnal oxygen consumption of the leopard frog, Rana pipiens. Comp. Biochem. Physiol. 49A, 583 - 601.

Table 1. Metabolic rate and routine metabolic scope (RMS) of the diamond-back water snake, Natrix rhombifera. Gas exchange rates are presented in $\text{ul g}^{-1} \text{hr}^{-1}$ STPD ($\bar{x} \pm \text{SE}$, N = animal hours).

Acclimation Temperature	$\bar{x} - 24 \text{ hrs}$			Q_{10}	$\dot{V}_{O_2} - \bar{x} \text{ Min}$	Time (CDT)	$\dot{V}_{O_2} - \text{Max}$	Time (CDT)	RMS	Increase
	$\dot{V}_{O_2}$	$\dot{V}_{CO_2}$	R							
15°C	29.5 (± 1.64 , 219)	23.3	0.79	5.28	18.7 (± 2.87 , 28)	2200-2400	45.6 (± 9.26 , 10)	1200	26.9	x 2.44
25°C	155.9 (± 4.77 , 240)	124.8	0.80		133.5 (± 9.99 , 20)	0100-0200	211.1 (± 37.75 , 10)	0800	77.6	x 1.58
35°C	356.2 (± 10.98 , 235)	256.4	0.72	2.28	277.3 (± 25.46 , 18)	1100-1200	485.6 (± 54.10 , 10)	0700	208.4	x 1.75

Table 2. Energy budget for 10 minutes forced activity in the diamond-back water snake, *Natrix rhombifera*.

Acclimation		Yield							Recovery	Oxygen Debt		
Temperature	Hour	N	Source	$\mu\text{l O}_2 \text{ g}^{-1}$	mg g^{-1} lactate	$\mu\text{M ATP g}^{-1}$	percent	cal g^{-1}	time	$\mu\text{l O}_2 \text{ g}^{-1}$	$\mu\text{M ATP g}^{-1}$	cal g^{-1}
(°C)	(CDT)			($\bar{x} \pm \text{SE}$)			contribution		(min $\pm \text{SE}$)	($\bar{x} \pm \text{SE}$)		
15°C	1400	10	Aerobic	4.9 (± 1.42)	-	1.42	8.0	0.03	153 (± 23.8)	100.9 (± 26.8)	27.04	0.49
		5	Anaerobic	-	0.97	<u>16.20</u>	92.0	<u>0.25</u>				
			Total			17.62		0.28				
25°C	1400	10	Aerobic	18.8 (± 2.05)	-	5.45	15.3	0.10	138 (± 19.5)	175.8 (± 20.5)	47.11	0.85
		5	Anaerobic	-	1.81	<u>30.23</u>	84.7	<u>0.47</u>				
			Total			35.68		0.57				
25°C	2200	10	Aerobic	16.6 (± 2.42)	-	4.81	21.4	0.08	118 (± 18.8)	182.3 (± 28.8)	48.86	0.88
		5	Anaerobic	-	1.06	<u>17.70</u>	78.6	<u>0.28</u>				
			Total			22.51		0.36				
35°C	1400	10	Aerobic	18.4 (± 3.81)	-	5.34	15.6	0.09	59 (± 11.0)	108.0 (± 25.1)	28.95	0.52
		5	Anaerobic	-	1.73	<u>28.89</u>	84.4	<u>0.45</u>				
			Total			34.23		0.54				

Table 3. Tissue metabolites (mean $\pm$ SE, N = number of animals) in Natrix rhombifera: controls and immediately after 10 min forced activity at 25°C. All values are presented as mg g⁻¹ tissue wet weight or mg ml⁻¹ plasma.

Time (CDT)	Experiment	Blood		Liver		Muscle		Total Body
		Glucose	Lactate	Glycogen	Lactate	Glycogen	Lactate	Lactate
1400	control	0.15 ($\pm$ 0.02,14)	0.09 ($\pm$ 0.09,10)	55.45 ($\pm$ 10.3,12)	0.30 ($\pm$ 0.03,8)	10.46 ($\pm$ 1.41,10)	1.36 ($\pm$ 0.13,8)	0.34 ($\pm$ 0.02,5)
	activity	0.702 ($\pm$ 0.13,5)	1.08 ($\pm$ 0.08,7)	44.23 ($\pm$ 7.6,7)	0.93 ($\pm$ 0.58,7)	3.93 ($\pm$ 0.94,7)	3.03 ($\pm$ 0.12,6)	2.15 ($\pm$ 0.06,5)
2200	control	0.23 ($\pm$ 0.03,8)	0.07 ($\pm$ 0.004,8)	51.04 ($\pm$ 8.8,12)	0.32 ($\pm$ 0.03,9)	11.41 ($\pm$ 1.98,9)	1.18 ($\pm$ 0.10,9)	0.40 ($\pm$ 0.04,5)
	activity	0.52 ($\pm$ 0.04,5)	1.15 ($\pm$ 0.07,7)	42.74 ($\pm$ 11.4,7)	1.13 ($\pm$ 0.12,7)	8.32 ($\pm$ 1.53,7)	2.71 ($\pm$ 0.24,7)	1.46 ($\pm$ 0.18,5)

FIGURE LEGENDS

Fig. 1. Diurnal cycles of oxygen consumption (half-shaded dots and dotted line) and routine activity (closed dots and solid line) in the diamond-back water snake, Natrix rhombifera, at three acclimation temperatures. Hatched areas beneath graphs indicate scotophase. Each point is the mean of 5 - 10 animals (8 - 12 determinations).

Fig. 2. The effect of temperature and hour of measurement on metabolic scope for activity in N. rhombifera. Circle is routine $\dot{V}O_2$ at hour indicated, square is maximum $\dot{V}O_2$ during activity and triangle is maximum $\dot{V}O_2$ during recovery. Vertical line is one standard error of the mean. Brackets indicate aerobic scope in $ul\ O_2\ g^{-1}\ hr^{-1}$ STPD and parentheses indicate the index of metabolic expansibility for each temperature and hour.

Fig. 3. Tissue metabolite levels in N. rhombifera during recovery from 10 min forced activity at 25°C, 1400 hrs. C = control value; I = immediately after activity, numbers represent hours of recovery. Each point is the mean from 5 - 14 animals except for liver glycogen, the last two data of which are based on three animals.

Fig. 4. Total energy cost of 10 min forced activity in N. rhombifera at temperatures and hours indicated on the abscissa. Shaded areas represent aerobic and anaerobic capacities and open areas the oxygen debt. Numbers above each bar are the caloric equivalent ($cal\ g^{-1}$) of the total energy expenditure.

Fig. 1

Fig. 2

Fig 3

Fig. 4

APPENDIX I

Diurnal variation in oxygen consumption in the diamond-back water snake, Natrix rhombifera. Gas volumes are presented as $\mu\text{l g}^{-1} \text{ hr}^{-1}$ STPD. Data correspond to those presented in Fig. 1. Each table is followed by the results of a two-way analysis of variance performed on the data. Following all tables are the results of the Duncan multiple range test. To determine if any two hours are significantly different find the point of intersection of the two hours. A blank indicates no difference, 5 indicates $P < 0.05$, 1 indicates $P < 0.01$.

15°C Acclimation

Hour (CDT)	Mean $\dot{V}_{O_2}$	N	Standard Error
0100	21.7	9	6.40
0200	27.4	10	4.96
0300	32.0	9	7.43
0400	26.8	7	6.78
0500	23.5	9	5.26
0600	39.6	10	8.57
0700	25.5	9	7.07
0800	26.9	10	5.92
0900	31.2	9	9.34
1000	34.3	9	11.07
1100	37.2	10	9.93
1200	45.6	10	9.68
1300	36.2	9	9.96
1400	36.9	9	10.09
1500	32.0	9	10.24
1600	34.7	8	12.59
1700	31.6	10	10.05
1800	23.2	10	7.46
1900	22.4	9	5.11
2000	26.8	8	4.12
2100	36.1	8	9.49
2200	16.6	9	4.03
2300	22.1	10	5.72
2400	17.5	9	4.88

Source of Variation	DF	Mean Square	F	P
Between snakes	9	6913.012	24.16	0.001
Between hours	23	512.628	1.79	0.05
Error	207	286.080		

25°C Acclimation

Hour (CDT)	Mean $\dot{V}_{O_2}$	N	Standard Error
0100	138.4	10	15.39
0200	128.6	10	12.59
0300	154.4	10	27.40
0400	149.6	10	25.66
0500	178.1	10	33.43
0600	170.1	10	28.75
0700	181.7	10	33.33
0800	211.1	10	39.34
0900	169.3	10	32.51
1000	143.6	10	16.95
1100	139.5	10	14.15
1200	161.9	10	28.03
1300	149.7	10	19.72
1400	154.7	10	19.87
1500	138.1	10	15.26
1600	140.7	10	17.93
1700	148.6	10	17.88
1800	163.4	10	27.98
1900	159.5	10	26.90
2000	152.7	10	19.24
2100	138.4	10	17.77
2200	166.5	10	20.96
2300	152.9	10	16.09
2400	151.4	10	19.61

Source of Variation	DF	Mean Square	F	P
Between snakes	9	69427.625	28.27	0.001
Between hours	23	2927.000	1.19	ns
Error	207	2456.094		

35°C Acclimation

Hour (CDT)	Mean $\dot{V}O_2$	N	Standard Error
0100	360.6	10	66.34
0200	307.6	10	30.63
0300	327.8	10	32.01
0400	331.3	10	27.35
0500	345.7	10	30.73
0600	415.9	10	47.39
0700	485.6	10	57.26
0800	447.7	10	43.53
0900	404.4	10	32.41
1000	361.5	10	41.04
1100	282.3	9	32.45
1200	272.2	9	39.28
1300	383.0	9	78.02
1400	327.0	9	75.73
1500	345.6	9	77.65
1600	420.0	10	92.75
1700	382.1	10	70.28
1800	348.8	10	58.60
1900	345.8	10	57.22
2000	332.4	10	47.25
2100	310.0	10	44.85
2200	308.6	10	33.64
2300	335.9	10	46.03
2400	349.1	10	69.91

Source of Variation	DF	Mean Square	F	P
Between Snakes	8	210709.500	18.97	0.001
Between hours	23	28021.250	2.52	0.005
Error	184	11109.590		

15°C Acclimation

	2400	2300	2200	2100	2000	1900	1800	1700	1600	1500	1400	1300	1200	1100	1000	0900	0800	0700	0600	0500	0400	0300	0200	0100
0100													1											
0200																								
0300																								
0400																								
0500																								
0600	5		5										5											
0700													5											
0800																								
0900																								
1000																								
1100	5		5																					
1200	1	5	1			5	5																	
1300																								
1400			5																					
1500																								
1600																								
1700																								
1800																								
1900																								
2000																								
2100																								
2200																								
2300																								
2400																								

At 25°C the only significant differences are that oxygen consumption at 0800 is significantly greater than that at 0100, 0200, 1500 and 2100 at the $P < 0.01$ level, and greater than 0300, 0400, 1000, 1100, 1300, 1400, 1600, 1700, 2000, 2300 and 2400 at the $P < 0.05$ level.

35°C Acclimation

	2400	2300	2200	2100	2000	1900	1800	1700	1600	1500	1400	1300	1200	1100	1000	0900	0800	0700	0600	0500	0400	0300	0200	0100
0100																		5						
0200																	5	1						
0300																	5	1						
0400																	5	1						
0500																		5						
0600													5	5										
0700	5	1	1	1	1	5	5			5	1		1	1	5									
0800			5	5	5						5		1	1										
0900													5	5										
1000																								
1100									5															
1200									1															
1300																								
1400																								
1500																								
1600																								
1700																								
1800																								
1900																								
2000																								
2100																								
2200																								
2300																								
2400																								

APPENDIX II

Diurnal variation in routine activity in the diamond-back water snake, Natrix rhombifera. Values presented are percent of total activity and correspond to those presented in Fig. 1. The table of values is followed by the results of the analyses of variance performed on the arcsin transformed data and the results of the Duncan multiple range test, also performed on the arcsin transformed data.

Acclimation temperature						
Time (GDT)	15°C N = 12		25°C N = 11		35°C N = 10	
	Mean %	Standard error	Mean %	Standard error	Mean %	Standard Error
0100	0.0	0.0	0.9	0.45	0.8	0.35
0200	0.0	0.0	1.1	0.51	1.8	0.63
0300	0.0	0.0	2.1	1.30	3.4	0.97
0400	0.0	0.0	0.6	0.46	3.5	0.73
0500	0.8	0.83	0.7	0.35	4.8	1.40
0600	0.3	0.35	0.7	0.36	12.1	1.68
0700	9.4	4.70	1.2	0.63	20.0	2.36
0800	8.3	3.64	1.1	0.50	16.3	1.80
0900	7.9	3.67	1.8	0.78	15.0	2.10
1000	5.2	2.01	3.2	0.92	8.5	1.27
1100	7.0	3.58	3.8	1.49	2.9	0.70
1200	9.7	4.48	7.3	1.67	0.9	0.63
1300	2.8	1.92	7.8	3.36	0.02	0.02
1400	2.6	1.45	15.5	4.02	0.04	0.04
1500	2.1	1.48	10.3	3.46	0.1	0.07
1600	3.6	2.12	2.0	1.06	0.5	0.25
1700	7.2	4.76	5.1	1.35	0.3	0.19
1800	4.4	2.42	6.7	2.10	0.1	0.08
1900	24.1	4.59	10.7	3.34	3.7	1.59
2000	4.3	2.33	3.1	1.32	1.1	0.37
2100	0.2	0.15	1.9	0.78	1.3	0.44
2200	0.0	0.0	2.8	0.92	0.7	0.25
2300	0.2	0.15	6.3	3.93	1.3	0.35
2400	0.0	0.0	1.9	0.59	0.8	0.34

15°C Acclimation

Source of Variation	DF	Mean Square	F	P
Between snakes	11	22.473	0.25	ns
Between hours	23	445.918	4.98	0.001
Error	253	8934.688		

25°C Acclimation

Source of Variation	DF	Mean Square	F	P
Between snakes	10	21.876	0.31	ns
Between hours	23	242.998	3.44	0.001
Error	230	16224.988		

35°C Acclimation

Source of Variation	DF	Mean Square	F	P
Between snakes	9	34.169	2.16	0.01
Between hours	23	611.645	38.66	0.001
Error	207	15.823		

15°C Acclimation

	2400	2300	2200	2100	2000	1900	1800	1700	1600	1500	1400	1300	1200	1100	1000	0900	0800	0700	0600	0500	0400	0300	0200	0100
0100						1							1			5	5	1						
0200						1							1			5	5	1						
0300						1							1			5	5	1						
0400						1							1			5	5	1						
0500						1							5			5	5	5						
0600						1							5			5	5	5						
0700	1	5	1	1		1										5	5	5						
0800	5	5	5	5		1																		
0900	5	5	5	5		1																		
1000						1																		
1100						1																		
1200	1	5	1	5		1																		
1300						1																		
1400						1																		
1500						1																		
1600						1																		
1700						1																		
1800						1																		
1900	1	1	1	1	1																			
2000																								
2100																								
2200																								
2300																								
2400																								

25°C Acclimation

	2400	2300	2200	2100	2000	1900	1800	1700	1600	1500	1400	1300	1200	1100	1000	0900	0800	0700	0600	0500	0400	0300	0200	0100
0100						1	5			1	1	5	5											
0200						1				5	1	5	5											
0300						5				5	1													
0400						1	5	5		1	1	5	1											
0500						1	5			1	1	5	5											
0600						1	5			1	1	5	1											
0700						1				5	1	5	5											
0800						1				5	1	5	5											
0900						5				5	1													
1000											1													
1100											1													
1200																								
1300																								
1400	1	1	1	1	1			5	1															
1500				5																				
1600																								
1700																								
1800																								
1900	5		5	5	5																			
2000																								
2100																								
2200																								
2300																								
2400																								

35°C Acclimation

	2400	2300	2200	2100	2000	1900	1800	1700	1600	1500	1400	1300	1200	1100	1000	0900	0800	0700	0600	0500	0400	0300	0200	0100
0100															1	1	1	1	1		1	1		
0200							5	5		5	1	1			1	1	1	1	1	1	5			
0300	1		1		5		1	1	1	1	1	1	1		1	1	1	1	1					
0400	1	5	1	5	5		1	1	1	1	1	1	1		1	1	1	1	1					
0500	1	1	1	1	1		1	1	1	1	1	1	1		1	1	1	1	1					
0600	1	1	1	1	1	1	1	1	1	1	1	1	1	1				1						
0700	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	5								
0800	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1									
0900	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1									
1000	1	1	1	1	1	1	1	1	1	1	1	1	1	1										
1100	5		5		5		1	1	1	1	1	1	1											
1200						1																		
1300		5		5	5	1																		
1400		5		5	5	1																		
1500		5		5		1																		
1600						1																		
1700						1																		
1800		5		5		1																		
1900	5		5		5																			
2000																								
2100																								
2200																								
2300																								
2400																								

APPENDIX III

Oxygen consumption volumes (V_{O_2}) and maximum sustained rates ($\dot{V}_{O_2}$) during activity and recovery from 10 minutes forced activity in the diamond-back water snake, Natrix rhombifera. Gas volumes are presented as $\mu\text{l } O_2 \text{ g}^{-1}$ and rates as $\mu\text{l } O_2 \text{ g}^{-1} \text{ hr}^{-1}$ STPD, mean $\pm$ Standard error, N = animal hours.

Acclimation Temperature and time of measurement	Mean Weight (g)	V_{O_2} activity	$\dot{V}_{O_2}$ activity	V_{O_2} recovery	$\dot{V}_{O_2}$ recovery	Recovery time (min)
15°C, 1400 hrs	694.6	4.9 ($\pm 1.42, 10$)	108.5 ($\pm 16.39, 10$)	100.9 ($\pm 29.94, 9$)	143.3 ($\pm 20.45, 9$)	153.2 ($\pm 26.6, 9$)
50 25°C, 1400 hrs	642.9	18.8 ($\pm 2.16, 10$)	317.7 ($\pm 20.54, 10$)	175.8 ($\pm 20.47, 10$)	382.2 ($\pm 32.23, 10$)	137.8 ($\pm 20.6, 10$)
25°C, 2200 hrs	590.0	16.6 ($\pm 2.54, 11$)	309.9 ($\pm 41.67, 11$)	182.3 ($\pm 30.23, 11$)	361.8 ($\pm 36.22, 11$)	117.8 ($\pm 19.7, 11$)
35°C, 1400 hrs	210.9	18.4 ($\pm 3.81, 10$)	480.7 ($\pm 94.76, 10$)	108.0 ($\pm 25.11, 9$)	493.2 ($\pm 93.92, 9$)	58.8 ($\pm 11.6, 9$)

APPENDIX IV

Metabolite concentrations before, immediately after and during recovery from 10 min forced activity at 25°C, 1400 hrs in the diamond-back water snake, Natrix rhombifera. For each metabolite, the data summary is followed by the results of the analysis of variance and the Duncan multiple range test performed on the data. Significance levels are presented as: blank = no significance; 5 = $P < 0.05$; 1 = $P < 0.01$.

Blood Glucose

Time (CDT)	Mean (mg ml ⁻¹)	Sample size	Standard Error
Control (C)	0.15	14	0.02
Immediately after (I)	0.70	5	0.12
1 hour of recovery	0.79	5	0.10
3 hours of recovery	1.16	6	0.10
5 hours of recovery	1.25	6	0.10
7 hours of recovery	1.11	5	0.17

Source of variation	DF	Mean Square	F	P
Between samples	5	16476.86	36.81	0.001
Within samples	34	447.57		
Total	39			

Time	C	I	1	3	5	7
7	1	1	5			
5	1	1	1			
3	1	1	5			
1	1					
I	1					
C						

Blood Lactate

Time (CDT)	Mean (mg ml ⁻¹)	Sample size	Standard error
Control (C)	0.09	10	0.02
Immediately after (I)	1.08	7	0.08
1 hour of recovery	1.46	5	0.06
3 hours of recovery	0.99	6	0.10
5 hours of recovery	0.48	6	0.08
7 hours of recovery	0.05	5	0.02

Source of variation	DF	Mean Square	F	P
Between samples	5	19381.13	71.58	0.001
Within samples	32	270.75		
Total	37			

Time	C	I	1	3	5	7
7		1	1	1	1	
5	1	1	1	1		
3	1		1			
1	1	1				
I	1					
C						

Liver Glycogen

Time (CDT)	Mean (mg g ⁻¹)	Sample size	Standard error
Control (C)	55.45	12	7.31
Immediately after (I)	44.23	7	7.01
1 hour of recovery	44.62	5	5.86
3 hours of recovery	46.23	3	10.92
5 hours of recovery	x*	x	x
7 hours of recovery	52.53	3	10.89

* no data for this hour

Source of variation	DF	Mean Square	F	P
Between samples	4	1608138.00	0.36	ns
Within samples	24	4451707.00		
Total	28			

The Duncan multiple range test showed no hours to be significantly different from any others.

Liver Lactate

Time (CDT)	Mean (mg g ⁻¹)	Sample size	Standard error
Control (C)	0.30	8	0.03
Immediately after (I)	0.93	7	0.06
1 hour of recovery	1.20	5	0.11
3 hours of recovery	0.81	6	0.07
5 hours of recovery	0.46	6	0.08
7 hours of recovery	0.34	5	0.04

Source of variation	DF	Mean Square	F	P
Between samples	5	7181.59	29.40	0.001
Within samples	30	244.24		
Total	35			

Time	C	I	1	3	5	7
7		1	1	1		
5		1	1	1		
3	1		1			
1	1	1				
I	1					
C						

Muscle Glycogen

Time (CDT)	Mean (mg g ⁻¹)	Sample size	Standard error
Control (C)	10.5	10	1.41
Immediately after (I)	3.9	7	0.94
1 hour of recovery	6.8	5	1.70
3 hours of recovery	8.4	5	2.21
5 hours of recovery	14.8	6	1.51
7 hours of recovery	13.5	5	2.56

Source of variation	DF	Mean Square	F	P
Between samples	5	990760.19	5.61	0.001
Within samples	31	176559.69		
Total	36			

Time	C	I	1	3	5	7
7		1	5	5		
5		1	1	5		
3						
1						
I	5					
C						

Muscle Lactate

Time (CDT)	Mean (mg g ⁻¹)	Sample size	Standard error
Control (C)	1.36	8	0.13
Immediately after (I)	3.10	6	0.11
1 hour of recovery	2.48	5	0.21
3 hours of recovery	1.43	6	0.16
5 hours of recovery	1.20	6	0.12
7 hours of recovery	1.34	5	0.13

Source of variation	DF	Mean Square	F	P
Between samples	5	38194.16	35.52	0.001
Within samples	29	1075.44		
Total	34			

Time	C	I	1	3	5	7
7		1	1			
5		1	1			
3		1	1			
1	1	5				
I	1					
C						

Total Body Lactate

Time (CDT)	Mean (mg g ⁻¹)	Sample size	Standard error
Control (C)	0.34	5	0.02
Immediately after (I)	2.15	5	0.06
1 hour of recovery	1.29	5	0.13
3 hours of recovery	0.87	5	0.14
5 hours of recovery	0.44	5	0.07

Source of variation	DF	Mean Square	F	P
Between samples	4	27084.20	59.71	0.001
Within samples	20	453.58		
Total	24			

Time	C	I	1	3	5
5		1	1	1	
3	1	1	1		
1	1	1			
I	1				
C					

APPENDIX V

Results of the regression analysis of "standard" metabolic rate vs acclimation temperature in the diamond-back water snake, N. rhombifera.

Acclimation Temperature	N	Mean $\dot{V}O_2$ ($\mu\text{l g}^{-1} \text{ hr}^{-1}$)	Standard error
15°C	28	18.7	2.87
25°C	20	133.5	9.99
35°C	18	277.3	25.46

Regression equation : $\dot{V}O_2 = 12.11044 T_{\text{accl}} - 166.5509$

95% confidence limits of the slope are: 3.7655 and 20.4553

Analysis of variance

Source of variation	DF	Mean Square	F	P
Groups	2	327172.4375	100.18	0.001
linear	1	652427.3125	340.23	0.05
deviations	1	1917.6250	0.59	ns
Error	63	3265.8569		
Total	65			