THE UNIVERSITY OF OKLAHOMA
GRADUATE COLLEGE

DIFFERENCES IN NITRATE REDUCTASE ACTIVITY BETWEEN
SPECIES OF DIFFERENT STAGES IN OLD FIELD SUCCESSION

A DISSERTATION
SUBMITTED TO THE GRADUATE FACULTY
in partial fulfillment of the requirements for the
degree of
DOCTOR OF PHILOSOPHY

BY
JEFFREY LYNN SMITH
Norman, Oklahoma
1982

DIFFERENCES IN NITRATE REDUCTASE ACTIVITY BETWEEN
SPECIES OF DIFFERENT STAGES IN OLD FIELD SUCCESSION

APPROVED BY

Elroy L. Rice
Herbert F. Jones
Thomas R. Cole
Att. Milby
Lion S. Glicks

DISSERTATION COMMITTEE

ACKNOWLEDGEMENTS

I wish to thank Dr. Elroy Rice for his help and guidance during this study, and to Drs. Leonard Beevers, James Estes, T.H. Milby, and Leon Ciereszko for their critical reading and comments on the writing. Appreciation is extended to Drs. Linda Wallace, Charles Daghljan, William Hurkman, and John Fletcher for their interest and helpful suggestions. I would especially like to thank Jim Spruell for his friendship and encouragement. I thank my parents and family for their love and understanding during this time of study. My deepest gratitude and love is to Deb Kuriger; whose assistance, support and love made this project possible.

TABLE OF CONTENTS

	Page
LIST OF TABLES.	v
LIST OF ILLUSTRATIONS	vi
TEXT	
I. SUMMARY	1
II. INTRODUCTION.	1
III. METHODS AND MATERIALS	5
IV. RESULTS	10
V. DISCUSSION.	13
VI. REFERENCES.	29

LIST OF TABLES

TABLE		Page
1.	Variations in mean Nitrate Reductase Activity in <u>Ambrosia trifida</u>	19
2.	Optimal conditions for <u>in vivo</u> Nitrate Reductase Activity.	20
3.	Nitrate Reductase Activity of leaf-high species	22
4.	Nitrate Reductase Activity of root-high species	23
5.	Nitrate Reductase Activity of species grown on different nitrogen treatments.	24
6.	Shoot dry weights of species grown on different nitrogen treatments . . .	26

LIST OF ILLUSTRATIONS

FIGURE	Page
1. Nitrate Reductase Activity of eight week old <u>Bromus japonicus</u> roots grown on nitrate or ammonium only.	28

FORMAT STYLE

This work will be submitted for publication in *Oecologia*. Format and style follow instructions by the journal.

Summary

The ratio of nitrate/ammonium in the soil decreases from a high level in the early stages of old field succession to a low level in the climax in the tall grass prairie region of central Oklahoma. Characteristic plant species from different stages of succession were grown in culture solutions where the total nitrogen content was held constant, but the form varied from nitrate only to ammonium only. The nitrate reductase activity of leaves and roots was measured in vivo. The results indicated that four pioneer species had relatively high levels of enzyme activity while four climax species had relatively low levels. Nitrate reductase activity was consistently higher in either the leaves or the roots of a species than in the other organ regardless of the nitrogen treatment. Enzyme activity generally decreased with decreasing nitrate concentrations, but seven cases were found where there were no significant differences in enzyme activity between any of the nitrogen treatments. All eight species had their highest shoot dry weights on the ammonium only treatment.

Introduction

Booth (1941) reported that secondary succession in abandoned fields in the tall grass prairie region of southeastern Kansas and central Oklahoma has four prominent stages of revegetation: (1) the pioneer weed stage which lasts 2-3 years, (2) the annual grass stage

lasting 9-13 years, (3) the bunch grass stage lasting 40 years or more, and (4) the climax tall grass prairie. The change in species composition during this succession has been of particular interest. Some species such as Helianthus annuus and Ambrosia trifida are found only in the recently abandoned fields, while others such as Sorghastrum nutans and Panicum virgatum are found only in the climax prairie. The mechanisms involved in the changes in composition of species during old field succession in central Oklahoma have been extensively investigated. The seed dispersal and mineral nutrition requirements of the species change during succession (Rice et al., 1960). Allelopathic effects of the pioneer weeds are important in the rapid disappearance of the pioneer weed stage (Abdul-Wahab and Rice, 1967; Neill and Rice, 1971; Parenti and Rice, 1969; Rice, 1976; Wilson and Rice, 1968) and also in the early inhibition of nitrogen fixation (Kapustka and Rice, 1976; Rice, 1964, Rice, 1968). Aristida oligantha is allelopathic to the nitrogen-fixing bacteria and species from several stages are allelopathic to nitrogen-fixing bluegreen algae (Parks and Rice, 1969). Species from several stages are also inhibitory to the nitrifiers (Rice and Pancholy, 1972; 1973; 1974) with the climax species being the most inhibitory.

Rice and Pancholy (1972) found that the concentration of ammonium nitrogen increased from a low value in the pioneer weed stage to a high value in the climax, whereas the concentration of nitrate nitrogen decreased from a high value in the pioneer weed stage to a very low value in the climax. They also found that the number of nitrifiers was high in the pioneer weed stage but was low in the climax. They hypothesized that the soil under the climax vegetation was low in nitrate due to an

inhibition of nitrification by the climax plant species. This was thought to be a logical successional development of ecosystems as a result of the increased conservation of nitrogen and the reduced energetic cost to incorporate ammonium instead of nitrate.

The result of the inhibition of nitrification would be a shift in form of nitrogen during the succession. Pioneer species would have mostly nitrate available to them while the climax species would have mainly ammonium. The utilization of a particular nitrogen form by a species has been reported previously. Some species of conifers grow best on nitrate (Krajina et al., 1973), others on ammonium (Bigg and Daniel, 1978; McFee and Stone, 1968), and still others on a mixture of both forms (Addoms, 1937; Van Den Driessche, 1971). Brassica hirta grows best on nitrate alone (Kirkby, 1968) but Vaccinium macrocarpon is unable to survive on nitrate and requires ammonium (Greidanus et al., 1972). Other species are reported to be able to use a mixture of the two forms (Haynes and Goh, 1978; Moris and Giddens, 1963; Schrader et al., 1972). Gigon and Rorison (1972) reported that some ecologically distinct species respond best to the nitrogen form which is available in their particular habitat. Calcifuge species grew best on ammonium while calcareous species grew best on nitrate. They also found that populations of Scabiosa columbaria which grows on both soil types had become adapted to the nitrogen form of the habitat. Seedlings grown from seeds of the calcifuge population grew best on ammonium whereas seedlings grown from seeds of the calcareous population grew best on nitrate. Reports of a change in preference in the form of nitrogen across several successional stages are relatively few. However, Wiltshire (1973) reported that

climax grasses showed a greater preference for ammonium than did earlier seral species or crop cultivars.

Species that are adapted to a particular successional stage must be adapted to the nitrogen form available to them. In particular, the enzymatic systems for absorbing and incorporating nitrogen should reflect the form of nitrogen used. Nitrate reductase (henceforth labeled NR) is the enzyme involved in the first step of the reduction of nitrate into a form that a plant can use (Beevers and Hageman, 1980). This enzyme controls the ability of a plant to use nitrate as a nitrogen source. Species that use mostly nitrate as their nitrogen source should have relatively high levels of nitrate reductase activity (henceforth labeled NRA). Species that use mostly ammonium as their nitrogen source should have relatively low levels of NRA.

Havill et al. (1974) reported that calcareous species showed high levels of NRA. Calcifuge species (mostly members of the Ericaceae) showed no NRA in the field. They were able to demonstrate low levels of NRA in calcifuge species under laboratory conditions, but only after feeding the plants nitrate. Studies that correlate the NRA of a species with its successional stage are few. Bate and Heelas (1975) reported that Sporobolus pyramidalis, a pioneer grass, had relatively high levels of NRA while Hyparrhenia filipendula, a climax grass, had relatively low levels of NRA. Franz and Haines (1977) reported that the NRA of vascular plants decreased during an old field to forest succession in South Carolina.

To date, no studies have been reported correlating the NRA of a species with its successional stage in the tall grass prairie region.

I hypothesized that the pioneer species in the tall grass prairie region should be adapted to nitrate as their primary nitrogen source and have relatively high levels of NRA. Climax species should be adapted to ammonium as their primary nitrogen source and should have relatively low levels of NRA. One goal of this study, therefore, was to determine if the NRA of a species depends on the successional stage in which the species occurs. Another goal was to determine the growth responses of species from the different successional stages to various ratios of nitrate to ammonium.

Methods and Materials

1. Growth of the plants

Eight species were used in the study. Seeds of two species from the pioneer weed stage (Ambrosia trifida L. and Helianthus annuus L.), one species from the annual grass stage (Aristida oligantha Michx.), and three climax species (Panicum virgatum L., Schizachyrium scoparium Nash., and Sorghastrum nutans (L.) Nash.) were collected in 1979 in the Norman, Oklahoma area. Representative specimens for each seed source are deposited in the Bebb Herbarium at the University of Oklahoma (OKL). Seeds for the pioneer species, Bromus japonicus Thunb., had been previously collected in the spring of 1978. Seeds collected locally for Andropogon gerardii Vitman failed to germinate, therefore

seeds from a non-local source were substituted in order that all four dominant grass species of the climax prairie could be tested. Seeds were stratified if necessary and germinated in plastic petri dishes on one sheet of Whatman No. 1 filter paper. The seedlings were transplanted into vermiculite in glazed ceramic pots, 12 cm in diameter and 20 cm deep. Planting densities of the seedlings were: 1 plant/pot (Helianthus annuus), 2 plants/pot (Ambrosia trifida), or 3 plants/pot (all grass species). Seedlings that died within the first 10 days after transplanting were replaced. The pots were then placed in a growth chamber immediately after transplanting. The growing conditions were: 28°C day/20°C night with a 16 hr light/8 hr dark photoperiod. Ambrosia trifida and Helianthus annuus were grown for ten weeks and all the grass species were grown for eight weeks before the determination of NRA.

The nitrogen treatments used were based on a modified Evan's minus-N solution (Moore, 1974). The final concentrations of the mineral salts used were: CaSO_4 - 6 mM, MgSO_4 - 2 mM, K_2SO_4 - 1.6 mM, KH_2PO_4 - 0.17 mM, K_2HPO_4 - 0.83 mM, HBO_3 - 23 μM , MnCl_2 - 5 μM , ZnSO_4 - 0.76 μM , CuSO_4 - 0.32 μM , CoSO_4 - 0.35 μM , and H_2MoO_4 - 0.28 μM . Iron was added at a 50 μM concentration as iron chelated with ethylenediaminetetraacetic acid. Nitrogen was given as $\text{Ca}(\text{NO}_3)_2$ or NH_4Cl with a total nitrate and/or ammonium concentration of 0.33 mM. Calcium chloride was added as necessary to balance the Ca^{+2} content. The nitrogen form was varied so that a nitrogen treatment contained either nitrate only, a 2:1 ratio of nitrate/ammonium, a 1:2 ratio of nitrate/ammonium, or ammonium only. There were 10 replicate pots per

nitrogen treatment. I found in preliminary experiments that ammonium concentrations higher than 0.33 mM caused toxicity. Consequently, the nitrate/ammonium concentrations were maintained below this level. The total nitrogen concentration used here was approximately twice that reported in the soil by Rice and Pancholy (1972), but it was still in an ecologically meaningful range. No apparent ammonium toxicity was observed at the 0.33 mM nitrogen content. Maintaining the pH of the nutrient solutions is important (Barker et al., 1966; Karim and Vlamis, 1962) and the pH of the solutions was adjusted to 6.0 by the addition of NaOH or HCl before the nutrient solution was given to the plants. Each pot was given 100 ml of a nutrient solution three times a week for the first four weeks after transplanting, then five times a week for the remainder of the growing period. Distilled water was given as necessary to prevent desiccation. The pots were flooded on two consecutive days per week to prevent pH shifting and accumulation of the mineral salts. Spot checks of the pH of pot exudates indicated that the pH was 6.0 ± 0.1 . N-serve 24E (2-chloro-6-(trichloromethyl)pyridine, Dow Chemical Co.) was added at a rate of 0.1 ml/l nutrient solution once per week to inhibit the oxidation of ammonium to nitrate by nitrifying bacteria (Goring, 1962a; 1962b).

2. Enzyme assay

A modified in vivo assay system for NRA was used (Klepper et al., 1971; Jaworski, 1971). A series of preliminary experiments were run

for each species on the roots and leaves of plants grown in nitrate only. These experiments were designed to maximize NRA by finding the optimal combinations of phosphate buffer, substrate (KNO_3), incubation time length, and n-propanol for the incubation. The pH of the incubation medium was held at 7.5 (Kannangara and Woolhouse, 1967). The optimal combination of variables determined for the leaves and roots of each species was then used in the incubation for determining the NRA of that organ for all nitrogen treatments. The incubations were conducted two to three hours after the light period had started in order to maximize the NRA (Duke et al., 1978). The leaves were prepared for incubation by removing the uppermost expanded leaves from each plant in the pot (Chu et al., 1980; Hallam and Blackwood, 1979; Peirson and Elliott, 1981). The leaves of Helianthus annuus and Ambrosia trifida were cut into 1 cm^2 pieces and the leaves of the grasses were cut into pieces approximately 1 cm long. The leaf segments for each pot were pooled and kept on ice. The roots were prepared by washing them free of vermiculite, then cutting them into 1 cm long pieces. The root segments from each pot were then pooled and kept on ice. The incubations were conducted in foil-wrapped two dram vials. The incubation medium (minus n-propanol) had been previously added to the vials and chilled to 5°C 12 hr before the incubations. Approximately 0.2-0.5 g fresh weight of the segments was added to a vial. It was found that larger samples were sometimes necessary to accurately measure the NRA in species that had low levels of activity. N-propanol was added after the segments had been placed in the vials. It was found that high levels (10%v/v) were necessary in order to wet and sink the leaf segments of some of

the grass species when the n-propanol was added to the incubation medium prior to addition of the sample. These high levels of n-propanol caused an interference in the determination of nitrite by allowing the leaching of some substance into the incubation medium that absorbed at 540 nm. Adding the n-propanol as a pulse after the segments had been placed in the vials allowed for the thorough wetting and sinking of the sample at lower n-propanol concentrations (2-5%v/v) and avoided the interference problem. The roots and leaves of the plants from each pot were tested in triplicate. The vials were sealed and were incubated in a water bath at 30°C. At the end of the incubation period, a 0.4 ml sample was withdrawn from each vial. Color formation for nitrite determination was obtained by adding 0.3 ml of a 1% sulfanilamide reagent and 0.3 ml of a 0.02% N-naphthylethylene diamine HCl solution. The sample was diluted to 5 ml with distilled water and absorption was measured at 540 nm. The amount of nitrite present was computed from a standard curve generated the day of the test. The tissue was removed from the incubation medium, rinsed twice with distilled water, and dried at 80°C for 24 hr. The dry weight of the sample was obtained and the NRA was calculated as $\mu\text{M NO}_2/\text{g dry wt/hr}$. The NRA was calculated on a dry weight rather than a fresh weight basis because of the differing water content of the organs within a species and between species. The data were found to be heterogenous; therefore, a log transformation was made before a nested Analysis of Variance (ANOVA) was calculated. The triplicate readings for the leaves or roots of the plants from each pot were nested within each replicate pot of a nitrogen treatment. Differences between means were determined by a Student-Newman-Keuls test. The

total shoot material per pot was saved and dried for 48 hr at 80°C, and the shoot dry weights were recorded. The shoot data were analyzed by ANOVA and differences between means were determined as before.

Results

The preliminary experiments to obtain the incubation conditions for maximal NRA were necessary. The NRA could vary significantly depending on the combination of conditions used (Table 1). The incubation conditions that were used in the determination of NRA for the roots and leaves of each species are given in Table 2. The optimal concentration of phosphate buffer and substrate was found to vary between the roots and leaves within a species and between species. The time required for a steady state of NRA was the same for both organs of the species, with the exception of Bromus japonicus and Andropogon gerardii, where the roots required less time than the leaves. There did not appear to be any correlation between the optimal incubation conditions and the successional stage of a species.

Both organs of each species tested showed some NRA. The amount of the activity varied between the roots and leaves, with one organ in a given species consistently showing higher NRA than the other under all nitrogen treatments. Wallace and Pate (1965; 1967) and Oghoghorie and Pate (1972) categorized plants based on the organ of highest NRA. Five species were found to resemble Xanthium pennsylvanicum by having most

of their NRA in the leaves (Table 3) and three species were found to resemble Pisum arvense by having most of their NRA in the roots (Table 4). There was no apparent correlation between the organ with the highest NRA and the successional stage of the species.

The NRA of the leaves of the leaf-high pioneer species Ambrosia trifida, Aristida oligantha, and Helianthus annuus was always significantly higher ($\alpha=.01$) than that of the leaves of the climax species Andropogon gerardii and Panicum virgatum, regardless of the nitrogen treatment (Table 3). The same pattern occurred in the roots except in the ammonium only treatment, where Helianthus annuus and Andropogon gerardii were not different. There were also some significant differences between the NRA of the leaves and of the roots within the pioneer species and the climax species for one or more of the nitrogen treatments. The root-high species (Table 4) showed a similar correlation between the successional stage and NRA. The pioneer species, Bromus japonicus, was always significantly higher ($\alpha=.01$) in NRA than the two climax species Sorghastrum nutans and Schizachyrium scoparium, regardless of the organ or nitrogen treatment. Sorghastrum nutans was also always higher in NRA than Schizachyrium scoparium.

The pattern of NRA between nitrogen treatments for organs of a species was variable (Table 5). In general, NRA decreased with decreasing nitrate in the treatment. This was expected because NR is a substrate induced enzyme (Beevers et al., 1965; Beevers and Hageman, 1969). In some instances, such as Helianthus annuus leaves, the nitrate only, 2:1, and 1:2 treatments did not show any significant difference in NRA, but all three treatments were significantly higher in NRA than the

leaves from the ammonium only treatment. In cases such as Ambrosia trifida roots, the NRAs of the 2:1 and 1:2 nitrogen treatments were not significantly different from each other, but they were intermediate between the NRA of the nitrate and ammonium only treatments. Sorghastrum nutans and Schizachyrium scoparium roots showed a pattern of NRA in which the nitrate only and 2:1 treatments grouped together, the 1:2 treatment was intermediate and the ammonium only treatment was significantly less. Seven cases were found where there was no apparent inducibility of the enzyme because there were no significant differences between any of the nitrogen treatments. Four of these cases (Schizachyrium scoparium leaves, Sorghastrum nutans leaves, Andropogon gerardii roots, and Aristida oligantha roots) occurred in the organ that had the lowest NRA for that species. Three cases (Panicum virgatum leaves, Ambrosia trifida leaves, and Bromus japonicus roots) occurred in the organ with the highest NRA of that species. One of these cases was re-examined (Bromus japonicus roots) in an attempt to discover why the enzyme was apparently not inducible. It was determined that four week old plants did show substrate inducibility, but this ability was lost when the plants were eight weeks old. A time plot of NRA during the incubation of eight week old Bromus japonicus roots (Fig. 1) indicated that the nitrate grown plants showed NRA almost immediately while the ammonium grown plants showed a lag of 30-45 min before NRA could be detected. The final NRA shown by the ammonium grown plants was somewhat less than that of the nitrate grown plants, but the difference was not significant.

The shoot dry weights indicated a correlation between successional stage and response to the nitrogen treatment (Table 6). All eight species

attained their highest shoot dry weight with the ammonium only treatment. Three of the pioneer species (Helianthus annuus, Aristida oligantha, and Bromus japonicus) had no significant differences in shoot dry weight between the nitrate only, 2:1, and 1:2 nitrogen treatments. The shoot dry weight in the ammonium only treatment was significantly higher than in any combination of nitrate/ammonium or nitrate only. Ambrosia trifida had an increase in shoot dry weight with the shifting of the nitrate/ammonium ratio from nitrate only to ammonium only. All climax species had two groups of shoot dry weight response. The nitrate only and the 2:1 treatments grouped together and the 1:2 and ammonium only treatments grouped together. The all or mostly nitrate group had a significantly lower shoot dry weight than the all or mostly ammonium group.

Discussion

The pattern of NRA was correlated with the successional stage of the species as had been hypothesized. These results are in complete agreement with the information obtained in other studies. Havill et al. (1974) reported that ruderal plants had the highest NRA of any species they tested. Both Bate and Heelas (1975) and Franz and Haines (1977) reported that the pioneer species had high NRA while the climax species had low NRA. The field NRA values obtained by Franz and Haines were very close to the values reported here. They found an average NRA

of $47 \mu\text{M NO}_2/\text{g dry wt/hr}$ for the herbs in their old field stage 1, and an average value between 1 and $2 \mu\text{M NO}_2/\text{g dry wt/hr}$ for the leaves of the climax tree species (personal communication).

It is difficult to make direct comparisons of the NRA values reported in this study with other NRA values found in the literature. Much of the literature information is based on seedlings of cultivated plants grown at ecologically unrealistic levels of available nitrogen. In some cases there was no indication as to whether preliminary tests were made to maximize the NRA. For example, in contrast to my findings, Weissman (1972) reported that Helianthus annuus had most of its NRA in the roots and little in the leaves. Also, no NRA was detected for plants grown in ammonium only. He used 19 day old seedlings of cultivated Helianthus annuus that were grown on 10 mM of NH_4Cl or 10 mM KNO_3 , and the enzyme analysis was done in vitro. Weissman (1964) also reported that the shoot dry weight of Helianthus annuus seedlings was highest on a mixture of nitrate and ammonium rather than on nitrate or ammonium alone. It is difficult to make meaningful comparisons between our results because of the difference in the methods and materials. The ecological relativity of studies using cultivated species grown at high nitrogen concentrations is questionable. One difference between the studies might possibly be explained, however, on the basis that the relative amounts of root and leaf NRA can shift from one organ to another during a growing season (Van Berkum and Sloger, 1981; Yuan and Shien, 1980).

The occurrence of NRA in both leaves and roots of ammonium grown plants for all eight species tested would appear to be contradictory to

the induction of the enzyme by substrate. This also suggests that low levels of constitutive enzyme are present and that induction increases the NRA above this base level. Several studies have indicated, however, that either NR or a NR precursor is present in ammonium grown plants (Ferguson, 1969; Funkhouser and Ramadoss, 1980; Funkhouser et al., 1980) or the plants have preformed messenger RNA for NR synthesis (Hipkin and Syrett, 1977). Other workers have indicated that both light (Jolly and Tolbert, 1978; Tischner and Hüttermann, 1978) and substrate (Pistorius et al., 1976) induction of NR involves the shift of NR from an inactive to an active form. Orebamjo and Stewart (1975) suggested that the shifting of inactive to active NR can be reversed and may involve protein-protein interaction. They hypothesized that inactivation of NR by an ammonium triggered protein would enable the plant to use ammonium preferentially and prevent the energy expense of reducing nitrate. Such an inhibitor could be removed when the ammonium concentration falls below some threshold level, reactivating the NR. Ferguson and Bollard (1969) reported that Spirodela oligorrhiza could use ammonium preferentially when both ammonium and nitrate were present. Little nitrate was reduced until the ammonium had been removed from the medium. Shen (1969) reported a similar response in rice seedlings, but also found that the addition of ammonium shut down NRA and that the NRA was resumed only when the ammonium was depleted. The lag phase observed in this study in the NRA of ammonium grown Bromus japonicus roots suggests that this species may have an inactivated NR caused by the presence of ammonium, but that the NR can be activated by the nitrate in the incubation medium. This could explain why the NRA values for

Bromus japonicus roots were not significantly different between the nitrogen treatments. It is not known whether a similar situation exists in the six other cases that did not show NR inducibility. In all seven cases, one of the two organs showed inducibility by substrate while the other organ did not. This suggests that at least one organ of each species could change NRA in response to changing nitrate concentrations while the other organ may not. Radin (1975) suggested that roots and shoots of cotton plants responded differently to ammonium inhibition and that each tissue might be under a different control system. Similar differences in control systems could exist for the seven cases of non-inducibility found here.

The ability to inhibit NR in order to utilize ammonium preferentially could be very important ecologically. A plant could use ammonium as the more efficient nitrogen source when it is available and reduce the energy expense of incorporating nitrogen into its metabolism. If the ammonium should become depleted, the NR could be activated and nitrate could be utilized, although at a greater energetic cost. The shoot dry weight data showed that the pioneer and climax species differ in their growth response to changing nitrate/ammonium ratios. Three of the pioneer species had the same shoot dry weights regardless of the nitrate/ammonium ratio. Their dry weights were increased significantly solely in the presence of ammonium only. All four climax species had the same shoot dry weight in the 1:2 or all ammonium treatments. Thus they demonstrate a greater preference for ammonium than do the pioneer species. The low levels of NR in the climax species may require lower concentrations of ammonium for inactivation than the higher levels of

NR in the pioneer species. This would enable the climax species to inactivate their NR at low ammonium concentrations and use ammonium preferentially as their primary nitrogen source. Pioneer plants can use ammonium for their nitrogen source, but it could take a higher concentration of ammonium to inactivate their NR and allow them to preferentially utilize ammonium. Because the soil of the pioneer stage is low in ammonium, it would be a disadvantage for these species to inactivate their NR and use ammonium as their primary nitrogen source. Therefore, these species would need to have high NRA in order to reduce and utilize the more abundant nitrate in order to meet their metabolic nitrogen demands. The soil of the climax stage is higher in ammonium than nitrate, therefore, the climax species can inactivate their NR and utilize ammonium preferentially without suffering from nitrogen deficiency. It would not be advantageous for them to produce NR and have high NRA. It should be noted that all four climax species grew perfectly well on nitrate only, even though they have low levels of NRA. Thus, they can obviously use nitrate if it is the only nitrogen source available to them. This observation that these species can use nitrate indicates that the low levels of NRA are not attributable to restricted uptake of the nitrate. Therefore, the climax species resemble the NR-deficient mutants reported by Warner and Kleinhofs (1981) in that they have low levels of NRA but are still able to grow normally on nitrate only. The present study does not conclusively prove the NR-active to NR-inactive switch, but it does provide evidence that several factors may be involved in obtaining and utilizing soil nitrogen.

It is interesting to note that there is a significant difference between the NRA of the species within each successional group of plants. Part of the patterning observed in the pioneer and climax stage might possibly be related to the NRA differences and the patchiness of nitrogen in the soil.

Rice and Pancholy's (1972) hypothesis that climax vegetation inhibits nitrification has been examined in a large variety of successional systems. In some studies the evidence supports the theory (Haines, 1977; Lodhi, 1979; 1981; Smith et al., 1968; Todd et al., 1975) and in other studies it does not (Johnson and Edwards, 1979; Lamb, 1980; Purchase, 1974; Rennie et al., 1977; Robertson and Vitousek, 1981). In some instances the evidence is not clear (Montes and Christensen, 1979) or the inhibition is dependent on the parent material from which the soils under the climax vegetation were derived (Nakos, 1977). Alternative hypothesis have also been suggested (Gorham et al., 1979; Vitousek, 1977; Vitousek and Reiners, 1975). It would be premature to generalize that all successional systems exhibit the same NRA pattern that was found in this study. If the soil nitrogen form is observed to shift from nitrate to ammonium during the succession, a similar adaptation of the NRA of the species might be expected. The data presented here show that the distribution of plant species across a sere can be correlated with their ability to utilize the nitrogen form available to them. The conclusion is that the successional position of a species is correlated to its physiological adaptation to the nitrogen form in the soil.

Table 1. Variations in mean Nitrate Reductase Activity ($\mu\text{M NO}_2/\text{g}$ dry wt/hr) in Ambrosia trifida leaves and roots according to the phosphate buffer and nitrate concentrations. The pH was 7.5, the temperature was 30°C , and the period of incubation was 2 hr in all cases.

Substrate (KNO_3)	<u>Phosphate Buffer</u>			
	1 mM	10 mM	50 mM	100 mM
<u>10 mM</u>				
leaves	33	28	24	30
roots	38	37	19	12
<u>50 mM</u>				
leaves	89	80	33	39
roots	40	18	9	12
<u>100 mM</u>				
leaves	98	88	65	31
roots	40	42	15	13
<u>150 mM</u>				
leaves	104	89	34	33
roots	26	33	15	14
<u>200 mM</u>				
leaves	75	71	40	34
roots	26	30	13	11

Table 2. Optimal conditions for in vivo Nitrate Reductase Activity

	mM PO_4 Buffer	mM KNO_3	Time(hr)	n-propanol(v/v)
<u>Pioneer Species</u>				
<u>Helianthus annuus</u>				
leaves	50	50	4	3%
roots	100	200	4	5%
<u>Ambrosia trifida</u>				
leaves	1	150	2.5	1%
roots	1	50	2.5	5%
<u>Aristida oligantha</u>				
leaves	10	50	4	0%
roots	10	50	4	0%
<u>Bromus japonicus</u>				
leaves	1	50	4	0%
roots	50	10	2	0%

Table 2. Continued

	mM PO ₄ Buffer	mM KNO ₃	Time(hr)	n-propanol(v/v)
<u>Climax Species</u>				
<u>Schizachyrium scoparium</u>				
leaves	100	150	2	2%
roots	100	50	2	3%
<u>Sorghastrum nutans</u>				
leaves	1	50	2.5	6%
roots	10	50	2.5	3%
<u>Panicum virgatum</u>				
leaves	50	100	4	2%
roots	50	50	4	4%
<u>Andropogon gerardii</u>				
leaves	50	50	2	1%
roots	50	50	1.5	0%

Table 3. Nitrate Reductase Activity of leaf-high species ($\mu\text{M NO}_2/\text{g}$ dry wt/hr, antilog of log mean $\pm$ standard deviation). Comparisons are made vertically in the table. Numbers that share the same letter are not statistically different at the .01 level.

	<u>Nitrogen Treatment</u>			
	NO_3	2:1	1:2	NH_4
<u>Leaves</u>				
<u>Pioneer Species</u>				
<u>Ambrosia trifida</u>	53.9 $\pm$ 1.2a	45.1 $\pm$ 1.3a	45.4 $\pm$ 1.2a	31.9 $\pm$ 1.7a
<u>Aristida oligantha</u>	45.1 $\pm$ 1.2a	41.9 $\pm$ 1.3a	43.1 $\pm$ 1.2a	28.5 $\pm$ 1.3a
<u>Helianthus annuus</u>	22.4 $\pm$ 1.3ab	22.5 $\pm$ 1.3b	22.4 $\pm$ 1.1b	13.1 $\pm$ 1.4b
<u>Climax Species</u>				
<u>Andropogon gerardii</u>	10.5 $\pm$ 1.7bc	8.0 $\pm$ 1.7c	7.7 $\pm$ 1.4c	5.6 $\pm$ 1.6c
<u>Panicum virgatum</u>	7.4 $\pm$ 1.4c	5.5 $\pm$ 1.7d	5.2 $\pm$ 1.7d	4.8 $\pm$ 1.7c
<u>Roots</u>				
<u>Pioneer Species</u>				
<u>Ambrosia trifida</u>	19.2 $\pm$ 1.3a	15.8 $\pm$ 1.2a	14.6 $\pm$ 1.4a	4.6 $\pm$ 1.6a
<u>Aristida oligantha</u>	12.2 $\pm$ 2.5b	5.3 $\pm$ 3.0b	11.9 $\pm$ 2.6a	5.4 $\pm$ 2.3a
<u>Helianthus annuus</u>	8.6 $\pm$ 1.4c	6.7 $\pm$ 1.8b	7.6 $\pm$ 1.3b	2.4 $\pm$ 1.6b
<u>Climax Species</u>				
<u>Andropogon gerardii</u>	4.3 $\pm$ 2.2d	3.4 $\pm$ 2.1c	3.8 $\pm$ 1.9c	3.0 $\pm$ 1.6b
<u>Panicum virgatum</u>	3.3 $\pm$ 2.2d	2.1 $\pm$ 3.0d	1.8 $\pm$ 3.2d	1.0 $\pm$ 3.3c

Table 4. Nitrate Reductase Activity of root-high species. ($\mu\text{M NO}_2/\text{g}$ dry wt/hr, antilog of log mean $\pm$ standard deviation).

Letters as in Table 3.

	<u>Nitrogen Treatment</u>			
	NO_3	2:1	1:2	NH_4
<u>Leaves</u>				
<u>Pioneer Species</u>				
<u>Bromus japonicus</u>	27.7 $\pm$ 1.5a	32.4 $\pm$ 1.3a	27.4 $\pm$ 1.6a	19.5 $\pm$ 1.7a
<u>Climax Species</u>				
<u>Sorghastrum nutans</u>	4.6 $\pm$ 3.0b	4.7 $\pm$ 3.7b	6.5 $\pm$ 2.2b	4.4 $\pm$ 2.7b
<u>Schizachyrium scoparium</u>	2.0 $\pm$ 1.6c	1.8 $\pm$ 1.7c	1.5 $\pm$ 1.6c	1.5 $\pm$ 1.7c
<u>Roots</u>				
<u>Pioneer Species</u>				
<u>Bromus japonicus</u>	61.1 $\pm$ 3.5a	82.3 $\pm$ 1.8a	88.9 $\pm$ 1.9a	79.8 $\pm$ 1.8a
<u>Climax Species</u>				
<u>Sorghastrum nutans</u>	12.1 $\pm$ 2.2b	12.4 $\pm$ 1.2b	7.8 $\pm$ 1.5b	5.8 $\pm$ 1.7b
<u>Schizachyrium scoparium</u>	4.9 $\pm$ 1.7c	4.2 $\pm$ 1.8c	3.1 $\pm$ 2.8c	1.5 $\pm$ 1.8c

Table 5. Nitrate Reductase Activity of species grown on different nitrogen treatments ($\mu\text{M NO}_2/\text{g dry wt/hr}$, antilog of $\log \text{mean} \pm \text{standard deviation}$). Comparisons are made horizontally in the table. Numbers that share the same letters are not significantly different at the .01 level.

	NO ₃	<u>Nitrogen Treatment</u>		
		2:1	1:2	NH ₄
<hr/>				
<u>Pioneer Species</u>				
<u>Helianthus annuus</u>				
leaves	22.4±1.3a	22.5±1.3a	22.4±1.1a	13.0±1.4b
roots	8.6±1.4a	6.7±1.8b	7.6±1.3ab	2.4±1.6c
<u>Ambrosia trifida</u>				
leaves	53.9±1.2a	45.0±1.3a	45.5±1.2a	31.9±1.7a
roots	19.2±1.3a	15.8±1.2b	14.6±1.4b	4.6±1.6c
<u>Aristida oligantha</u>				
leaves	45.1±1.2a	43.1±1.3a	41.9±1.2a	28.5±1.3b
roots	12.6±2.5a	11.9±3.0a	5.3±2.6a	5.4±2.3a
<u>Bromus japonicus</u>				
leaves	27.7±1.5a	32.4±1.3a	27.4±1.6a	19.5±1.7b
roots	61.1±3.5a	82.3±1.8a	88.9±1.9a	79.8±1.8a

Table 5. Continued

		<u>Nitrogen Treatment</u>			
		NO ₃	2:1	1:2	NH ₄
<hr/>					
<u>Climax Species</u>					
<u>Schizachyrium scoparium</u>					
leaves	2.0±1.6a	1.8±1.7a	1.5±1.6a	1.5±1.7a	
roots	4.8±1.7a	4.2±1.8a	3.1±2.8b	1.5±1.8c	
<u>Sorghastrum nutans</u>					
leaves	4.6±3.0a	4.7±3.7a	6.5±2.2a	4.4±2.7a	
roots	12.1±2.2a	12.4±1.2a	7.8±1.5b	5.8±1.7c	
<u>Panicum virgatum</u>					
leaves	7.4±1.4a	5.5±1.7a	5.2±1.7a	4.8±1.7a	
roots	3.3±2.2a	2.1±3.0b	1.9±3.2b	1.0±3.5c	
<u>Andropogon gerardii</u>					
leaves	10.5±1.7a	8.1±1.7b	7.7±1.4b	5.6±1.6c	
roots	4.3±2.2a	3.8±2.1a	3.4±1.9a	3.0±1.6a	

Table 6. Shoot dry weight (g) of species grown on different nitrogen treatments (mean $\pm$ standard deviation). Comparisons are made horizontally in the table. Numbers that share the same letter are not significantly different at the .01 level except for species marked with an *, where the comparison is at the .05 level.

	<u>Nitrogen Treatment</u>			
	NO_3	2:1	1:2	NH_4
<u>Pioneer Species</u>				
<u>Helianthus annuus</u>	2.76 $\pm$ 0.27a	3.31 $\pm$ 1.01a	2.66 $\pm$ 0.25a	4.36 $\pm$ 0.76b
<u>Ambrosia trifida</u>	2.29 $\pm$ 0.30a	2.83 $\pm$ 0.33ab	3.24 $\pm$ 0.38bc	3.35 $\pm$ 0.81c
<u>Aristida oligantha*</u>	0.81 $\pm$ 0.26a	0.90 $\pm$ 0.43a	1.13 $\pm$ 0.44a	1.49 $\pm$ 0.41b
<u>Bromus japonicus*</u>	1.10 $\pm$ 0.13a	0.99 $\pm$ 1.14a	1.12 $\pm$ 0.15a	1.34 $\pm$ 0.20b
<u>Climax Species</u>				
<u>Schizachyrium scoparium</u>	1.73 $\pm$ 0.36a	2.12 $\pm$ 0.30a	2.65 $\pm$ 0.37b	2.79 $\pm$ 0.40b
<u>Sorghastrum nutans</u>	0.68 $\pm$ 0.19a	0.79 $\pm$ 0.23a	1.36 $\pm$ 0.24b	1.54 $\pm$ 0.23b
<u>Panicum virgatum*</u>	1.37 $\pm$ 0.33a	1.71 $\pm$ 0.30a	2.03 $\pm$ 0.34b	2.09 $\pm$ 0.36b
<u>Andropogon gerardii</u>	0.59 $\pm$ 0.13a	0.61 $\pm$ 0.13a	0.98 $\pm$ 0.17b	0.82 $\pm$ 0.10b

Figure 1. Nitrate Reductase Activity of eight week old Bromus japonicus roots grown on nitrate only ($\bullet$ $y = 45.25x - 15.57x^2 + 1.71x^3 - 15.12$, $r^2 = .96$) or ammonium only ($\blacktriangle$ $y = 42.22x - 11.52x^2 + 0.99x^3 - 25.45$, $r^2 = .97$).

References

- Abdul-Wahab AS, Rice EL (1967) Plant inhibition by Johnson grass and its possible significance in old field succession. Bull Torrey Bot Club 94: 486-497
- Addoms RM (1937) Nutritional studies on loblolly pine. Plant Physiol 12: 199-205
- Barker AV, Volk RJ, Jackson WA (1966) Growth and nitrogen distribution patterns in bean plants (Phaseolus vulgaris L.) subjected to ammonium nutrition. I. Effects of carbonate and acidity control. Soil Sci Soc Amer Proc 30: 228-232
- Bate GC, Heelas BV (1975) Studies on the nitrate nutrition of two indigenous Rhodesian grasses. J Appl Ecol 12: 941-952
- Beevers L, Hageman RH (1969) Nitrate reduction in higher plants. Annual Rev Plant Physiol 20: 495-522
- Beevers L, Hageman RH (1980) Nitrate and nitrite reduction. The Biochemistry of Plants Vol 5. Academic Press Inc. p 115-168
- Beevers L, Schrader LE, Flesher D, Hageman RH (1965) The role of light and nitrate in the induction of nitrate reductase in radish cotyledons and maize seedlings. Plant Physiol 40: 691-698
- Bigg WL, Daniel TW (1978) Effects of nitrate, ammonium, and pH on the growth of conifer seedlings and their production of nitrate reductase. Plant and Soil 50: 371-385
- Booth WE (1941) Revegetation of abandoned fields in Kansas and Oklahoma. Amer J Bot 28: 415-422

- Chu C-C, Sweet RD, Ozbun JL, Kaplan SL (1980) Nitrate reduction in redroot pigweed (Amaranthus retroflexus) and common lambsquarters (Chenopodium album) as a function of leaf age and photosynthetic photon flux density. *Weed Sci* 28: 484-486
- Duke SH, Friedrich JW, Schrader LE, Koukkari WL (1978) Oscillations in the activities of enzymes of nitrate reduction and ammonia assimilation in Glycine max and Zea mays. *Physiol Plant* 42: 269-276
- Ferguson AR (1969) The nitrogen metabolism of Spirodela oligorrhiza II. Control of the enzymes of nitrate assimilation. *Planta* 88: 353-363
- Ferguson AR, Bollard EG (1969) Nitrogen metabolism of Spirodela oligorrhiza I. Utilization of ammonium, nitrate and nitrite. *Planta* 88: 344-352
- Franz EH, Haines BL (1977) Nitrate reductase activities of vascular plants in a terrestrial sere: relationship of nitrate to uptake and the cybernetics of biogeochemical cycles. *Bull Ecol Soc Amer* 58: 62
- Funkhouser EA, Ramadoss CS (1980) Synthesis of nitrate reductase in Chlorella II. Evidence for synthesis in ammonia-grown cells. *Plant Physiol* 65: 944-948
- Funkhouser EA, Shen T-C, Aakermann R (1980) Synthesis of nitrate reductase in Chlorella I. Evidence for an inactive protein precursor. *Plant Physiol* 65: 939-943

- Gigon A, Rorison IH (1972) The response of some ecologically distinct plant species to nitrate- and ammonium-nitrogen. *J Ecol* 60: 93-102
- Gorham E, Vitousek PM, Reiners WA (1979) The regulation of chemical budgets over the course of terrestrial ecosystem succession. *Ann Rev Ecol Syst* 10: 53-84
- Goring CAI (1962a) The control of nitrification by 2-chloro-6-(trichloromethyl)pyridine. *Soil Sci* 93: 211-218
- Goring CAI (1962b) Control of nitrification of ammonium fertilizers and urea by 2-chloro-6-(trichloromethyl)pyridine. *Soil Sci* 93: 431-439
- Greidanus T, Peterson LA, Schrader LE, Dana MN (1972) Essentiality of ammonium for cranberry nutrition. *J Amer Soc Hort Sci* 97: 272-277
- Haines BL (1977) Nitrogen uptake: apparent pattern during old-field succession in southeastern United States. *Oecologia* 26: 295-303
- Hallam R, Blackwood GC (1979) Nitrate reductase activity in wheat (Triticum aestivum L.) I. Distribution throughout the leaves. *New Phytol* 82: 407-415
- Havill DC, Lee JA, Stewart GR (1974) Nitrate utilization by species from acidic and calcareous soils. *New Phytol* 73: 1221-1231
- Haynes RJ, Goh KM (1978) Ammonium and nitrate nutrition of plants. *Biol Rev* 53: 465-510
- Hipkin CR, Syrett PJ (1977) Post-transcriptional control of nitrate reductase formation in green algae. *J Exp Bot* 28: 1270-1277

- Jaworski EG (1971) Nitrate reductase assay in intact plant tissues.
Biochem Biophys Res Comm 43: 1274-1279
- Johnson DW, Edwards NT (1979) The effects of stem girdling on
biogeochemical cycles within a mixed deciduous forest in eastern
Tennessee. II. Soil nitrogen mineralization and nitrification
rates. Oecologia 40: 259-271
- Jolly SO, Tolbert NE (1978) NADH-Nitrate reductase inhibitor from
soybean leaves. Plant Physiol 62: 197-203
- Kannangara CC, Woolhouse HW (1967) The role of carbon dioxide, light
and nitrate in the synthesis and degradation of nitrate reductase
in leaves of Perilla frutescens. New Phytol 66: 553-561
- Karin AGH, Vlamis J (1962) Comparative study of the effects of ammonium
and nitrate nitrogen in the nutrition of rice. Plant and Soil 16:
32-41
- Kapustka LA, Rice EL (1976) Acetylene reduction (N_2 fixation) in soil
and old field succession in central Oklahoma. Soil Biol Biochem
8: 497-503
- Kirkby EA (1968) Influence of ammonium and nitrate nutrition on the
cation-anion balance and nitrogen and carbohydrate metabolism of
white mustard plants grown in dilute nutrient solutions. Soil Sci
105: 133-141
- Klepper L, Flesher D, Hageman RH (1971) Generation of reduced
nicotinamide adenine dinucleotide for nitrate reduction in green
leaves. Plant Physiol 48: 580-590
- Krajina VJ, Madoc-Jones S, Mellor G (1973) Ammonium and nitrate in the
nitrogen economy of some conifer growing in Douglas-fire communities

- of the pacific northwest of America. *Soil Biol Biochem* 5: 143-147
- Lamb D (1980) Soil nitrogen mineralisation in a secondary rainforest succession. *Oecologia* 47: 257-263
- Lodhi MAK (1979) Inhibition of nitrifying bacteria, nitrification and mineralization in spoil soils as related to their successional stages. *Bull Torrey Bot Club* 106: 284-289
- Lodhi MAK (1981) Accelerated soil mineralization, nitrification and revegetation of abandoned fields due to the removal of crop-soil phytotoxicity. *J Chem Ecol* 7: 685-694
- McFee WW, Stone EL Jr (1968) Ammonium and nitrate as nitrogen sources for Pinus radiata and Picea glauca. *Soil Sci Soc Amer Proc* 32: 379-384
- Montes RA, Christensen NL (1979) Nitrification and succession in the Piedmont of North Carolina. *Forest Sci* 25: 287-297
- Moore TC (1974) *Research Experiences in Plant Physiology, a Laboratory Manual*. Springer-Verlag, New York p 421
- Moris HD, Giddens J (1963) Response of several crops to ammonium and nitrate forms of nitrogen as influenced by soil fumigation and liming. *Agron J* 55: 372-374
- Nakos G (1977) Interaction of nitrogen fertilizers and forest humus. II. Beech and Oak. *Soil Biol Biochem* 9: 423-426
- Neill RL, Rice EL (1971) Possible role of Ambrosia psilostachya on pattern and succession in old fields. *Amer Midland Natur* 86: 344-357

- Oghoghoric CGO, Pate JS (1972) Exploration of the nitrogen transport system of a nodulated legume using ^{15}N . *Planta* 104: 35-49
- Orebamjo TO, Stewart GH (1975) Ammonium inactivation of nitrate reductase in Lemna minor L. *Planta* 122: 37-44
- Parenti RL, Rice EL (1969) Inhibitional effects of Digitaria sanguinalis and possible role in old-field succession. *Bull Torrey Bot Club* 96: 70-78
- Parks JM, Rice EL (1969) Effect of certain plants of old-field succession on the growth of blue-green algae. *Bull Torrey Bot Club* 96: 345-360
- Peirson DR, Elliott JR (1981) In vivo nitrite reduction in leaf tissue of Phaseolus vulgaris L. *Plant Physiol* 68: 1068-1072
- Pistorius E, Gewitz H-S, Voss H, Vennesland B (1976) Reversible inactivation of nitrate reductase in Chlorella vulgaris in vivo. *Planta* 128: 73-80
- Purchase BS (1974) Evaluation of the claim that grass root exudates inhibit nitrification. *Plant and Soil* 41: 527-539
- Radin JW (1975) Differential regulation of nitrate reductase in roots and shoots of cotton plants. *Plant Physiol* 55: 178-182
- Rennie RJ, Reyes VG, Schmidt EL (1977) Immunofluorescence detection of the effects of wheat and soybean roots on Nitrobacter in soil. *Soil Sci* 124: 10-15
- Rice EL (1964) Inhibition of nitrogen-fixing and nitrifying bacteria by seed plants I. *Ecology* 45: 824-837
- Rice EL (1968) Inhibition of nodulation of inoculated legumes by pioneer

- plant species from abandoned fields. Bull Torrey Bot Club 95: 346-358
- Rice EL (1976) Allelopathy and grassland improvement. In: JR Estes and RJ Tyr1 (eds) The grasses and grasslands of Oklahoma. Ann Okla Acad Sci No. 6, Noble Foundation. Ardmore, Oklahoma. p 90-111
- Rice EL, Pancholy SK (1972) Inhibition of nitrification by climax ecosystems. Amer J Bot 59: 1033-1040
- Rice EL, Pancholy SK (1973) Inhibition of nitrification by climax ecosystems II. Additional evidence and possible role of tannins. Amer J Bot 60: 691-702
- Rice EL, Pancholy SK (1974) Inhibition of nitrification by climax ecosystems III. Inhibitors other than tannins. Amer J Bot 61: 1095-1103
- Rice EL, Penfound WT, Rohrbaugh LH (1960) Seed dispersal and mineral nutrition in succession in abandoned fields in central Oklahoma. Ecology 41: 224-228
- Robertson GP, Vitousek PM (1981) Nitrification potentials in primary and secondary succession. Ecology 62: 376-386
- Schrader LE, Domska D, Jung PE Jr, Peterson LA (1972) Uptake and assimilation of ammonium-N and nitrate-N and their influence on the growth of corn (Zea mays L.). Agron J 64: 690-695
- Shen TC (1969) The induction of nitrate reductase and the preferential assimilation of ammonium in germinating rice seedlings. Plant Physiol 44: 1650-1655
- Smith WH, Bormann FH, Likens GE (1968) Response of chemoautotrophic nitrifiers to forest cutting. Soil Sci 106: 471-473

- Tischner R, Hüttemann A (1978) Light-mediated activation of nitrate reductase in synchronous Chlorella. Plant Physiol 62: 284-286
- Todd RL, Swank WT, Douglass JE, Keer PC, Brockway DL, Monk CP (1975) The relationship between nitrate concentration in the southern Appalachian mountain streams and terrestrial nitrifiers. Agro-Ecosystem 2: 127-132
- Van Berkum P, Sloger C (1981) Ontogenetic variation in nitrogenase, nitrate reductase, and glutamine synthetase activities in Oryza sativa. Plant Physiol 68: 722-726
- Van Den Driessche R (1971) Response of conifer seedlings to nitrate and ammonium sources of nitrogen. Plant and Soil 34: 421-439
- Vitousek PM (1977) The regulation of element concentrations in mountain streams in the northeastern United States. Ecol Monog 47: 65-87
- Vitousek PM, Reiners WA (1975) Ecosystem succession and nutrient retention: a hypothesis. BioScience 25: 376-381
- Wallace W, Pate JS (1965) Nitrate reductase in the field pea (Pisum arvense L.). Ann Bot 29: 655-671
- Wallace W, Pate JS (1967) Nitrate assimilation in higher plants with special reference to the cocklebur (Xanthium pennsylvanicum Wallr.). Ann Bot NS 31: 213-228
- Warner RL, Kleinhofs A (1981) Nitrate utilization by nitrate reductase-deficient barley mutants. Plant Physiol 67: 740-743
- Weissman GS (1964) Effect of ammonium and nitrate nutrition on protein level and exudate composition. Plant Physiol 39: 947-952

- Weissman GS (1972) Influence of ammonium and nitrate nutrition on enzymatic activity in soybean and sunflower. *Plant Physiol* 49: 138-141
- Wilson RE, Rice EL (1968) Allelopathy as expressed by Helianthus annuus and its role in old field succession. *Bull Torrey Bot Club* 95: 432-448
- Wiltshire GH (1973) Response of grasses to nitrogen source. *J Appl Ecol* 10: 429-435
- Yuan H-F, Shien Y-S (1980) Seasonal variation of nitrate reductase, glutamate dehydrogenase and the soluble nitrogenous compounds during rice growth. *Bot Bull Acad Sin* 21: 35-52