

THE INFLUENCE OF HABITAT QUALITY ON THE
POPULATION DECLINE OF THE RED-COCKADED
WOODPECKER IN THE MCCURTAIN COUNTY
WILDERNESS AREA, OKLAHOMA

By

JEFFREY F. KELLY

Bachelor of Science

University of Maine

Orono, Maine


1987

Submitted to the Faculty of the
Graduate College of the
Oklahoma State University
in partial fulfillment of
the requirements for
the Degree of
MASTER OF SCIENCE
May, 1991


Thesis
1991
K29,
cop. 2


THE INFLUENCE OF HABITAT QUALITY ON THE
POPULATION DECLINE OF THE RED-COCKADED
WOODPECKER IN THE MCCURTAIN COUNTY
WILDERNESS AREA, OKLAHOMA

Thesis Approved:



Thesis Adviser







Dean of the Graduate College

ACKNOWLEDGEMENTS

I would like to thank Dr. David M. Leslie Jr. for his advice and support throughout this research project. Sandra Pletschet also unselfishly provided invaluable assistance in every phase of this project. In addition, I thank the members of my graduate committee Drs. Tracy Carter and Stanley Fox. I would like to express my gratitude to Frank James, Ron Masters, Scott McMurry, John Skeen, Bill Stuart, and James Whitehead for their help in collecting data reported in this thesis. Special thanks to Don Wood for providing information about the 1977 Red-cockaded Woodpecker survey and to Dr. Claypool for providing statistical advice. This study was supported by Section 6 funds through the Oklahoma Department of Wildlife Conservation.

TABLE OF CONTENTS

Chapter	Page
I. INTRODUCTION.	1
II. THE INFLUENCE OF FORAGING HABITAT QUALITY ON THE DECLINE OF THE RED-COCKADED WOODPECKER IN OKLAHOMA. 2	
Abstract	3
Introduction	5
Study Area	6
Methods.	7
Results.	11
Discussion	13
Acknowledgements	17
Literature cited	18
III. HABITAT ASSOCIATIONS OF ACTIVE RED-COCKADED WOODPECKER CAVITY TREES IN OKLAHOMA	26
Abstract	27
Introduction	28
Study area	30
Methods.	30
Results.	34
Discussion	36
Acknowledgements	41
Literature cited	42

LIST OF TABLES

Table	Page
Chapter II	
1. Comparison of Red-cockaded Woodpecker clans, clusters, and active cavity tree sites located in 1990 with those located in 1977. . . .	22
2. Habitat characteristics in a 510-ha area of McCurtain County Wilderness Area and the relationship (r^2) of each variable to the distance from the nearest active cluster. . . .	23
3. Comparison of a 510-ha study area in McCurtain County Wilderness Area with home ranges of 18 clans in the Francis Marion National Forest and foraging habitat requirements stipulated in the Red-cockaded Woodpecker Recovery Plan	24
Chapter III.	
1. Comparison of hardwood variables between cluster and non-use site base tree plots using Wilcoxon's signed ranks test.	46
2. Comparison of pine variables between cluster and non-use site base tree plots using Wilcoxon's signed ranks test.	47
3. Comparison of basal area of pines and hardwoods between cluster and non-use site 31.3 and 71.3 m sample points.	48
4. Comparison of pine and hardwood basal area between random sample points and cluster site sample points.	49

LIST OF FIGURES

Figure	Page
Chapter II	
1. Locations of active clusters and 510-ha study area within the McCurtain County Wilderness Area	25
Chapter III	
1. Base-tree plot sampling design used at 18 cluster and non-use sites.	50

CHAPTER I

INTRODUCTION

This thesis is composed of two manuscripts written for submission to the Auk. Each manuscript is complete as written and does not require additional support material. Chapter I is an introduction to the rest of the thesis. Chapter II is entitled "The influence of foraging habitat quality on the population decline of the Red-cockaded Woodpecker in Oklahoma." Chapter III is entitled "Habitat associations of active Red-cockaded Woodpecker cavity trees in Oklahoma."

CHAPTER II

THE INFLUENCE OF FORAGING HABITAT QUALITY ON
THE POPULATION DECLINE OF THE RED-COCKADED
WOODPECKER IN OKLAHOMA.

Jeffrey F. Kelly

Oklahoma Cooperative Fish and Wildlife Research Unit,
Department of Zoology, Oklahoma State University,
Stillwater, OK 74078

ABSTRACT.--We searched 4,600 ha of the McCurtain County Wilderness Area, Oklahoma, in 1989-90 and located 15 Red-cockaded Woodpecker (Picoides borealis) clans. In 1977, 83% (3,795 ha) of this area was searched, and 29 Red-cockaded Woodpecker clans were found (Wood 1977). There was a 62% decline (from 29 to 11) in the number of Red-cockaded Woodpecker clans and a 74-76% decline in the number of individuals (from 86-92 to 22) in the resurveyed area (3,795 ha) since 1977. Within the resurveyed area, we found only 3 of 22 clusters of cavity trees (active and abandoned) >400 m from a cluster that was active in 1977 (Wood, unpubl. data), which indicated a low rate of colonization of new areas. The productivity of the population was low during the study period; only 0.69 young per nesting attempt were fledged.

We measured vegetative characteristics in a 510-ha study area to assess quality of foraging habitat. Vegetative characteristics measured were basal area, DBH (diameter at 1.4 m above the ground), number of pines ≥ 25.4 cm DBH/ha, and number of midstory stems (<12.7 cm DBH). Linear regression was used to examine the relationship between foraging habitat quality and distance from the nearest active clusters. The distance from the sample points to the nearest active cluster did not explain a large portion of the variation in any of the vegetation variables for pines or hardwoods. Overall, foraging habitat in the McCurtain County Wilderness Area meets requirements established in the

Red-cockaded Woodpecker Recovery Plan (USFWS 1985).

Geographic isolation and competition are postulated as causes of the decline in population abundance.

The Red-cockaded Woodpecker (Picoides borealis) is endemic to mature pine forests of the southeastern United States and has been listed as an endangered species since 1970. The decline of this species has been caused by loss and fragmentation of mature pine forests (USFWS 1985). Aside from documenting decreasing abundance, little research has been done on small (<25 clans) isolated populations of Red-cockaded Woodpeckers (Ortego and Lay 1988, Eddleman and Clawson 1987, Baker 1983, Thompson 1976). We have found no published instances where small isolated populations have maintained or increased in abundance.

The McCurtain County Wilderness Area in southeastern Oklahoma contains a small isolated population of Red-cockaded Woodpeckers (Wood 1977, Masters et al. 1989). Timber harvesting has never occurred in McCurtain County Wilderness Area; however, private timber lands managed on an even-age, short-rotation basis (<60 yrs) surround the wilderness area. The nearest population of Red-cockaded Woodpeckers from the wilderness area is in the Ouachita National Forest which is 40 km to the northeast (Burnside 1983).

A 1977 survey of 3,795 ha of the wilderness area located 29 clans and 86-92 individuals (Wood 1977). In 1985, the area within 400 m of these 29 active clusters was resurveyed for Red-cockaded Woodpecker activity; 14 active clusters were found in this area (Masters et al. 1989). Fires in the

wilderness area have been suppressed since 1926 (Carter 1967). Concern has been expressed that fire suppression in the wilderness area has or will preclude pine regeneration and result in a hardwood-dominated forest that is unsuitable for the Red-cockaded Woodpecker (Wood 1977, Masters et al. 1989).

The objectives of our research on the Red-cockaded Woodpecker in McCurtain County Wilderness Area were to: (1) assess the current status of the population; (2) determine if foraging habitat was limiting the population abundance; and (3) assess the overall condition of the foraging habitat.

STUDY AREA

The McCurtain County Wilderness Area is a 5,700-ha shortleaf pine (*Pinus echinata*)-mixed hardwood forest located in southeastern Oklahoma. Pines are most abundant on south facing slopes (Carter 1967, Masters et al. 1989, pers. obs.); ridge tops and north facing slopes are dominated by hardwoods, primarily oaks (*Quercus sp.*) and hickories (*Carya sp.*). The terrain is rolling to steep, and elevations range from 183 m to 439 m (Masters et al. 1989). The wilderness area has been owned by the State of Oklahoma since 1917 and managed by the Oklahoma Department of Wildlife Conservation since 1926 (Carter 1967).

METHODS

Clusters.--We used the term clan to refer to a group of Red-cockaded Woodpeckers that foraged together and used closely associated cavity trees; none of the cavity trees located were >300 m from the nearest cavity tree of the same cluster. Clan also referred to single birds that were not associated with other birds. A cavity tree was any tree that contained ≥ 1 Red-cockaded Woodpecker cavity. An active cluster referred to the cavity tree(s) currently used by a clan (Walters et al. 1988 and Rudolph et al. 1990) A group of cavity trees that were not currently being used by a clan were termed an abandoned cluster.

In 1989-90, 4,600 ha of the McCurtain County Wilderness Area were searched along transects placed at 60-m intervals. All pine trees within 30 m of the transects were examined for Red-cockaded Woodpecker cavities (Fig. 1). This technique was identical to the primary technique use by Wood (1977). All of the 3,795 ha searched by Wood in 1977 were resurveyed. An additional 120 ha in the McCurtain County Wilderness Area were assumed to contain no active clusters because infrared aerial photographs revealed that this area contained few pines and therefore, was not searched. Tape recordings of Red-cockaded Woodpecker calls were played intermittently throughout the searched area to attract birds.

When an active cluster was located, the number of birds in the clan was determined by watching the cavity trees at dawn, dusk, or during nesting on ≥ 3 occasions in 1989 and ≥ 6 occasions in 1990. Clan size censuses were conducted in both years for clusters located in 1989. Nests were located by observing bird behavior and by listening for nestlings begging for food. Clans were observed weekly to determine the presence of a nest throughout the nesting season (15 April - 1 July). Nests were monitored on a weekly basis until young fledged or the nest failed. The number of fledglings was determined by monitoring each clan after completion of nesting. Fledglings were easily distinguished by their appearance, awkward bark-scaling, and begging behavior (Jackson 1983).

At each cavity tree we measured: (1) number of cavities; (2) tree diameter at 1.4 m above the ground (DBH); (3) tree height; (4) height of lowest limb; (5) cavity height; (6) cavity orientation; (7) slope; and (8) aspect. Heights and slopes were measured with a clinometer, and diameter was measured with a DBH tape.

Foraging habitat measurements.--Vegetative characteristics were measured at 127 points on 12 randomly located transects within a 510-ha study area (Fig. 1). Points along transects were spaced at 80-m intervals; transects contained from 3-19 points. The sample points ranged from 50 to 1650 m from the nearest active cluster ($\bar{x} = 560 \pm 412$ m, $n = 127$). At each

point, we measured basal area and DBH of pines and hardwoods (>12.7 cm DBH) with a 10-factor prism (Grosenbaugh 1952, Hovind and Rieck 1970) and counted stems (<12.7 cm DBH) >1 m tall within 3.6 m of the point. Classification by DBH was used to separate trees that were potential foraging habitat (>12.7 cm DBH) from those that were too small to be important for foraging and may be selected against because of Red-cockaded Woodpeckers affinity for forests with an open midstory (USFWS 1985, Porter and Labinsky 1986).

To assess overall quality of the foraging habitat, vegetative characteristics that we measured were compared to suitable foraging habitat parameters as defined in the Red-cockaded Woodpecker Recovery Plan (USFWS 1985). We also compared our observations to the habitat within the home range of 18 Red-cockaded Woodpecker clans in the Francis Marion National Forest (Hooper and Harlow 1986).

We hypothesized that if poor quality foraging habitat was responsible for the decline in the number of Red-cockaded Woodpeckers in the McCurtain County Wilderness Area, the remaining active clusters would be located near high quality foraging habitat and the quality of the habitat would decline further from colonies. We defined high quality foraging habitat as having: (1) more basal area of pines and less basal area of hardwoods; (2) pines with larger DBH and hardwoods with smaller DBH; (3) more large pines (>25.4 cm DBH); and (4) fewer midstory stems (<12.7 cm DBH)

particularly hardwood stems (USFWS 1985, Hooper and Harlow 1986, Porter and Labinsky 1986).

Statistical Analysis.--SYSTAT (version 2) was used to conduct all analyses (Wilkinson 1989). Linear regression analysis was used to determine relationships between the distribution of quality foraging habitat and the distance of the sample points from the nearest active cluster. Bartlett's test for homogeneity of variance was used to test for differences in variance in samples with ≥ 30 observations. t-tests were used to compare cavity tree characteristics when sample sizes were ≥ 30 . When variances were unequal, a separate variance t-test was employed (Sokal and Rohlf 1981). Mann-Whitney U-tests were used for comparisons of samples with < 30 observations. All deciduous tree species were grouped for analysis.

Cavity orientation and aspect data were analyzed by breaking down each angle (orientation) into its sine and cosine vectors. The mean sine and cosine vectors for all the orientations were calculated. The angle characterized by the mean sine and cosine was used as the mean orientation. The mean sine and cosine were squared and summed; the square root of the sum was a measure of distribution (r) of the data around the mean, which was tested against a random distribution (Batschelet 1981).

RESULTS

Population status.--Fifteen active Red-cockaded Woodpecker clans were located (Fig. 1); 11 of them (73%) within the 3,795 ha surveyed in 1977. This represented a 62% decline (29 to 11) in the number of clans in the 1977 survey area. Each of the 15 clans used one active cluster. Of the 11 active clusters in the resurveyed area, 10 were within 400 m of an active cluster located in 1977. In addition, 10 of 12 abandoned clusters (83%) located were within 400 m of an active cluster location reported by Wood (1977). This suggests that Red-cockaded Woodpeckers in the McCurtain County Wilderness Area have not frequently formed new clans or established clusters in new locations, and are restricted in the areas that they use.

The modal clan size was 2 ($n = 15$) with a range from 1-4 individuals. A total of 31 individuals was found. Of these, 22 individuals were within the resurveyed area, which represented a 74-76% decline. Birds were never located in areas without active cavity trees; however, a member of a clan had no known roost cavity in 2 cases.

Cavity enlargement by competitors was examined as a possible contributing factor to population decline. Of all cavities that were located (active and abandoned; $n = 64$), 23% had been enlarged by other species. No Red-cockaded Woodpeckers were observed using enlarged cavities. Of the

32 cavities known to be active at some point during this study, 28% (n = 9) were later abandoned. Only 2 of these abandoned cavities were noticeably enlarged. It is not known if enlargement occurred prior or subsequent to abandonment.

During the 1989 nesting season, only 6 clans had been located; 5 of these nested. A single young fledged at 1 nest; the other 4 nests did not fledge young that survived until the time of our next weekly census. Nests were located at 8 of 15 colonies in 1990, and all 8 nests produced a single fledgling. During both years, at least 10 of the 13 nests had ≥ 2 nestlings. However, brood reduction occurred in all successful nests; generally only 1 nestling remained in the nest 1 week prior to fledging. Nine fledglings were produced in these 13 nests, or an average of 0.69 young/nest.

We found clans with fewer individuals and clusters with fewer cavity trees than were reported by Wood (1977) (Table 1). Slope at the cluster was the only other characteristic that differed between 1977 and 1990 (Table 1). Cavity orientation (\bar{x} = 254 degrees, n = 39, r = 0.55, p < 0.001) and aspect (\bar{x} = 231.6 degrees, n = 32, r = 0.42, p < 0.004) found in 1990 had significantly directional distributions. The directional tendency of the aspect may be related to the higher abundance of pines on south facing slopes.

Habitat condition.--There were only 2 significant

relationships between the distance from the nearest active cluster and habitat variables, and only a small percentage of the variation was accounted for (Table 2). We believe that proximity of clusters to quality foraging habitat was not a primary influence on the specific locations of active clusters.

Overall, our study area met foraging habitat requirements stipulated by the Red-cockaded Woodpecker Recovery Plan (USFWS 1985) and contained comparable habitat characteristics to those in the home ranges of Red-cockaded Woodpeckers in the Francis Marion National Forest (Hooper and Harlow 1986) (Table 3). Average DBH and basal area were similar to those found to be preferred foraging habitat for Red-cockaded Woodpeckers in the Apalachicola National Forest (Porter and Labinsky 1986). Despite the suitability of the foraging habitat, the number of clans declined in the sampled area, which suggested that the decline of the Red-cockaded Woodpecker in the McCurtain County Wilderness Area has not been exclusively due to a lack of foraging habitat.

DISCUSSION

Our data indicate that the distribution of quality foraging habitat was probably not a primary factor determining the distribution or abundance of Red-cockaded Woodpeckers in the McCurtain County Wilderness Area. Other factors have the potential to limit the population; e.g.,

availability of suitable trees for cavity construction, which can be addressed indirectly. Characteristics of cavity trees in the McCurtain County Wilderness Area in 1990 were similar to those reported from the same area in 1977 (Table 1). Also, except for greater cavity height and tree age, cavity tree characteristics in 1977 and 1990 were similar to those reported elsewhere (Ligon 1970, Baker 1971, Hopkins and Lynn 1971, Hooper 1988). Between 1989 and 1990, 6 cavities were abandoned in 5 active clusters; however, we located 5 new cavity trees in these 5 clusters in 1990. In addition, 94% (29 of 31) of the birds that we located had known roost cavities. These factors suggested that availability of trees that could be used for cavities was probably not limiting in the areas where colonies were located. Also, the number of pines/ha ≥ 25.4 cm DBH throughout the foraging area indicated that the number of trees of the size suitable for cavity construction was not detrimental to the Red-cockaded Woodpecker population (Table 2).

The trend of forest succession in the wilderness area could not be assessed directly because comparable vegetation measurements from 1977 were not available for comparison. However, our habitat measurements demonstrate a relatively dense midstory that is dominated by hardwoods (Table 2). The dominance of hardwoods in the midstory probably indicates a successional shift from pine dominance to

hardwood dominance due to fire suppression in the wilderness area (Cain 1987). Previous research has indicated that high hardwood density decreases the suitability of forests for Red-cockaded Woodpeckers (Lennartz et al., 1983, USFWS 1985, Hooper and Harlow 1986). Unfortunately, the specific level at which hardwood density becomes problematic for Red-cockaded Woodpeckers is unknown (USFWS 1985).

Productivity of Red-cockaded Woodpeckers in the McCurtain County Wilderness Area (0.69 fledglings/nest) was lower than has been reported elsewhere. Average productivity in coastal South Carolina was 1.8 fledglings/nest/year (USFWS 1985); in Florida the fledgling/nest average was 1.6 (n = 9 nests) (Ligon 1970). We found that the mean annual number of fledglings produced/breeder was 0.35 compared to the 0.74 reported by Reed et al. (1988) in the North Carolina Sandhills. This disparity in productivity may be caused by several factors, such as inbreeding due to the low number of individuals in the populations, low resource availability, competition, and/or predation.

The amount of gene flow from outside populations and its consequences for genetic fitness are important considerations in examining the decline of the Red-cockaded Woodpecker in the McCurtain County Wilderness Area. Theoretical estimates of minimal viable population size range from 500 to 1,018 individuals (USFWS 1985, Reed et al. 1988). Red-cockaded Woodpeckers dispersing into the

wilderness area from the nearest population (the Ouachita National Forest) would have to travel 40 km, which is further than the maximum (31.5 km) and 3.4-8.8 times the mean dispersal distance (depending on the age, sex and status of the individual) reported by Walters et al. (1988). Lack of immigration and isolation were suggested as causes of the extirpation of a Red-cockaded Woodpecker population that was separated from a larger population by only 10 km (Baker 1983). Dispersal problems would be expected to increase as the distance between clans within a population and distance between populations increases.

The effect of population density on reproductive success of the population needs to be addressed. Most research that has been conducted on the Red-cockaded Woodpecker has focused on relatively large populations in forests managed for timber harvest. Factors that limit the abundance of large populations (i.e., amount of suitable habitat) may not be as important in populations that have declined below a density where dispersing individuals can easily locate mates. Clan density probably influences ability of dispersing Red-cockaded Woodpeckers to locate mates or clans other than the natal clan. Low clan density could lead to increased mortality of dispersers and increased probability of inbreeding within clans of close proximity. Dispersing males have considerably lower survival than helpers in areas where populations are relatively large (Walters et al.

1988). A combination of low survival rate among dispersers, poor reproductive success, and few dispersing individuals (of both sexes due to population size) may limit the possibility of forming new colonies and explain why 10 of 11 colonies located in the resurveyed area were within 400 m of colonies sites identified in 1977.

Cavity competition from Pileated (Dryocopus pileatus) and Red-bellied woodpeckers (Melanerpes carolinus) and flying squirrels (Glaucomys volans) also may contribute to the population decline of Red-cockaded Woodpeckers in McCurtain County Wilderness Area. Previous research has not found cavity competition from Pileated Woodpeckers to be important (Rudolph et al. 1990, Harlow and Lennartz 1983); however, McCurtain County Wilderness Area is predominantly a mixed forest and Pileated Woodpeckers are abundant (pers. obs.). The influence of cavity competition on individual clans may be significant. The frequency and overall impact of enlargement of active Red-cockaded Woodpecker cavities in McCurtain County Wilderness Area is unknown.

ACKNOWLEDGEMENTS

This study was supported by Section 6 funds through the Okla. Dep. of Wildl. Cons. We thank F. James, S. McMurry, R. Masters, B. Stivers, J. Skeen, B. Stuart, and J. Whitehead for help collecting data. We also are grateful to D. Wood for information concerning the 1977 survey.

LITERATURE CITED

- Baker, W.W. 1971. Progress report on life history studies of the Red-cockaded Woodpecker at Tall Timbers Research Station. Pp. 44-59 in Symposium on the Red-cockaded Woodpecker. (R.L. Thompson, Ed.). Tallahassee, Florida, Bur. Sport Fish. Wildl., U.S. Dept. Interior, and Tall Timbers Research Station.
- _____. 1983. Decline and extirpation of a population of Red-cockaded Woodpeckers in northwest Florida. Pp. 44-45 in Proc. Red-cockaded Woodpecker Symp.II. (D.A. Wood, Ed.). Gainesville, Florida. Game Fresh Water Fish Comm., U.S. Fish Wildl. Serv. and U.S. For. Serv.
- Batschelet, B. 1981. Circular statistics in biology. London, England, Academic Press.
- Burnside, F.L. 1983. The status and distribution of the Red-cockaded Woodpecker in Arkansas. Am. Birds 37(2):142-145.
- Cain, M.D. 1987. Survival patterns of understory woody species in a pine-hardwood forest during 28 years without timber management. Pp. 141-147 in Proc. of the central hardwood forest conference VI. (R.L. Hay, F.W. Woods, & H. DeSelm, Ed.). Knoxville, Tennessee.
- Carter, W.A. 1967. Ecology of the nesting birds of McCurtain Game Preserve, Oklahoma. Wilson Bull. 79:259-272.

- Eddleman, W.R., & R.L. Clawson. 1987. Population status and habitat conditions for the Red-cockaded Woodpecker in Missouri. *Trans. Missouri Acad. Sci.* 21:105-117.
- Grosenbaugh, L.R. 1952. Plotless timber estimates, new, fast, easy. *J. For.* 50:32-37.
- Harlow, R.F., & M.R. Lennartz. 1983. Interspecific competition for Red-cockaded Woodpecker cavities during the nesting season in South Carolina. Pp. 41-43 in *Proc. of Red-cockaded Woodpecker Symp. II.* (D.A. Wood, Ed.). Gainesville, Florida, Game Fresh Water Fish Comm., U.S. Fish Wildl. Serv. and U.S. For. Serv.
- Hooper, R.G. 1988. Longleaf pines used for cavities by Red-Cockaded Woodpeckers. *J. Wildl. Manage.* 52:392-398.
- _____, & R.F. Harlow. 1986. Forest stands selected by foraging Red-cockaded Woodpeckers. *Res. Pap. SE-259.* Asheville NC: USDA, For. Serv., Southeastern Forest Experiment Station. 10 Pp.
- Hopkins, M.L., & T.E. Lynn. 1971. Some characteristics of Red-cockaded Woodpecker trees and management implications in South Carolina. Pp. 140-169 in *Symposium on the Red-cockaded Woodpecker.* (R.L. Thompson, Ed.). Tallahassee, Florida, Bur. Sport Fish. Wildl., U.S. Dept. Interior and Tall Timbers Research Station.
- Hovind, H.J., & C.E. Rieck. 1970. Basal area and point sampling interpretation and application. Wisconsin Dept. of Nat. Res. *Tech. Bull. no. 23.*

- Jackson, J.A. 1983. Morphological and behavioral Development of post-fledging Red-cockaded Woodpeckers. Pp. 30-38 in Proc. Red-cockaded Woodpecker Symp. II (D.A. Wood, Ed.). Gainesville, Florida Game Fresh Water Fish Comm., U.S. Fish Wildl. Serv. and U.S. Forest Serv.
- Lennartz, M.R., H.A. Knight, J.P. McClure, & V.A. Rudis. 1983. Status of Red-cockaded Woodpecker nesting habitat in the south. Pp. 13-19 in Proc. Red-cockaded Woodpecker Symp. II. (D.A. Wood, Ed.). Gainesville, Florida Game Fresh Water Fish Comm., U.S. Fish Wildl. Serv. and U.S. For. Serv.
- Ligon, J.D. 1970. Behavior and breeding biology of the Red-cockaded Woodpecker. Auk 87:255-278.
- Masters, R.E., J.E. Skeen, & J.A. Garner. 1989. Red-cockaded Woodpecker in Oklahoma: an update of Wood's 1974-77 study. Proc. Okla. Acad. Sci. 69:27-31.
- Ortego, B., & D. Lay. 1988. Status of Red-cockaded Woodpecker colonies on private land in east Texas. Wildl. Soc. Bull 16:403-405.
- Porter, M.L., & R.F. Labinsky. 1986. Home range and foraging habitat of Red-cockaded Woodpeckers in northern Florida. J. Wildl. Manage. 50:239-247.
- Reed, J.M., P.D. Doerr, & J.R. Walters. 1988. Minimum viable population size of the Red-cockaded Woodpecker. J. Wildl. Manage. 52:385-391.

- Rudolph, D.C., R.N. Conner, & J. Turner. 1990. Competition for Red-cockaded Woodpecker roost and nest cavities: effects of resin age and entrance diameter. *Wilson Bull.* 102:23-36.
- Sokal, R.R., & F.J. Rohlf. 1981. *Biometry*. San Francisco, California, W.H. Freeman and Company.
- Thompson, R.L. 1976. Changes in status of Red-cockaded Woodpecker colonies. *Wilson Bull.* 88:491-492.
- U.S. Fish and Wildlife Service. 1985. Red-cockaded Woodpecker Recovery Plan. U.S. Fish and Wildl. Serv. Atlanta, Georgia.
- Walters, J.R., P.D. Doerr, & J.H. Carter III. 1988. The cooperative breeding system of the Red-cockaded Woodpecker. *Ethology* 78:275-305.
- Wilkinson, L. 1989. *SYSTAT: The system for statistics*. Evanston Illinois, SYSTAT Inc.
- Wood, D.A. 1977. Status, habitat, home range, and notes on the behavior of the Red-cockaded Woodpecker in Oklahoma. Stillwater, Oklahoma, Oklahoma State Univ. Master's thesis.

Table 1. Comparison of Red-cockaded Woodpecker clans, clusters, active cavity trees, and active cavity tree sites located in 1990 with those located in 1977. Sample sizes are in parentheses.

	1990	1977 ^a	Test	
	$\bar{X} \pm SD$	$\bar{X} \pm SD$	statistic	P
Clan				
Number of birds	2.1 \pm 0.8 (15)	3.1 \pm 1.2 (29)	U = 112	0.008
Cluster				
Number of trees	1.9 \pm 0.7 (15)	4.4 \pm 2.0 (29)	U = 57	0.001
Cavity tree				
Height(m)	25.7 \pm 4.8 (33)	24.9 \pm 4.1 (154)	t = 0.98	0.329
Diameter(cm)	43.9 \pm 6.3 (32)	44.6 \pm 8.2 (154)	t = 0.39	0.693
Low limb height(m)	9.6 \pm 3.6 (32)	10.7 -- ^b (154)	--	--
Cavities/tree	1.3 \pm 0.6 (32)	1.2 -- (155)	--	--
Cavities				
Height(m)	12.0 \pm 3.0 (32)	13.1 -- (224)	--	--
Orientation(degrees)	260.0 \pm 55.0 (32)	251.1 \pm 56.7 (205)	--	--
Site				
Slope(percent)	12.0 \pm 4.9 (31)	26.0 \pm 12.8 (153)	t = 5.27	0.001
Aspect(degrees)	232.0 \pm 62.0 (32)		--	--

^a Data from Wood (1977)

^b Data as collected in 1977 precluded analysis

Table 2. Habitat characteristics in a 510-ha area of McCurtain County Wilderness Area and the relationship (r^2) of each variable to the distance from the nearest active cluster. All sample sizes = 127

Habitat variable	$\bar{X} \pm SD$	r^2	P
Basal Area (m²/ha)			
Pine	10.8 \pm 7.6	0.006	0.37
Hardwood	7.4 \pm 4.8	0.028	0.06
DBH (cm)			
Pine	29.9 \pm 15.1	0.018	0.13
Hardwood	24.0 \pm 12.3	0.008	0.33
Number/ha >25.4 cm DBH			
Pine	58.8 \pm 47.6	0.010	0.25
Number /ha <12.7 cm DBH			
Pine	281.7 \pm 867.2	0.033	0.04
Hardwood	1487.0 \pm 1194.4	0.079	0.01

Table 3. Comparison of a 510 ha study area in McCurtain County Wilderness Area (MCWA) with home ranges of 18 clans in the Francis Marion National Forest (FMNF) and the foraging habitat requirements stipulated in the Red-cockaded Woodpecker recovery plan (RCWRP) (Hooper and Harlow 1986, USFWS 1985). Sample sizes are in parentheses.

	MCWA	FMNF	RCWRP
	$\bar{X} \pm SD$	$\bar{X} \pm SD$	
Pine			
Basal area (m ² /ha)	10.8 ± 7.6 (127)	11.5 ± 7.7 (276)	6.7-20.7
Percent	59.0 --- ---	67.3 --- ---	>50.0
Number/ha >25.4 cm DBH	58.8 ± 47.6 (127)	93.4 ± 72.1 (276)	>59.3 ^a
Hardwood			
Basal area (m ² /ha)	7.5 ± 4.9 (127)	5.6 ± 9.4 (276)	0.0-10.4
Percent	41.0 --- ---	32.7 --- ---	<50.0
Total			
Basal area (m ² /ha)	18.3 ± 7.6 (127)	17.1 --- (276)	13.8-20.7

^a Red-cockaded Woodpecker Recovery Plan stipulates 59.3 Pines/ha >24.0 cm DBH rather than 25.4 cm DBH




Figure 1. Locations of active clusters, the area used to measure foraging habitat, and the area surveyed for active clusters in the McCurtain County Wilderness Area, Oklahoma.

CHAPTER III

HABITAT ASSOCIATIONS OF ACTIVE RED-COCKADED
WOODPECKER CAVITY TREES IN OKLAHOMA

Jeffrey F. Kelly

Oklahoma Cooperative Fish and Wildlife Research Unit,
Department of Zoology, Oklahoma State University,
Stillwater, OK 74078

ABSTRACT.--We compared habitat at 18 active Red-cockaded Woodpecker (Picoides borealis) cavity trees (cluster sites) and 18 paired non-use sites in McCurtain County Wilderness Area, Oklahoma. The area surrounding active cavity trees (0.04 ha) had significantly shorter hardwoods and less hardwood basal area than non-use sites. Within cluster sites, the 0.01-ha quarter directly in front of the cavity opening also had significantly shorter hardwoods, hardwoods with smaller DBHs, and less hardwood basal area than the remainder of the area surrounding the cavity tree. In addition, basal areas of pines 31.3 m from cluster sites was significantly higher than randomly sampled habitats, but there was no significant difference in the density of hardwoods between these 2 samples. Hardwood density at cluster sites was greater than recommended by the Red-cockaded Woodpecker Recovery Plan but was comparable to hardwood densities documented elsewhere in the range of the Red-cockaded Woodpecker. Our observations suggest that Red-cockaded Woodpeckers selected habitat on at least 2 levels; low hardwood densities were selected for cavity sites and high pine densities were selected for foraging habitat.

Red-cockaded Woodpeckers (Picoides borealis) excavate cavities exclusively in mature southern pines (e.g., Pinus palustris, Pinus taeda, Pinus echinata). Cavities are usually placed in trees >60 years old if they are available (USFWS 1985, Conner and O'Halloran 1987). Several studies have described characteristics of cavity trees and investigated causes of cavity abandonment (Hopkins and Lynn 1971, Teitelbaum and Smith 1985, Conner and O'Halloran 1987, Hooper 1988). Size and age of pines and density of hardwood midstory have been proposed as the primary factors that influence placement of cavities by Red-cockaded Woodpecker (Van Balen and Doerr 1978, Locke et al., 1983, USFWS 1985, Hovis and Labinsky 1985, Conner and O'Halloran 1987, Conner and Rudolph 1989). Red-cockaded Woodpeckers select cavity trees that are surrounded by relatively open forests (within 200 m) (Hovis and Labinsky 1985). Some evidence exists that Red-cockaded Woodpeckers select cavity trees that have sparse hardwood midstory within several meters and that the midstory nearest the cavity entrance differs from the remainder of the midstory in the vicinity of the cavity tree (Van Balen and Doerr 1978). However, the importance and differences in habitat selection at different levels (i.e., distance from a prospective cavity tree) have not been investigated.

The number of Red-cockaded Woodpeckers in the McCurtain County Wilderness Area, Oklahoma, has declined by 75%

between 1977 and 1990, and there were only 15 known clans and 31 individuals in 1990 (Kelly et al. 1991).

Availabilities of pines of sufficient size in which to construct cavities and quality foraging habitat probably did not cause this decline (Kelly et al. 1991). Fires have been suppressed in McCurtain County Wilderness Area since 1926. Because of this, encroachment of hardwood midstory on active cavities has been proposed as the cause for the decline in the number of Red-cockaded Woodpeckers (Masters et al. 1989).

To examine habitat selection and the influence of hardwood encroachment on active cavities of Red-cockaded Woodpeckers in the wilderness area, we measured vegetative characteristics around cavity trees and compared them to the habitat around randomly chosen paired non-use sites (without cavities) and randomly placed sites. Our objectives were: (1) to describe the habitat at active cavities in the McCurtain County Wilderness Area and compare this habitat to habitat descriptions from elsewhere in the Red-cockaded Woodpeckers range; and (2) to compare habitat at cavity trees to that of non-use sites and randomly placed sites and thereby assess habitat preferences of Red-cockaded Woodpeckers.

STUDY AREA

Our study was conducted in the 5,700-ha McCurtain County Wilderness Area in southeastern Oklahoma. The area was dominated by shortleaf pine (*Pinus echinata*)-mixed hardwood forest. Pines were most abundant on south facing slopes (Carter 1967, Masters et al. 1989, pers. obs.); ridgetops were dominated by hardwoods, primarily oaks (*Quercus* spp.) and hickories (*Carya* spp.). The terrain was rolling to steep and elevations ranged from 183 to 439 m (Masters et al. 1989). The wilderness area was purchased by the State of Oklahoma in 1917 and has been managed by the Oklahoma Department of Wildlife Conservation since 1926.

METHODS

We defined a clan as a group of Red-cockaded Woodpeckers that foraged together and used closely associated cavity trees; individuals in the same clan always used cavity trees within 300 m of each other. Clan also referred to single birds. The term cluster referred to the cavity trees used by a clan. Cavity trees were determined to be active by observing roosting and nesting behavior of clans. Vegetative characteristics were measured in 15 active clusters at 18 active cavity trees. At the 15 active clusters, an active cavity tree was selected to be the plot center for habitat measurements (base trees) if it had been

used as a nest tree ($n = 2$). If the nesting history was unknown at a particular cluster, a base tree was chosen at random ($n = 13$). At 3 clusters, there was an active cavity tree >180 m from the initially selected base tree; these 3 trees were measured as independent base tree sites. The DBH (diameter at 1.4 m above the ground) of each base tree and slope and aspect of each site were measured with a DBH tape, a SUUNTO clinometer, and a hand held compass, respectively. Circular 0.04-ha plots were established with the base tree at the center (Fig. 1). Within each base-tree plot, we measured height and DBH of all plants ≥ 1 m tall and ≥ 5.0 cm DBH and counted saplings (>1 m tall and <5.0 cm DBH). All species of deciduous trees were grouped as "hardwoods" for analysis. Each plot was divided into 4 quadrants; the direction of cavity orientation determined the midpoint of quadrant 1 (Fig. 1). Foliage cover was estimated using a forest densiometer at the point where each quadrant line intersected the 0.04-ha circular plot boundary. Basal area measurements were taken with a 10-factor prism at 31.3 m and 71.3 m from the center of the base tree along the quadrant boundary lines (Hovind and Rieck 1970).

Active and non-use sites were paired by the DBH of the base tree and slope and aspect of the site. Non-use sites were randomly located on a topographic map in areas that were: (1) within the 4,600 ha that had been searched for clusters (Kelly et. al 1991); (2) >500 m from active

clusters; and (3) >500 m from the boundary of the searched area. These criteria were used to eliminate the possibility that the non-use sites were being used by Red-cockaded Woodpeckers. We located these random points in the field and then found the site closest to the random point that had a pine within 5 cm DBH of the base tree and a slope within 3 degrees and aspect within 10 degrees of the paired cluster site. If no suitable site was found within 200 m of the random point, another point was chosen. Tape recordings of Red-cockaded Woodpeckers were played intermittently over a 3-hour period at non-use sites to ensure that the area was not currently being used by Red-cockaded Woodpeckers. Orientation of quadrant directions at each non-use site base tree were the same as quadrant directions at the paired cluster site base tree. Comparisons were made among quadrants within cluster sites to identify habitat differences that were related to cavity orientation. Identical comparisons were made among quadrants of non-use sites to determine if site characteristics alone could account for differences found among quadrants in cluster sites. Cluster sites were compared to non-use sites by quadrant and overall.

Basal area point samples taken at 31.3 m from the cluster site base trees along the quadrant boundary lines were compared to 127 randomly located basal area point samples that were measured in another study to evaluate foraging

habitat (Kelly et al. 1991). Trees <12.7 cm DBH were eliminated from the cluster site samples so that the techniques used in collecting the 2 samples were identical.

Comparisons between and within cluster and non-use sites were made using SYSTAT computer software (Wilkinson 1989). Comparison among quadrants within sites were made with Friedman's 2-way ANOVA (Conover and Iman 1981). Wilcoxon's signed ranks tests were used to make comparisons between cluster and non-use sites for each habitat variable (Sokal and Rohlf 1981). Because 5 Wilcoxon's tests were conducted for each habitat variable, a sequential Bonferroni correction was employed to control Type 1 error (Rice 1989). An initial critical P value of 0.01 (0.05/5 tests) was used to indicate significance. If none of the 5 tests had a P value <0.01 then none were significant. If 1 of the 5 tests had a P value <0.01 then it was significant and the critical P value was adjusted to 0.0125 (0.05/(5-1 tests)) and the remaining 4 P values were reexamined to determine if they were <0.0125. The critical P value was increased each time a test was significant (i.e., for the 5th test the critical P value was 0.05 if the previous 4 tests were significant). Discriminant function analysis was used to distinguish between cluster and non-use sites. The comparisons between 31.3 m basal area point samples and basal area point samples from Kelly et al. (1991) were made with t-tests.

RESULTS

Within site comparisons.--No significant differences were found in the number of pine saplings, DBH of pine trees, basal area of pines, or height of pines among quadrants in either cluster or non-use sites. Quadrant 1 of cluster sites had shorter hardwoods than the other 3 quadrants (Friedman's statistic = 14.5, $P = 0.002$, $df = 3$; Sum of ranks: quadrant 1 = 28, quadrant 2 = 54, quadrant 3 = 53, quadrant 4 = 45). The difference in the height of hardwoods between quadrant 1 and the other 3 quadrants in cluster sites indicated that Red-cockaded Woodpeckers selected areas with shorter hardwoods in which to orient their cavity openings. At non-use sites, quadrant 1 also had shorter hardwoods (Friedman's statistic = 9.5, $P = 0.023$, $df = 3$; Sum of Ranks: quadrant 1 = 32, quadrant 2 = 44, quadrant 3 = 55, quadrant 4 = 49), which suggested that shorter hardwoods may be inherent to the cavity orientations and type of sites (i.e., slope, aspect, and tree size) selected by Red-cockaded Woodpeckers. There were no significant differences in number of hardwood saplings, hardwood DBH, or hardwood basal area among quadrants within either cluster or non-use sites. Forest cover (measured with a densiometer) did not differ significantly among quadrants within sites. Basal area of pines and hardwoods 31.3 m and 71.3 m from the base tree did not differ significantly among quadrants within

either cluster or non-use sites.

Between site comparisons--Overall (i.e., in all 4 quadrants combined), cluster sites had less hardwood basal area and shorter hardwoods than non-use sites. Quadrant 1 was the most disparate between the cluster and non-use sites (Table 1). Additionally, hardwoods in quadrant 3 were significantly shorter in cluster sites than in non-use sites. There was significantly less foliage cover in cluster sites ($82.1 \pm 4.1\%$, $n = 18$) than in non-use sites ($84.2 \pm 4.9\%$, $n = 18$, $Z = 2.44$, $P = 0.015$). Although this difference was small and it is unclear if it is biologically significant, it may reflect smaller, sparser hardwoods at cluster sites than at non-use sites. The shorter, sparser hardwoods in quadrant 1 and cluster sites as a whole indicated selection against hardwoods near cavity trees by Red-cockaded Woodpeckers.

Overall and in quadrant 1, cluster sites had significantly more pine saplings than non-use sites (Table 2). There were no other significant differences in pines between cluster and non-use site, base-tree plots. The higher pine density, particularly the number of saplings, in cluster sites compared to non-use sites was probably due to the open (in terms of hardwoods) character of the forest selected for cluster sites.

Discriminant function analysis using basal area, average height, and average DBH of hardwoods in quadrant 1 correctly

classified 28 of the 36 sites (78%) as either cluster or non-use ($X^2 = 48.4$, $df = 4$, $P < 0.001$), which further emphasizes the importance of hardwoods in quadrant 1 in selection of cavity sites. There were 4 cluster and 4 non-use sites that were incorrectly classified.

Quadrant 3 at cluster sites had significantly more pine and less hardwood basal area 31.3 m and 71.3 m from the base tree than at non-use sites (Table 3). There was an overall trend toward more pine and less hardwood in cluster sites when compared to non-use sites at both 31.3 and 71.3 m from the base trees. There was significantly more pine basal area 31.3 m from cluster site base trees than at randomly selected sites (Table 4). These 2 samples did not differ significantly in the amount of hardwood basal area. Apparently, pine density is of greater importance than hardwood density in habitat selection 31 m from the cavity tree.

DISCUSSION

Dense hardwood midstory has been proposed as a cause of abandonment of Red-cockaded Woodpecker cavities (Hopkins and Lynn 1971, Hovis and Labinsky 1985, Kalisz and Boettcher 1991). There is some evidence that clusters in areas with high hardwood densities are more likely to be abandoned (Conner 1989). However, the reason why high hardwood density leads to abandonment is unclear. Possible

explanations include: (1) hardwood encroachment on the cavity entrance obstructs the entrance making the cavity unsuitable to Red-cockaded Woodpeckers; (2) hardwood encroachment makes the cavity accessible to competitors and/or predators by allowing them to avoid the resin barrier; (3) dense midstory hardwoods decrease foraging habitat quality resulting in low productivity; and (4) forests with dense midstory hardwoods have denser populations of species that compete with or prey upon Red-cockaded Woodpeckers.

Habitat quality.--Hardwood density in the McCurtain County Wilderness Area was within the range of densities that have been reported elsewhere within the species range (Van Balen and Doerr 1978, Conner 1989, Kalisz and Boettcher 1991). However, hardwoods surrounding Red-cockaded Woodpecker cavity trees in the wilderness area were generally denser (Table 1) than recommended in the Red-cockaded Woodpecker Recovery Plan (4.6 m²/ha, USFWS 1985).

Hardwood encroachment on the cavity entrance is probably not a major threat to the remaining Red-cockaded Woodpecker population in McCurtain County Wilderness Area. Heights of Red-cockaded Woodpecker cavities in McCurtain County Wilderness Area were greater (12-13 m) than have been reported elsewhere (Wood 1977, Kelly et al. 1991), but the gap between the cavity openings and the midstory (>5 m) was similar to that reported elsewhere (Van Balen and Doerr

1978, Kalisz and Boettcher 1991). This suggests that the Red-cockaded Woodpeckers have placed their cavities in areas where hardwood encroachment will not cause abandonment. In addition, average midstory height in unburned shortleaf pine stands has been found to remain constant (8-11.5 m) after the stand reaches 50 years of age (Billings 1938). The height of hardwoods in the wilderness area (Table 1) was generally within the range reported by Billings (1938). Because fires have been suppressed in the wilderness area for 65 years, it is possible that the hardwood midstory has reached a maximum average height and has not increased in height over the last 15 years.

McCurtain County Wilderness Area is one of the few forests that contain Red-cockaded Woodpeckers and has never been logged. Wood (1977) found that average age of cavity trees in the wilderness area was 149 years and average cavity tree age from other forests was 81 years. Cavity trees in the wilderness area may be used for short periods of time because of they are old and have high mortality rates and low resin flow. Cavities that are used for shorter periods of time are less likely to be encroached upon by hardwoods.

In summary, our data suggest that hardwood encroachment is probably not a major threat to the remaining active cavities because of (1) the relationship of cavity height to the height of the hardwoods in quadrant 1, (2) the probable stability in average midstory height, and (3) the relatively

short period that old cavity trees can produce adequate resin. Wood (1983) reached a similar conclusion concerning the impact of midstory encroachment on cavity abandonment during his 1977 study of Red-cockaded Woodpeckers in McCurtain County Wilderness Area.

The primary negative impact of hardwood density on this population of Red-cockaded Woodpeckers is probably a reduction in productivity and potential for expansion caused by (1) limitation of the number of suitable cluster sites, (2) degradation of foraging habitat as a result of succession from a pine to hardwood dominated forest, and/or (3) competition from Red-bellied (Melanerpes carolinus) and Pileated Woodpeckers (Dryocopus pileatus) which are more abundant in dense hardwood than in open pine forests (USFWS 1985, Renken and Wiggers 1989).

Habitat selection.--We suggest that Red-cockaded Woodpeckers in McCurtain County Wilderness Area select habitat on at least 2 levels and use different selection criteria at each of these levels. We found that structure of the hardwoods midstory was most important within 11.3 m of the cavity tree and pine density was more important than hardwood density at 31.3 m from the cavity tree. In general, cavities were constructed in areas where hardwoods were shorter and less dense. Specifically, hardwoods in quadrant 1 were less dense and smaller than in the rest of the quadrants; and overall, habitat in the 0.04 ha

surrounding cluster sites had smaller hardwoods than were found at paired non-use sites. This indicated that selection against hardwoods was occurring in the immediate vicinity (within 11.3 m) of the cavity tree. The difference in the size and amount of hardwoods (particularly height) between quadrant 1 and the other 3 quadrants indicated that the hardwood density immediately in front of the cavity may be more important than overall forest composition in selection of a specific cavity site. Basal area samples taken at 31.3 m from cluster site base trees did not differ from randomly placed basal area samples in the amount of hardwood basal area, suggesting that selection against hardwoods may cease to be important at that distance (Table 4).

The area 31.3 m from the cavity tree is probably important foraging habitat for Red-cockaded Woodpeckers during the breeding season; the high density of pine at these sample points reflects selection of foraging habitat by the resident clan. In addition, the trend of higher pine basal area at cluster sites when compared to non-use sites is consistent with prior observations that Red-cockaded Woodpeckers prefer to forage in areas with relatively high pine densities (Hooper and Harlow 1986, Porter and Labinsky 1986).

While our data do not address availability of habitat within the McCurtain County Wilderness Area, they do suggest

that habitat selection by Red-cockaded Woodpeckers of characteristics may occur on several levels. Large pines were selected for foraging habitat and low hardwood density was selected for cavity sites. It appears that selection against hardwoods by Red-cockaded Woodpeckers is strongest in the area nearest the cavity entrance; this trend also was found by Van Balen and Doerr (1978). Selection against midstory hardwoods in the vicinity of the cavity entrance may prevent cavity abandonment caused by hardwood encroachment even in forests with relatively high hardwood densities.

ACKNOWLEDGEMENTS

This study was supported by Section 6 funds through the Oklahoma Department of Wildlife Conservation. We thank F. James, manager of the McCurtain County Wilderness Area for his assistance in various aspects of this study.

LITERATURE CITED

- Billings, W.D. 1938. The structure and development of old field shortleaf pine stands and certain associated physical properties of the soil. *Ecol. Monogr.* 8:437-499.
- Carter, W.A. 1967. Ecology of the nesting birds of McCurtain Game Preserve, Oklahoma. *Wilson Bull.* 79:259-272.
- Conner, R.N. 1989. Injection of 2,4-D to remove hardwood midstory within Red-cockaded Woodpecker colony areas. Res. Pap. SO-251. New Orleans, LA: USDA, For. Serv., Southern Forest Experiment Station. 4 Pp.
- _____, & K.A. O'Halloran. 1987. Cavity-tree selection by Red-cockaded Woodpeckers as related to growth dynamics of southern pines. *Wilson Bull.* 99:398-412.
- _____, & C.D. Rudolph. 1989. Red-cockaded Woodpecker colony status and trends on the Angelina, Davy Crockett and Sabine National Forests. Res. Pap. SO-251. New Orleans, LA: USDA, For. Serv., Southern Forest Experiment Station. 15 Pp.
- Conover W.J. and R.L. Iman. 1981. Rank transformations as a bridge between parametric and nonparametric statistics. *The Am. Statistician* 35:124-129.

- Hooper, R.G. 1988. Longleaf pines used for cavities by Red-cockaded Woodpeckers. *J. Wildl. Manage.* 52:392-398.
- _____, & R.F. Harlow. 1986. Forest stands selected by foraging Red-cockaded Woodpeckers. Res. Pap. SE-259. Asheville, NC: USDA, For. Serv., Southeastern Forest Experiment Station. 10 Pp.
- Hopkins, M.L., & T.E. Lynn. 1971. Some characteristics of Red-cockaded Woodpecker cavity trees and management implications in South Carolina. Pp. 140-169 in Symposium on Red-cockaded Woodpecker. (R.L. Thompson, Ed.). Tallahassee, Florida, Bur. Sport Fish. Wildl., U.S. Dept. Interior, and Tall Timbers Research Station.
- Hovind, H.J., & C.E. Rieck. 1970. Basal area and point sampling interpretation and application. Wisconsin Dept. of Nat. Res. Tech. Bull. no. 23.
- Hovis, J.A., & R.F. Labinsky. 1985. Vegetative associations of Red-cockaded Woodpecker colonies in Florida. *Wildl. Soc. Bull.* 13:307-314.
- Kalisz, P.J., & S.E. Boettcher. 1991. Active and abandoned Red-cockaded Woodpecker habitat in Kentucky. *J. Wildl. Mange.* 55:146-154.
- Kelly, J.F., S.M. Pletschet, & D.M. Leslie Jr. 1991. Influence of foraging habitat quality on the population decline of the Red-cockaded Woodpecker in Oklahoma. In prep.

- Locke, B.A., R.N. Conner, & J.C. Kroll. 1983. Factors influencing colony site selection by Red-cockaded Woodpeckers. Pp. 46-50 in Proc. of Red-cockaded Woodpecker Symp. II. (D. A. Wood, Ed.). Gainesville, Florida, Game Fresh Water Fish Comm., U.S. Fish Wildl. Serv. and U.S. For. Serv.
- Masters, R.E., J.E. Skeen, & J.A. Garner. 1989. Red-cockaded Woodpecker in Oklahoma: An update of Wood's 1974-1977 Study. Proc. Okla. Acad. Sci. 69:27-31.
- Porter, M.L., & R.F. Labinsky. 1986. Home range and foraging habitat of Red-cockaded Woodpeckers in northern Florida. J. Wildl. Manage. 50:239-247.
- Renken, R.B. and E.P. Wiggers. 1989. Forest characteristics related to Pileated Woodpecker territory size in Missouri. Condor 91:642-652.
- Rice, W.R. 1989. Analyzing tables of statistical tests. Evolution 43:223-225.
- Sokal, R.R., & F.J. Rohlf. 1981. Biometry. San Francisco, California, W.H. Freeman and company.
- Teitelbaum, R.D., & W.P. Smith. 1985. Cavity-site characteristics of the Red-cockaded Woodpecker in Fontainbleau State Park, Louisiana. Proc. Louisiana Acad. Sci. 48:116-122.
- U.S. Fish and Wildlife Service. 1985. Red-cockaded Woodpecker Recovery Plan. U.S. Fish and Wildl. Serv., Atlanta, Georgia. 88 Pp.

- Van Balen, J.B., & P.D. Doerr. 1978. The relationship of understory vegetation to Red-cockaded Woodpecker activity. Proc. annu. conf. SE Assoc. Fish & Wildl. Agencies 32:82-92.
- Wilkinson, L. 1989. SYSTAT: The system for statistics. Evanston, Illinois, SYSTAT Inc.
- Wood, D.A. 1983. Foraging and colony habitat characteristics of the Red-cockaded Woodpecker in Oklahoma. Pp. 51-58 in Proc. of Red-cockaded Woodpecker Symp. II. (D.A. Wood, Ed.). Gainesville, Florida, Game Fresh Water Fish Comm., U.S. Fish Wildl. Serv. and U.S. For. serv.

Table 1. Comparisons of hardwood characteristics between cluster and non-use site base tree plots by quadrant and overall using Wilcoxon's paired signed ranks test. Variables compared were: (1) number of hardwood saplings; (2) DBH of hardwoods (cm); (3) total basal area of hardwoods (m²/ha); and (4) height of hardwoods (m). All sample sizes = 18.

Hardwood	<u>Cluster</u>	<u>Non-use</u>		<u>Cluster</u>	<u>Non-use</u>	
Variable	Median(range)	Median(range)	P	Median(range)	Median(range)	P
	Quadrant 1			Quadrant 2		
Saplings	3.0 (0.0-17.0)	2.0 (0.0-26.0)	0.777	4.0 (0.0-15.0)	3.0 (0.0-26.0)	0.593
DBH	8.9 (0.0-13.7)	11.2 (7.7-16.3)	0.004 ^a	10.3 (7.8-15.5)	12.5 (7.2-30.2)	0.085
Basal area	5.9 (0.0-14.8)	11.2 (5.4-19.1)	0.001 ^a	8.8 (2.0-16.0)	12.6 (3.6-28.5)	0.058
Height	6.9 (0.0- 9.9)	8.6 (3.4-14.6)	0.003 ^a	8.5 (6.9-12.0)	9.9 (7.1-14.5)	0.028
	Quadrant 3			Quadrant 4		
Saplings	3.5 (0.0-10.0)	1.0 (0.0-17.0)	0.678	3.0 (0.0- 8.0)	3.0 (0.0-12.0)	0.757
DBH	9.8 (7.1-21.9)	12.7 (7.4-19.4)	0.170	9.6 (8.3-18.5)	10.6 (8.1-24.4)	0.446
Basal area	7.0 (3.8-22.9)	12.9 (1.8-32.5)	0.199	7.2 (1.8-28.2)	9.7 (4.2-31.1)	0.184
Height	8.6 (7.3-12.7)	11.5 (6.2-17.8)	0.011 ^a	7.8 (6.2-15.5)	9.3 (6.6-18.2)	0.035
	Overall					
Saplings	13.5 (1.0-44.0)	13.5 (0.0-79.0)	0.913			
DBH	10.3 (7.9-12.6)	12.0 (9.1-16.7)	0.018			
Basal area	8.6 (3.4-14.6)	11.3 (7.9-14.5)	0.008 ^a			
Height	7.8 (6.7-10.7)	9.5 (7.5-12.8)	0.002 ^a			

^a P value is significant when using the sequential Bonferroni technique to control Type 1 error for each variable; initial critical value = 0.01 (0.05/5 tests).

Table 2. Comparisons of pine characteristics between cluster and non-use site base tree plots by quadrant and overall using Wilcoxon's paired signed ranks test. Variables compared were: (1) number of pine saplings; (2) DBH of pines (cm); (3) total basal area of pines (m^2/ha); and (4) height of pines (m). All sample sizes = 18.

Pine Variable	Cluster		P	Non-use		P
	Median(range)	Median(range)		Median(range)	Median(range)	
	Quadrant 1			Quadrant 2		
Saplings	0.0 (0.0-59.0)	0.0 (0.0- 2.0)	0.012 ^a	0.0 (0.0-42.0)	0.0 (0.0- 5.0)	0.161
DBH	24.5 (0.0-53.2)	22.3 (0.0-58.4)	0.984	22.2 (0.0-49.3)	18.1 (0.0-37.2)	0.744
Basal area	18.4 (0.0-45.9)	11.9 (0.0-26.8)	0.030	17.1 (0.0-67.4)	8.1 (0.0-37.3)	0.647
Height	14.3 (0.0-33.5)	14.9 (0.0-21.5)	0.679	14.2 (0.0-31.6)	14.7 (7.1-14.5)	0.744
	Quadrant 3			Quadrant 4		
Saplings	0.0 (0.0- 27.0)	0.0 (0.0- 1.0)	0.034	0.0 (0.0-91.0)	0.0 (0.0- 1.0)	0.161
DBH	19.8 (0.0- 42.9)	23.2 (0.0-39.6)	0.356	21.7 (0.0-46.0)	19.5 (0.0-43.6)	0.948
Basal area	13.5 (0.0- 56.1)	7.8 (0.0-26.8)	0.177	13.4 (0.0-36.8)	13.1 (0.0-57.8)	0.679
Height	14.1 (0.0- 31.0)	13.2 (0.0-23.3)	0.463	12.9 (0.0-30.8)	15.7 (0.0-22.4)	0.327
	Overall					
Saplings	3.5 (0.0-219.0)	0.0 (0.0- 5.0)	0.006 ^a			
DBH	21.6 (7.9- 42.6)	19.6 (12.0-27.7)	0.372			
Basal area	17.8 (8.0- 31.8)	12.0 (1.3-19.8)	0.028			
Height	12.9 (7.3- 25.1)	13.5 (3.1-22.2)	0.879			

^a P value is significant when using the sequential Bonferroni technique to control Type 1 error for each variable; initial critical value = 0.01 (0.05/5 tests).

Table 3. Comparisons of basal area (m^2/ha) of pines and hardwoods sampled at 31.3 and 71.3 m from the base tree between cluster and non-use sites by quadrant and overall using Wilcoxon's paired signed ranks test. All sample sizes = 18.

Quadrant	Hardwood basal area			Pine basal area		
	Cluster Median(range)	Non-use Median(range)	P	Cluster Median(range)	Non-use Median(range)	P
31.3 m from base tree						
1	10.3 (0.0-25.3)	11.5 (0.0-23.0)	0.297	16.1 (0.0-27.5)	16.1 (0.0-34.4)	0.553
2	11.5 (2.3-23.0)	11.5 (4.6-23.0)	0.568	12.6 (0.0-32.1)	14.9 (0.0-39.0)	0.717
3	9.2 (2.3-20.7)	12.6 (4.6-20.7)	0.008 ^a	20.7 (6.9-32.1)	13.8 (0.0-27.5)	0.004 ^a
4	10.3 (2.3-23.0)	11.5 (4.6-29.8)	0.121	17.2 (6.9-32.1)	11.5 (2.3-29.8)	0.161
Overall	11.2 (4.0-17.2)	11.8 (8.6-21.8)	0.013	16.4 (9.2-25.2)	13.8 (4.6-25.8)	0.472
71.3 m from base tree						
1	11.5 (0.0-23.0)	14.9 (0.0-28.8)	0.312	16.1 (0.0-27.5)	18.4 (0.0-34.4)	0.337
2	13.8 (4.6-29.8)	13.8 (6.9-23.0)	0.421	17.2 (0.0-36.7)	6.9 (0.0-23.0)	0.019
3	11.5 (4.6-20.7)	13.8 (4.6-27.5)	0.263	16.1 (2.3-48.2)	12.6 (2.3-32.1)	0.162
4	11.5 (4.6-16.1)	14.9 (4.6-27.5)	0.041	16.1 (2.3-48.2)	6.9 (0.0-43.6)	0.041
Overall	12.1 (8.0-22.2)	14.1 (9.2-21.8)	0.016	15.2 (5.2-32.1)	12.3 (4.0-19.5)	0.020

^a P value is significant when using the sequential Bonferroni technique to control Type 1 error for each variable at each distance (31.3 and 71.3 m); initial critical value = 0.01 (0.05/5 tests).

Table 4. Comparison of pine and hardwood basal area (m^2/ha) between 127 randomly placed sample points and 72 cluster site sample points (4 at each base tree). Cluster site samples were taken 31 m from the base tree on the quadrant boundary lines.

Tree Type/			
Site	$\bar{X} \pm SD$	t-statistic	P
Pine			
Cluster	15.3 \pm 7.3	4.07	>0.001
Random	10.8 \pm 7.6		
Hardwood			
Cluster	7.8 \pm 4.2	0.53	0.595
Random	7.4 \pm 4.8		

★ 71.3 m Basal Area Sample

★ 31.3 m Basal Area Sample




Figure 1. Sampling design used at cluster and non-use site base-tree plots.

VITA

Jeffrey F. Kelly

Candidate for the Degree of
Master of Science

Thesis: THE INFLUENCE OF HABITAT QUALITY ON THE POPULATION
DECLINE OF THE RED-COCKADED WOODPECKER IN MCCURTAIN
COUNTY WILDERNESS AREA, OKLAHOMA

Major Field: Zoology

Biographical:

Personal Data: Born in San Francisco, California, 13
July, 1965, the son of Richard F. and Barbara
Kelly

Education: Graduated from Wayne Valley High School,
Wayne, New Jersey, in June 1983; received Bachelor
of Science Degree in Wildlife Management from
University of Maine in May, 1987; completed
requirements for the Master of Science degree at
Oklahoma State University in May, 1991.

Professional Experience: Graduate Research Assistant,
Oklahoma Cooperative Fish and Wildlife Research
Unit, Oklahoma State University, April 1989 to May
1990; Biological Technician, U.S. Fish and
Wildlife Service, Hawaii Research Group, Volcano
Hawaii, June 1987 to April 1989; Volunteer, U.S.
Fish and Wildlife Service, Anchorage, Alaska, May
to August 1986.