



US00PP16801P2

(12) **United States Plant Patent**  
**Taliaferro et al.**

(10) **Patent No.:** **US PP16,801 P2**

(45) **Date of Patent:** **Jul. 11, 2006**

(54) **PATRIOT TURF BERMUDAGRASS**

(50) Latin Name: *Cynodon dactylon* var. *dactylon*×*C. transvaalensis*  
Varietal Denomination: **Patriot**

(75) Inventors: **Charles M. Taliaferro**, Stillwater, OK (US); **Dennis L. Martin**, Stillwater, OK (US); **Jeffrey A. Anderson**, Stillwater, OK (US); **Michael P. Anderson**, Stillwater, OK (US)

(73) Assignee: **The Board of Regents for Oklahoma State University**, Stillwater, OK (US)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 1 day.

(21) Appl. No.: **10/873,809**

(22) Filed: **Jun. 22, 2004**

(51) **Int. Cl.**  
**A01H 5/00** (2006.01)

(52) **U.S. Cl.** ..... **Plt./389**

(58) **Field of Classification Search** ..... Plt./389  
See application file for complete search history.

(56) **References Cited**  
**PUBLICATIONS**

United States Department of Agriculture, Agricultural Research Service, National Bermudagrass Test Final Report 1997–2001, *NTEP* No. 02–7.

Anderson et al., Longer Exposure Durations Increase Freeze Damage to Turf Bermudagrasses, 2003, *Crop Sci.*, vol. 43, pp 973–977.

Anonymous, Shanghai Bermudagrass—The True Blue-Green Winter Hardy Grass for the South, Access Jul. 8, 2003, *SuperSod*, a division of Patten Seed Company, published on world wide web at <http://www.pattenseed.com/sod-bgsh.html>.

Hanna et al., Registration of ‘Tifton 10’ Bermudagrass, 1990, *Crop Sci.*, vol. 30(6), pp 1355–1356.

Landry, Bermudagrass Lawn Management, 1996, *Educational Support Services of the Cooperative Extension Service of The University of Georgia*, Circular 837.

*Primary Examiner*—Anne Marie Grunberg

*Assistant Examiner*—Annette H Para

(74) *Attorney, Agent, or Firm*—Fellers, Snider, Blankenship, Bailey & Tippens, P.C.

(57) **ABSTRACT**

‘Patriot’ is an improved clonal turf bermudagrass cultivar suitable for many applications including golf course fairways and tees athletic fields, home and commercial landscape sites, and parks and playgrounds. It is a tetraploid (2n=4x=36 chromosomes) interspecific F<sub>1</sub> hybrid from a cross of *Cynodon dactylon* by *C. transvaalensis* characterized by dark blue-green color, vigorous growth, good winter hardiness, and good turf quality.

**8 Drawing Sheets**

**1**

**BACKGROUND OF THE INVENTION**

1. Field of the Invention

‘Patriot’ bermudagrass is a clonally propagated F<sub>1</sub> hybrid plant from a cross of *Cynodon dactylon* var. *dactylon* by *C. transvaalensis*. The cultivar will be marketed as Certified class sod and/or Certified class sprigs.

Background of the Invention:

The maternal parent of ‘Patriot’ was ‘Tifton 10’ bermudagrass, *Cynodon dactylon* var. *dactylon*. The pollen parent of ‘Patriot’ was a selected plant of *Cynodon transvaalensis* designated by its field nursery location identity as 4200 TN 26-8. ‘Tifton 10’ is a commercial cultivar that differs from most *C. dactylon* plants in having 54 chromosomes (hexaploid; 2n=6x=54) compared to the usual 36 chromosomes (tetraploid; 2n=4x=36) and having a darker green foliage color (Hanna, et al., 1990). The *C. transvaalensis* pollen parent has 18 chromosomes (diploid; 2n=2x=18). ‘Patriot’ is a tetraploid with 2n=4x=36 chromosomes, presumably having inherited 27 chromosomes (3 genomes) from ‘Tifton 10’ and 9 chromosomes (1 genome) from the *C. transvaalensis* parent.

The cross between ‘Tifton 10’ and *C. transvaalensis* “4200 TN 26-8” was made in 1992. Seeds from this cross

**2**

and crosses between other parent plants were germinated in late winter 1993 and the seedlings individually planted in greenhouse pots. The potted plants were transplanted to a field screening nursery in spring 1993 as spaced (3 meter centers) plants. ‘Patriot’ was first vegetatively propagated in late 1994. The vegetative propagation occurred in a greenhouse on the Agronomy Farm, Oklahoma State University and was selected in late 1994 for advancement to more comprehensive testing including testing in replicated experiments through time and space to measure adaptation and performance characteristics. At the time of its first selection in 1994 it was identified as “OKC 18-4” and evaluated under this designation. ‘Patriot’ has been clonally propagated through many generations since 1994 without any indication of genetic change. Advanced generation clonal plants appear identical to the original plant in morphological phenotype and in genetic characteristics that have been measured.

**SUMMARY OF THE INVENTION**

‘Patriot’ bermudagrass produces a high quality turf suitable for most turf applications including golf course fairways and tees, home lawns, commercial lawns, playgrounds, parks, and athletic fields. ‘Patriot’ produces a dense, dark-green turf of high visual quality. It has medium-fine texture,

grows vigorously, and has good cold hardiness. These characteristics make it especially useful in climatic regions representing zones of transition between cool- and warm-season turfgrass species. 'Patriot' is an F<sub>1</sub> hybrid of the cross 'Tifton 10' by *C. transvaalensis* "4200 TN 26-8". It has 2n=4x=36 chromosomes having inherited 27 chromosomes (3 genomes) from 'Tifton 10' and 9 chromosomes (1 genome) from *C. transvaalensis*. 'Patriot' is highly sterile and is propagated asexually by sprigs and sod. 'Patriot' can be distinguished from other turf bermudagrass cultivars by morphological and/or genetic characteristics outlined herein.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a photograph of five-week old growth of 'Tifton 10' (left), 'Patriot' (center), and *C. transvaalensis* "4200 TN 26-8" (right). The intermediate size and texture of 'Patriot' relative to the parents are illustrated.

FIG. 2 illustrates stolons of 'Tifton 10' (bottom), 'Patriot' (second from bottom), 'U-3' (third from bottom), and 'Quicksand' (top). Stolons were taken from five-week old growth of potted plants in the greenhouse. The relative length and diameter of internodes is illustrated as are the relative length and width of leaves.

FIG. 3 illustrates laterally growing stolons of 'Patriot' during stand establishment. The photograph was taken in early October showing the response of 'Patriot' to cool temperatures in production of anthocyanin pigmentation.

FIG. 4 illustrates leaf tips of 'Patriot' (left), 'Tifton 10' (second from left), *C. transvaalensis* "4200 TN 26-8" (third from left), and 'U-3' (right). Leaf tips of 'Patriot' and 'Tifton 10' leaf tips tend to be more oval and less pointed than typically found in bermudagrass.

FIG. 5 illustrates a typical inflorescence of 'Patriot' with four racemes typically varying in length from 2.5 to 3.5 cm.

FIG. 6 are photographs of 'Patriot', 'Tifway', and 'Tifgreen' mowed twice weekly at 19 mm (3.4 inch) height. A) 'Patriot' and Tifway, B) 'Patriot' and Tifgreen, and C) close-up of 'Patriot'.

FIG. 7 are photographs of field plots of 'Patriot' (A & C) and 'Tifsport' (B & D) at Stillwater, Okla. on Apr. 5, 2002 and Apr. 12, 2002, demonstrating the typically earlier growth (green up) of 'Patriot' relative to Tifsport.

FIG. 8 illustrates the DNA profile of 'Patriot' and 12 other clonal turf bermudagrass cultivars using DAF (DNA amplification fingerprinting) primer 9111 (sequence GAAACGCC). The numbers on the Y axis indicate numbers of base pairs of nucleotides. The arrow points to a prominent band at the 220 bp region that is unique to 'Patriot' among these cultivars.

#### DETAILED BOTANICAL DESCRIPTION

'Patriot' experimental name "OKC 18-4") is a distinct cultivar of *Cynodon* from the interspecific cross of *C. dactylon* Pers. cv. 'Tifton 10' and *C. transvaalensis* Burt-Davy cv. '4200 TN 26-8'. It was developed and vegetatively propagated by the Oklahoma Agricultural Experiment Station, Stillwater, Okla. Identifying morphological characteristics of 'Patriot' are its dark green color, strong anthocyanin pigmentation of stolon internodes during periods of cool temperatures, and high shoot density. It is intermediate to its parents in plant size and texture as illustrated in FIG. 1. 'Patriot' has finer texture than 'Tifton 10', 'U-3', and 'Quickstand' as illustrated in FIG. 2 by the relative diameter

and length of stolons and stolon internodes. 'Patriot' is a stoloniferous sward-forming perennial with slender underground rhizomes; surface stolons are slender attaining a maximum diameter of approximately 1 mm, and a maximum internode length of approximately 6 cm. During establishment under good growing conditions stolons may attain a length of approximately 70 cm (FIG. 3). Shoot numbers emanating from nodes of stoloniferous runners range from 2 to 5 and average 3.0. Culms are slender varying in height on unmowed swards from about 10 to 25 cm. Culm density is high producing a dense sward (sod). Culm leaves are flat, glabrous on the abaxial and adaxial surfaces, and slightly serrate on the edges. Culm leaves emanating from the last visible node below the apex of shoots vary in width at their base from 2 to 4 mm and range in length from about 25 to 38 mm. The leaf tips resemble the Tifton 10 parent and are more rounded that is typical for *Cynodon* plants (FIG. 4) The ligule is a membranous rim about 0.2 mm long with very short hairs on the edge except at the outer edges where a tuft of much longer hairs grow. Leaf sheaths are glabrous. Inflorescence is a single whorl of 2 to 5, usually 3 or 4, racemes averaging 33 mm in length (FIG. 5). Racemes contain on average 25 spikelets spaced 2 mm apart and 2 mm in length. The lower and upper glumes are approximately ½ and ¾ the length of the spikelet.

'Patriot' was evaluated under the experimental name 'OKC 18-4' in the National Turf Evaluation Program (NTEP) bermudagrass test conducted at 21 locations from 1997 through 2001. This multi-environment testing provided the most definitive data on the performance characteristics of 'Patriot' relative to other major commercial clonal turf bermudagrass cultivars. The trademarked cultivar 'Shanghai' in this test is indicated by descriptions published by the Patton Seed Co. (2003) and Landry (2003) to be the same as 'Tifton-10'.

'Patriot' produces a high quality turf surface as indicated by visual ratings of turf quality (Table 1, FIG. 6). Under management schedule "A" ½ to ¾ in. mowing height), the mean turf quality rating of 'Patriot' (6.1) was less than that of Tifway (6.4) and Tifsport (6.5), equal to that of CN 2-9, and greater than all other cultivars, including 'Shanghai'. Under management schedule "B" ¾ to 1 inch mowing height), the mean turf quality rating of 'Patriot' (6.6) was significantly higher than those of all cultivars except 'Midlawn'.

'Patriot' is relatively stable in turfgrass quality performance across environments (Table 2). It was lowest of all clonal varieties in sum-of-ranks across locations and finished in the top 25% of all entries in 65% of the locations.

'Patriot' has not shown high susceptibility to the major disease or insect pests of turf bermudagrass in the northern half of geographic region of bermudagrass use in the U.S. where it is expected to primarily be used. Spring dead spot disease caused by *Ophiosphaerella herpotricha* is the most serious disease of turf bermudagrass in the upper south of the U.S. 'Patriot' was evaluated for response to spring dead spot disease in a controlled test at Stillwater, Okla. during 2000 through 2002. Replicated plots were inoculated in September 1997 with *Ophiosphaerella herpotricha*. Measurements of the size of necrotic patches indicated 'Patriot' to have good tolerance to the disease (Table 10), much higher than very susceptible cultivars such as Tifway. 'Patriot' has exhibited very little leaf disease indicating it to be highly resistant to such disease caused by *Bipolaris cynodontis* and other dematiaceous fungal pathogens. 'Patriot' has good resistance to the bermudagrass stunt mite,

*Aceria neocynodonis*, as indicated by minimal infestation when grown in field nurseries with other bermudagrasses that exhibited severe infestation. ‘Patriot’ has not been evaluated for response to fall armyworm, *Spodoptera frugiperda*, black cutworm, *Agriotis ipsilon*, or sod webworm, *Herpetogramma phaeopteralis*, all pests of bermudagrass. However, no significant injury was observed due to these pests during the time ‘Patriot’ was under evaluation.

Mean genetic color ratings of ‘Patriot’ were the same as those of Shanghai and CN 2-9 and greater than all other clonal cultivars (Table 3). ‘Patriot’ has nearly the same dark blue-green color of Tifton-10 (FIG. 1).

‘Patriot’ bermudagrass fertilized monthly during the growing season with nitrogen at rates of 1.0 to 1.5 Lbs. N/1000 ft<sup>2</sup> has the following color scale ratings using Munsell® Color Charts (Munsell Color, GretabMacbeth, LLC, 617 Little Britain Road, New Windsor, N.Y. 12553) for Plant Tissues.

1. ‘Patriot’ produces leaves of dark green color [7.5 GY (4/4 to 4/6)].
2. ‘Patriot’ produces leaf sheaths that vary in color from green [7.5 GY. (6/6to 6/8)] to dark purple [10RP (2/4 to 3/4)]. The purple anthocyanin pigmentation varies with age of the sheaths and with environmental conditions, being more pronounced in older tissues and when the plant is subjected to cool temperatures. However, anthocyanin pigmentation is usually present in leaf sheaths of ‘Patriot’ throughout the growing season, with the presence and intensity the pigmentation tending to be greater in leaf sheaths emanating from stolons compared to leaf sheaths of upright shoots. Pigmentation usually first develops at the base of the sheath and may ultimately occur throughout the sheath.
3. ‘Patriot’ produces internodes that vary in color from light green [10Y (5/4 to 6/4)] to dark purple [(10RP (2/4 to 3/4)]. As in the case of leaf sheaths, anthocyanin pigmentation in internodes varies with age of tissue and environmental conditions in internodes as in leaves. Pigmentation tends to be more pronounced in older tissues and when plants are subjected to cool conditions. The presence and intensity of pigmentation tends to be greater in internodes emanating from stolons compared to upright shoots.
4. ‘Patriot’ bears inflorescences of light green color [2GY (6/6 to 6/8)]. At maturity the inflorescences are straw colored [2.5YG (6/4/ to 7/4)].

Mean leaf texture of ‘Patriot’ was rated equal to Midlawn and CN 2-9, finer than Shanghai, and coarser than other clonal cultivars (Table 4). ‘Patriot’ has distinctly finer leaf texture than Tifton-10 (FIG. 1).

Mean stand density (Table 5) and mean percent living ground cover (Table 6) of ‘Patriot’ as indicated by ratings are greatest during summer months and on a par with other clonal cultivars throughout the growing season.

‘Patriot’ has good winter hardiness relative to other turf bermudagrass cultivars as indicated by winter injury ratings at Wichita, Kans. (Table 7) and laboratory measure of response to low freezing temperatures (Table 8). Its good low temperature tolerance reduces risk of winter injury relative to less winter hardy cultivars when grown in colder climatic regions where bermudagrass is used. The earlier growth of ‘Patriot’ relative to ‘TifSport’ at Stillwater, Okla. (36.06° N. Latitude; 9797.03 w. Longitude) is illustrated in FIG. 7.

‘Patriot’ has vigorous growth relative to other clonal cultivars as indicated by its spread during establishment at Fayetteville, Ark., Griffin, Ga., Lexington, Ky. and Starkville, Miss. (Table 9). The rate of spread during establishment of ‘Patriot’ and its growth potential are similar to Tifton 10 (Shanghai) although it has smaller plant size (texture). Ratings of seedhead abundance relative to other clonal cultivars may be found in Table 9a.

‘Patriot’ can be distinguished from other cultivars by DNA profiling. FIG. 8 shows a DAF (DNA Amplification Fingerprinting) profile using DAF primer 9111. The arrow points to a band at the 220 base pair region that is unique to ‘Patriot’ among the included cultivars. Other band differences also exist between ‘Patriot’ and the other cultivars in this profile.

The nucleotide sequence of DAF primer 9111 is ‘GAAACGCC’ (SEQ ID NO: 1). DAF primer 9111 was obtained from Integrated DNA Technologies, Inc., 1710 Commercial Park, Coralville, Iowa 52241.

TABLE 1

Cultivar	Maintenance Schedule <sup>3</sup>	
	“A”	“B”
Cardinal	5.4	5.7
CN2-9	6.1	5.8
Midlawn	5.8	6.5
Mini-Verde	5.6	5.1
Patriot	6.1	6.6
Shanghai	5.6	6.1
Tifsport	6.5	6.1
Tifway	6.4	6.2
Tifgreen	6.1	6.3
5% LSD	0.2	0.2
CV (%)	15.8	14.2

<sup>1</sup>Based on a visual rating scale of 1 to 9 with 1 being poorest and 9 representing ideal turf.

<sup>2</sup>Means over years 1997 through 2001.

<sup>3</sup>Schedule “A”: mowing height = ½ to ¾ inch, ¾ to 1 Lb. N/100 ft<sup>2</sup>/growing month, irrigation to prevent visual drought stress, mowed 3 to 5 times/month. Schedule “B”: mowing height = ¾ to 1 inch, ½ to ¾ Lb. N/1000 ft<sup>2</sup>/growing month, irrigation to prevent dormancy, mowed 1 to 2 times/week. Schedule “A” conducted at nine locations; Schedule “B” conducted at 11 locations.

TABLE 2

Summary of turfgrass quality ratings for ‘Patriot’ and eight other cultivars in the 1997 National Turfgrass Evaluation Program bermudagrass test.

Cultivar	Statistics for all locations			
	All locations		Sum of	
	Mean <sup>1</sup>	Rank <sup>2</sup>	Ranks <sup>3</sup>	Rank <sup>4</sup>
<b>Clonal Cultivars</b>				
Cardinal	5.5	13	327	15
CN 2-9	5.9	9	213	10
Midlawn	6.1	7	156	7
Mini-Verde	5.4	14	312	13
Patriot	6.3	4	112	2
Shanghai	5.9	10	190	9
Tifgreen	6.2	6	136	6
Tifsport	6.4	2	125	4
Tifway	6.3	3	126	5

TABLE 2-continued

Summary of turfgrass quality ratings for 'Patriot' and eight other cultivars in the 1997 National Turfgrass Evaluation Program bermudagrass test.

Seeded Cultivars				
Arizona Common	4.6	28	539	28
Blackjack	5.2	17	343	18
Blue-Muda	5.0	24	433	25
J-540	5.2	18	353	19
Jackpot	4.9	26	470	27
Majestic	5.2	19	340	17
Mirage	5.0	25	425	24
Numex-Sahara	4.9	27	469	26
Princess	6.3	5	121	3
Pyramid	5.0	23	416	23
Riviera	6.5	1	91	1
Savannah	5.3	15	317	14
ShangriLa	5.1	21	409	22
Southern Star	5.3	16	335	16
Sundevil II	5.1	22	394	21
SWI-11	5.8	11	234	12
Sydney	5.2	20	385	20
Transcontinental	5.7	12	215	11

Statistics for all locations			
	Highest Rank <sup>5</sup>	Lowest Rank <sup>6</sup>	Maximum in Top 25% <sup>7</sup>
Clonal Cultivars			
Cardinal	1	28	20
CN 2-9	2	25	25
Midlawn	1	27	60
Mini-Verde	5	28	30
Patriot	1	11	65
Shanghai	3	16	30
Tifgreen	1	22	70
Tifsport	1	20	60
Tifway	1	23	75
Seeded Cultivars			
Arizona Common	23	28	0
Blackjack	4	27	5
Blue-Muda	5	26	5
J-540	12	24	0
Jackpot	16	28	0
Majestic	13	26	0
Mirage	7	28	5
Numex-Sahara	16	27	0
Princess	1	12	60
Pyramid	9	27	0
Riviera	1	12	80
Savannah	12	23	0
ShangriLa	10	27	0
Southern Star	11	28	0
Sundevil II	13	28	0
SWI-11	2	20	25
Sydney	6	28	5
Transcontinental	3	21	20

<sup>1</sup>Mean—An average of all the turfgrass quality ratings from all locations.  
<sup>2</sup>Rank—Ranking of the mean of all quality ratings.  
<sup>3</sup>Sum of ranks—a sum of all the rankings from the various locations.  
<sup>4</sup>Rank—The ranking of the sum of ranks.  
<sup>5</sup>Highest rank—The highest ranking achieved by that entry at any one location.  
<sup>6</sup>Lowest rank—The lowest ranking achieved by that entry at any one location.  
<sup>7</sup>Maximum in top 25%—The percentage of locations where that entry finished in the top 25% of all entries.

TABLE 3

Mean genetic color ratings of 'Patriot' and eight other bermudagrass cultivars in the 1997 National Turf Evaluation Program bermudagrass test.<sup>1</sup>

Cultivar	Rating <sup>2</sup>
Cardinal	4.7
CN 2-9	7.2
Midlawn	6.5
Mini-Verde	6.7
Patriot	7.4
Shanghi	7.5
Tifgreen	6.2
Tifsport	7.0
Tifway	7.1
5% LSD	0.2
CV (%)	14.4

<sup>1</sup>Means over 21 locations and 5 years (1997–2001).  
<sup>2</sup>Based on a visual rating scale with 1 being light green and 9 being dark green.

TABLE 4

Mean leaf texture ratings of 'Patriot' and eight other bermudagrass cultivars in the 1997 National Turf Evaluation Program bermudagrass test.<sup>1</sup>

Cultivar	Rating <sup>2</sup>
Cardinal	8.2
CN 2-9	6.9
Midlawn	6.9
Mini-Verde	8.2
Patriot	6.7
Shanghai	5.1
Tifgreen	7.7
Tifsport	7.2
Tifway	7.3
5% LSD	0.2
CV (%)	14.4

<sup>1</sup>Means over 20 locations and 5 years (1997–2001).  
<sup>2</sup>An estimate of leaf width based on a visual rating scale with 1 equaling coarse and 9 equaling fine.

TABLE 5

Mean stand density ratings of 'Patriot' and eight other bermudagrass cultivars in the 1997 National Turf Evaluation Program bermudagrass test.<sup>1</sup>

Cultivar	Stand density rating <sup>2</sup>		
	Spring	Summer	Fall
Cardinal	7.6	7.3	7.4
CN 2-9	6.5	7.1	7.0
Midlawn	7.1	7.1	7.1
Mini-Verde	6.9	8.1	7.8
Patriot	6.7	7.5	7.2
Shanghai	6.2	6.8	6.4
Tifgreen	7.3	7.6	7.5
Tifsport	7.2	7.6	7.6
Tifway	7.3	7.7	7.6
5% LSD	0.4	0.3	0.3
CV (%)	18.1	12.3	13.8

<sup>1</sup>Means over 9 locations and 5 years (1997–2001).  
<sup>2</sup>An estimate of living plants or tillers per unit area based on a visual rating scale with 9 equaling maximum density.

TABLE 6

Mean percent living ground cover ratings of 'Patriot' and eight other bermudagrass cultivars in the 1997 National Turf Evaluation Program bermudagrass test.<sup>1</sup>

Cultivar	Ground cover rating <sup>1</sup>		
	Spring <sup>2</sup>	Summer <sup>3</sup>	Fall <sup>4</sup>
Cardinal	87.0	89.9	86.7
CN 2-9	70.1	88.3	91.6
Midlawn	85.9	90.8	88.2
Mini-Verde	59.3	78.8	87.7
Patriot	84.0	90.7	90.0
Shanghai	77.8	87.3	87.6
Tifgreen	78.7	91.4	91.0
Tifsport	71.4	89.5	93.9
Tifway	76.8	91.4	93.7
5% LSD	5.5	3.8	4.0
CV (%)	24.8	14.0	12.1

<sup>1</sup>Percent living ground cover based on surface area covered by the originally planted species.  
<sup>2</sup>Means over 14 locations and 5 years.  
<sup>3</sup>Means over 9 locations and 5 years.  
<sup>4</sup>Means over 8 locations and 5 years.

TABLE 7

Mean winter injury ratings of 'Patriot' and eight other bermudagrass cultivars in the 1997 National Turf Evaluation Program bermudagrass test at Wichita, KS in 2001.

Cultivar	Rating
Cardinal	5.0
CN 2-9	2.3
Midlawn	5.7
Mini-Verde	3.0
Patriot	6.7
Shanghai	5.3
Tifgreen	4.0
Tifsport	3.7
Tifway	2.7
5% LSD	2.3
CV (%)	32.7

<sup>1</sup>Injury due to low winter temperatures based on a rating scale with 1 indicating severe injury and 9 indicating no injury.

TABLE 8

Freeze tolerance of 'Patriot' and six other turf bermudagrass cultivars. T<sub>mid</sub> values represent the midpoints of survival-temperature response curves<sup>1</sup>.

Cultivar	T <sub>mid</sub> °C.
Princess	-6.9a <sup>†</sup>
Tifway	-7.9b
Tifsport	-7.9b
Riviera	-8.3bc
U-3	-8.9cd
Patriot	-9.7de
Midlawn	-10.3e

<sup>1</sup>Jeffrey A. Anderson et al. 2003. Longer exposure durations increase freeze damage to turf bermudagrass. Crop Sci. 43:973-977.  
<sup>†</sup>Means of four repetitions are separated by Duncan's New Multiple Range Test at P,0.05.

TABLE 9

Mean establishment measurements or ratings of Patriot and eight other bermudagrass cultivars in the 1997 National Turf Evaluation Program bermudagrass test.<sup>1</sup> 1997 Measurements by date or weeks after planting.

Cultivar	Fayetteville, AR	Griffin, GA			Lexington, KY
	September	2 Wks	4 Wks	6 Wks	3 Wks
Cardinal	88.3	28.3	68.3	91.0	51.3
CN 2-9	60.0	26.7	58.3	81.7	33.3
Midlawn	86.7	26.7	60.0	81.7	46.0
Mini-Verde	53.3	26.7	58.3	71.7	36.7
Patriot	93.0	33.3	70.0	97.7	56.0
Shanghai	95.0	38.3	71.7	95.0	50.3
Tifgreen	90.0	28.3	66.7	90.0	48.3
Tifsport	63.3	18.3	53.3	71.7	34.3
Tifway	71.7	26.7	58.3	76.7	29.3
5% LSD	15.7	11.0	10.1	10.0	11.5
CV (%)	12.4	19.4	8.7	7.1	16.6

Cultivar	Lexington, KY		Starkville, MS	
	5 Wks	13 Wks	4 Wks	6 Wks
Cardinal	94.0	99.0	58.3	93.0
CN 2-9	77.7	99.0	50.0	81.7
Midlawn	93.3	99.0	58.3	93.0
Mini-Verde	80.0	98.3	41.7	81.7
Patriot	95.3	99.0	71.7	96.0
Shanghai	96.3	99.0	76.7	94.7
Tifgreen	92.3	99.0	66.7	96.3
Tifsport	68.3	97.3	53.3	83.3
Tifway	75.0	98.7	45.0	78.3
5% LSD	9.4	1.2	12.0	10.8
CV (%)	6.8	0.7	12.9	6.8

TABLE 9a

Mean seedhead ratings of 'Patriot' and eight other bermudagrass cultivars in the 1997 National Turf Evaluation Program bermudagrass test.<sup>1</sup>

Cultivar	Riverside, CA, 1999		Stillwater, OK, 1999		
Cardinal	8.0	4.0	8.0	8.0	8.0
CN 2-9	8.7	8.0	8.7	7.7	9.0
Midlawn	8.7	5.3	9.0	8.0	9.0
Mini-Verde	9.0	8.7	9.0	9.0	9.0
Patriot	9.0	7.7	9.0	8.0	8.3
Shanghai	9.0	7.3	8.7	8.0	9.0
Tifgreen	9.0	6.0	8.3	7.3	9.0
Tifsport	8.7	7.3	8.7	8.0	9.0
Tifway	8.3	8.0	8.7	8.0	9.0
5% LSD	0.9	1.0	0.7	0.4	0.3
CV (%)	4.8	9.0	4.4	3.1	2.1

<sup>1</sup>Ratings of seedhead abundance on a scale of 1 to 9 with 9 indicating none.

TABLE 10

Mean spring dead spot necrotic patch area (cm<sup>2</sup>) of bermudagrass cultivars in the 1997 National Turf Evaluation Program Bermudagrass Trial<sup>1</sup> at Stillwater, OK. Plots were inoculated with a blend of *Ophiophaerella herpotricha* isolates KS107, KS112 and KS188 in September 1997.

Entry	2000	2001	2002
Arizona Common	622	1791	2221
Cardinal	25	28	76
CN-2-9	383	931	1904
Continental	806	2124	3178

TABLE 10-continued

Mean spring dead spot necrotic patch area (cm<sup>2</sup>) of bermudagrass cultivars in the 1997 National Turf Evaluation Program Bermudagrass Trial<sup>1</sup> at Stillwater, OK. Plots were inoculated with a blend of *Ophiosphaerella herpotricha* isolates KS107, KS112 and KS188 in September 1997.

Entry	2000	2001	2002
Blackjack	751	1372	2035
Blue Muda	806	1835	2377
GN-1	1367	1722	1410
Jackpot	682	1805	2293
J-540	876	2333	2180
Majestic	1081	2531	3096
Midlawn	20	45	99
Mini-Verde	1195	3677	4902
Mirage	747	1626	1923
OKC 19-9	68	133	429
Patriot (OKC 18-4)	377	754	766
Princess 77	1947	3727	3312
Pyramid	1093	2417	2886
Riviera (OKS 95-1)	795	1116	1595
Sahara	618	1734	2416
Savannah	717	1731	2186
Shanghai	779	1207	1277
Shangri La	742	1760	2476
Southern Star	809	1753	2263
Sundevil II	637	1555	2289
SW1-11	537	2184	2589
Sydney	834	1257	2642
Tifgreen	486	1118	2125
Tifsport	473	927	996
Tifway	469	1133	2754
Yukon (OKS 91-11)	167	345	678
LSD (0.05)	698	1096	678

<sup>1</sup>Yukon and GN-1 were included at the Stillwater, OK, NTEP Bermudagrass Test Site as local standards.

## SEQUENCE LISTING

```

<160> NUMBER OF SEQ ID NOS: 1

<210> SEQ ID NO 1
<211> LENGTH: 8
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer

<400> SEQUENCE: 1

gaaacgcc


```

8


What is claimed is:

1. A bermudagrass plant substantially as described and illustrated in the specification herein.

\* \* \* \* \*




*Fig. 1*




*Fig. 2*







*Fig. 3*




*Fig. 4*




*Fig. 5*



*Fig. 6*



*Fig. 7*



*Fig. 8*