



# Leaf Curl of Peaches and Nectarines

Jen Olson

Plant Disease Diagnostician

Leaf curl of peaches and nectarines is one of the most commonly encountered diseases in unsprayed orchards and home yards during cold, wet springs. The disease can cause distortion of blossoms, leaves, fruit and shoots of the plant. In some years, the disease can cause significant defoliation and yield loss.

## Symptoms

The symptoms first appear roughly two weeks after the leaves emerge. Reddish spots develop on the leaves and as the disease progresses, these areas will thicken and pucker (Figure 1). Individual leaves may be affected or entire shoot tips may exhibit curl and distortion (Figure 2). Puckered areas on leaves may turn yellow or reddish-brown and occasionally powdery to gray spores can be observed on the leaves. Diseased leaves will eventually wither and fall from trees. Defoliation can result in sunburn injury on exposed branches.

Although new leaves emerge from dormant buds, their growth requirements reduce yield and may weaken diseased trees. The second flush of leaves is seldom affected by the disease. Defoliation of trees in several successive seasons may result in their death.

## Disease Cycle

Leaf curl disease is caused by a fungus, *Taphrina deformans*. Spores of the leaf curl fungus survive the summer, fall and winter months on bud scales and twigs. During periods of moist weather in early spring, the overwintering ascospores form conidia and they are carried into the buds on a film of water. During bud swell and opening, the developing leaves




Figure 1. Typical symptoms of leaf curl disease on a peach leaf.

Oklahoma Cooperative Extension Fact Sheets are also available on our website at: <http://osufacts.okstate.edu>

become infected. Cool, wet weather slows down leaf development and provides more time for leaf infection. Infection occurs readily at 50 F to 70 F, with 68 F being optimum. As the leaves grow and mature, they become less susceptible to infection. Secondary infection rarely occurs and is unimportant. If the weather is dry during bud swell and bud break, leaf curl is limited, and may not occur. However, the fungus may survive to infect trees another season.

The leaf curl fungus not only infects leaves, but also tender growing shoots and, more rarely, blossoms and fruits. Infected twigs become slightly swollen, may be yellowish and remain stunted. Infected blossoms and fruit usually fall early in the season. However, some infected fruit may remain in the trees. Diseased fruit will show shiny, raised, warty, discolored areas (Figure 3).



Figure 2. Peach leaves infected by the leaf curl fungus. Note puckering, deformation and change in color. Color changes usually occur long after initial infection.



**Figure 3. Fruit infected by the leaf curl fungus.**

## **Control**

Leaf curl is one of the relatively few serious diseases which may be easily, economically and effectively controlled. If a recommended fungicide is applied at the correct time, 95 to 98 percent control is usually obtained. A fungicide spray is

most effective when applied just before swelling of the buds. Timing depends on temperature and rainfall pattern of the season in each area of the state, usually around March. In commercial practice, an earlier application is frequently made. Ordinarily only one application is required, but if bud opening is delayed due to cold weather or if frequent rainfall occurs, a second application may be required. A fall application after leaf drop may also be helpful.

It must be remembered that leaf curl cannot be controlled after the fungus has infected the leaves and that fungicide applications must be made before the buds begin to swell. Thorough and complete coverage of twigs, branches and trunks is required. For suggested fungicides refer to the current OSU Extension Agents' Handbook of Insect, Plant, Diseases and Weed Control.

When leaf curl is severe, maintenance of tree vigor is very important. Fruit should be thinned to compensate for the loss of leaves. Over-cropping a tree will weaken it and make it more susceptible to winter injury. Trees should be adequately irrigated, and additional nitrogen fertilizer may be helpful for trees that are defoliated.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, Title IX of the Education Amendments of 1972, Americans with Disabilities Act of 1990, and other federal laws and regulations, does not discriminate on the basis of race, color, national origin, gender, age, religion, disability, or status as a veteran in any of its policies, practices, or procedures. This includes but is not limited to admissions, employment, financial aid, and educational services.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, the Director of Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources and has been prepared and distributed at a cost of 42 cents per copy. 0413 Revised. GH.