


SOIL AND CROP FACTORS

for

FERTILIZER RECOMMENDATIONS

1957



DIVISION OF AGRICULTURE  
OKLAHOMA A. & M. COLLEGE  
Stillwater

SOIL AND CROP FACTORS  
FOR FERTILIZER RECOMMENDATIONS 1957

Department of Agronomy  
Oklahoma A. & M. College

Principal Factors Determining Fertilizer Requirements

There is no simple means of determining the type or amount of fertilizer that will give highest returns for a specific field and crop. The principal basic factors determining the kind, the amount of fertilizer, and the best methods of application are:

1. Characteristics and production capability of the soil.

Soil texture and physical condition, organic matter and permeability along with soil depth, slope and degree of erosion largely determine relative soil productive capability. Deep permeable soils in good physical condition have an advantage for increased profitable returns from fertilizers because of higher production capabilities.

2. Crop to be grown and yield possibilities.

Crops differ in fertilizer requirement. Most important cash crops give highly profitable returns from fertilizers used properly. A crop used primarily for forage has different fertilizer requirements than the same crop grown for grain yield. A crop under irrigation has higher fertility requirements than in dryland conditions. Irrigated crops should be fertilized for maximum yields.

3. Previous treatment of the particular field.

The previous cropping and fertilizer history of a field gives reliable information on soil fertility level and its productive ability. The nitrogen contribution of previously grown legume crops must be estimated from this information. Fertilizers containing phosphorus and potassium can be expected to give residual effects on succeeding crops in cases of crop failure.

4. The individual farmer or rancher, his farming system and his limitations in capital and equipment are very important factors determining what the fertilizer recommendation should be. If resources are limited, the farmer should use the fertilizer materials that give the greatest return from limited investment. How the fertilizer is applied is as important as the kind and amount of fertilizer used. Good soil management practices are necessary to maintain and increase soil productivity. Commercial fertilizers cannot do the job alone.

## Soil and Other Tests

Soil tests, plant tissue tests, and plant deficiency symptoms are aids which may be used in evaluating some of the above factors. They support basic information, but will not satisfactorily substitute for it; therefore, it is necessary that all of this basic information be obtained and used in connection with these tests for sound, practical fertilizer recommendations.

The importance of obtaining representative soil samples for the soil tests cannot be overemphasized, since the tests can be no better than the samples. The accuracy of the most precise analytical procedure depends upon the sample used. An interpretation of analytical results is entirely dependent on how well the samples submitted represent the area concerned.

## Method of Fertilizer Application

Proper fertilizer placement is of great importance, particularly for stand establishment. Fertilizers in contact with the seed seriously reduce the stand for most large-seeded plants. Banded fertilizer two to three inches below or to one side of the seed has proved best for establishment of most row crops, grasses and legumes. For drill planted crops, such as small grains, fertilizers should be drilled in the row at seeding.

## General Information

Plant nutrients most commonly deficient in cultivated soils are nitrogen, phosphorus, and potassium. Commercial fertilizers contain one or more of these three elements. Other minerals required in large amounts by plants are calcium, magnesium, and sulphur. Calcium and magnesium are supplied in lime. Sulphur is contained in most fertilizers supplying phosphorus. The minor or trace elements for plants include iron, manganese, copper, zinc, boron, and molybdenum. Various plants differ greatly in their ability to utilize plant nutrients in soil, and the ability of a single plant to use these nutrients differs greatly under many soil and climatic conditions.

## Soil Organic Matter

Organic matter is the first factor in maintaining soil productivity. The level of organic matter in the soil can be maintained only by the addition of nitrogen and by the systematic return of crop residues and organic materials. Legume green manure crops and applications of barnyard manure within the cropping system are practical means of restoring soil organic matter in depleted soils. Applications of nitrogen fertilizers to highly carbonaceous crop residues such as straw and corn stalks are effective means for maintaining organic matter levels in soils.

TABLE I. SOIL REACTION, SOIL TEXTURE, AND LIME REQUIREMENT

Soil Reaction	pH Range	Sand	Lime Requirement in Tons/Acre	
			Sandy Loam, Loam and Silt Loam	Clay Loam and Clay Soil
Slightly Acid	6.1 to 6.4	None	1	1 to 2
Moderately Acid	5.5 to 6.0	1	1 to 2	2 to 3
Strongly Acid	5.0 to 5.4	1 to 2	2 to 3	3 to 4
Very Strongly Acid	Below 5.0	2	3 to 4	4 to 5

Legumes are usually benefited more by lime applications than are most grains and grasses. Lime is usually not recommended above pH 6.4. Soil organic matter, subsoil characteristics, and previous liming history will influence the lime requirement.

TABLE II. -- GENERAL FERTILIZER RECOMMENDATIONS FOR OKLAHOMA, 1957.

		SOIL TEST RESULTS:				
		Phosphorus		Potassium		
		Low to Very Low	Medium to High	Low to Very Low	Medium to High	
		FERTILIZER RECOMMENDED - POUNDS PER ACRE				
Crop	Nitrogen N	Phosphorus (P <sub>2</sub> O <sub>5</sub> )	Phosphorus (P <sub>2</sub> O <sub>5</sub> )	Potash (K <sub>2</sub> O)	Potash (K <sub>2</sub> O)	Remarks
<u>Corn:</u>						
Bottomland						
At Planting	10-20	20-60	20-40	20-60	0-20	Example: 100 lbs. 10-20-10 fertilizer at planting supplies 10 lbs. nitrogen, 20 lbs. P <sub>2</sub> O <sub>5</sub> and 10 lbs. K <sub>2</sub> O. 150 lbs ammonium nitrate sidedressed at second cultivation supplies 50 lbs. nitrogen.
Sidedressed	30-60					
Upland						
At Planting	10-20	20-40	10-20	20-40	None	
Sidedressed	20-40					
<u>Cotton:</u>						
Central and Eastern Okla.						
At Planting	10-20	20-40	10-20	20-40	0-20	Sandy soils in Western Oklahoma usually give better response to fertilizers than hard lands because of their lower fertility level and better moisture relationships.
At Planting	10-15	20-30	0-20	15-30	0-10	
Western Okla.						
At Planting	10-15	20-30	0-20	15-30	0-10	
<u>Sorghums:</u>						
Grain						
At Planting	10-20	20-40	0-20	10-20	None	Forage type sorghums have shown response from sidedressing with nitrogen during favorable moisture seasons
At Planting	10-20	20-40	0-20	15-40	None	
Forage						
At Planting	10-20	20-40	0-20	15-40	None	
Sidedressed	0-40					
<u>Wheat (grain):</u>						
Central and Eastern Okla.						
At Planting	0-40*	20-40	20-30	20-40	None	Phosphorus is a limiting factor in wheat production on most soils in Oklahoma. Use of nitrogen as top-dressing is determined by moisture conditions, previous crop, and crop prospects.
At Planting	0-30*	20-30	10-20	20-30	None	
Western Okla.						
At Planting	0-30*	20-30	10-20	20-30	None	

\*The nitrogen fertilizer used may be applied all at planting; all topdressed in the spring or fall, or it may be split, i. e., 10 lbs. at planting and 20 lbs. topdressed.

TABLE II. -- GENERAL FERTILIZER RECOMMENDATIONS FOR OKLAHOMA, 1957 continued. (1)

Crop	Nitrogen N	SOIL TEST RESULTS:				Remarks
		Phosphorus		Potassium		
		Low to Very Low	Medium to High	Low to Very Low	Medium to High	
		FERTILIZER RECOMMENDED - POUNDS PER ACRE				
		Phosphorus (P <sub>2</sub> O <sub>5</sub> )	Phosphorus (P <sub>2</sub> O <sub>5</sub> )	Potash (K <sub>2</sub> O)	Potash (K <sub>2</sub> O)	
<u>Barley (grain):</u>						
At Planting	10-20	30-40	20-30	20-40	10-20	Barley is best adapted to high fertility soils and is more sensitive to soil acidity than other small grains.
Topdressed	20-50					
<u>Oats (grain):</u>						
At Planting	10-30	20-40	10-20	10-40	None	Oats require large amounts of nitrogen to produce high yields of forage and grain, particularly during wet cold seasons and on poorly drained soils.
Topdressed	20-50					
<u>Soybeans:</u>						
At Planting	10-20	30-40	20-30	10-20	None	
<u>Peanuts:</u>						
At Planting	10-20	20-40	10-20	20-30	None	
<u>Castor Beans:</u>						
At Planting	10-20	20-40	None	10-20	None	

(continued)

TABLE II. -- GENERAL FERTILIZER RECOMMENDATIONS FOR OKLAHOMA, 1957 continued. (2)

		SOIL TEST RESULTS:				
		Phosphorus		Potassium		
		Low to Very Low	Medium to High	Low to Very Low	Medium to High	
		FERTILIZER RECOMMENDED - POUNDS PER ACRE				
Crop	Nitrogen N	Phosphorus (P <sub>2</sub> O <sub>5</sub> )	Phosphorus (P <sub>2</sub> O <sub>5</sub> )	Potash (K <sub>2</sub> O)	Potash (K <sub>2</sub> O)	Remarks
<u>Alfalfa:</u>						
At Establishment	0-15	40-80	30-60	30-60	None	Alfalfa requires well drained soils and has a high requirement for lime, phosphorus, and potassium. Established stands should be fertilized annually. Use of potash fertilizers may be lowered where subsoils are high in potassium. 20 lbs. borax per acre is recommended for heavily fertilized stands showing bronzing or chlorosis of leaves.
Topdress annually		40-80	30-60	30-60	0-30	
<u>Sweet Clover and Sericea Lespedeza:</u>						
At Establishment	0-10	40-60	10-40	20-30	None	Alfalfa and Sweet Clover are relatively efficient in utilizing the relatively insoluble phosphorus in rock phosphate. If used, rock phosphate should usually not be applied at rates less than 500 to 1000 lbs. per acre and is not recommended for soils low in organic matter or for soil with surface and subsoil reactions higher than pH 6.0. Sericea Lespedeza is more tolerant of acid low fertility soils than sweet clover.
Topdress annually on Sericea Lespedeza		30-50	0-20	10-20	None	
<u>Blue Panic:</u>						
At Establishment	10-30	20-40	10-20	10-30	None	Blue Panic is not adapted to low fertility soils and is subject to winter kill in northern sections of the state. Annual application of nitrogen is required for high forage and seed production.
Topdress annually	20-100	20-40	10-20	10-30	None	

(continued)

TABLE II. -- GENERAL FERTILIZER RECOMMENDATIONS FOR OKLAHOMA, 1957 continued. (3)

		SOIL TEST RESULTS:				
		Phosphorus		Potassium		
		Low to Very Low	Medium to High	Low to Very Low	Medium to High	
		FERTILIZER RECOMMENDED - POUNDS PER ACRE				
Crop	Nitrogen N	Phosphorus (P <sub>2</sub> O <sub>5</sub> )	Phosphorus (P <sub>2</sub> O <sub>5</sub> )	Potash (K <sub>2</sub> O)	Potash (K <sub>2</sub> O)	Remarks
<b>PERMANENT PASTURES:</b>						
<u>Warm Season Grasses:</u>						
<u>Bermuda:</u>						
<u>Weeping Lovegrass:</u>						
<u>Dallis Grass:</u>						
	At Establishment	10-20	20-40	10-20	10-40	None
	Topdress annually	30-60	20-40	0-20	10-40	None
(no legumes)						
Overseeded with:						
<u>Hop Clover: (fall)</u>						
<u>Lespedezas: (spring)</u>						
	At Establishment	None	20-40	10-20	10-20	None
	Topdress annually	None	20-40	10-20	10-20	None
Overseeded fall with:						
<u>White, Ladino,</u>						
<u>Crimson Clovers:</u>						
<u>Vetch:</u>						
	At Establishment	0-10	30-60	20-30	20-40	0-20
	Topdress annually	None	30-60	20-30	20-40	0-20
<u>Cool Season Grasses:</u>						
<u>Smooth brome, tall fescue,</u>						
<u>orchard grass, and perennial</u>						
<u>ryegrass seeded (fall) with</u>						
<u>alfalfa, vetch, white, ladino,</u>						
<u>red or crimson clovers:</u>						
	At Establishment	10-30	40-80	20-40	20-60	0-20
	Topdress annually	20-40	40-80	20-40	20-60	0-20
<b>GRASS SEED PRODUCTION:</b> Topdressing with 30 to 60 lbs. of nitrogen and 20 to 40 lbs. of phosphorus (P <sub>2</sub> O <sub>5</sub> ) in established stands is recommended for seed production of Weeping Lovegrass, Smooth Brome, Tall Fescue Orchard Grass, and Blue Panic.						

(continued)



TABLE II. -- GENERAL FERTILIZER RECOMMENDATIONS FOR OKLAHOMA, 1957 continued (4)

SOIL TEST RESULTS:						
Phosphorus			Potassium			
Low to	Medium	Low to	Medium to			
Very Low	to High	Very Low	High			
FERTILIZER RECOMMENDED - POUNDS PER ACRE						
Crop	Nitrogen N	Phosphorus (P2O5)	Phosphorus (P2O5)	Potash (K2O)	Potash (K2O)	Remarks
<b>TEMPORARY PASTURE:</b>						
<u>Small Grains and common ryegrass alone and with vetch, crimson or sweet clover:</u>						
At Establishment	10-40	30-60	10-30	20-40	0-20	Barley is best adapted to high fertility soils and is more sensitive to acid soil conditions than other small grains. Oats require large amounts of nitrogen to produce high yields of forage particularly during cool wet seasons on poorly drained soils.
Small Grains only for pasture, topdress	20-50					
<u>Sudan Grass:</u>						
At Establishment	10-40	20-40	10-20	10-20	0-20	Sudan grass produces best on high fertility soils and gives good response to nitrogen fertilization.
Sidedress	20-60					