

RESEARCH ARTICLE

Open Access

Oral re-vaccination of Eurasian wild boar with *Mycobacterium bovis* BCG yields a strong protective response against challenge with a field strain

Christian Gortazar^{1*†}, Beatriz Beltrán-Beck^{1†}, Joseba M Garrido², Alicia Aranaz³, Iker A Sevilla², Mariana Boadella¹, Konstantin P Lyashchenko⁴, Ruth C Galindo¹, Vidal Montoro¹, Lucas Domínguez⁵, Ramón Juste² and Jose de la Fuente^{1,6}

Abstract

Background: Field vaccination trials with *Mycobacterium bovis* BCG, an attenuated mutant of *M. bovis*, are ongoing in Spain, where the Eurasian wild boar (*Sus scrofa*) is regarded as the main driver of animal tuberculosis (TB). The oral baiting strategy consists in deploying vaccine baits twice each summer, in order to gain access to a high proportion of wild boar piglets. The aim of this study was to assess the response of wild boar to re-vaccination with BCG and to subsequent challenge with an *M. bovis* field strain.

Results: BCG re-vaccinated wild boar showed reductions of 75.8% in lesion score and 66.9% in culture score, as compared to unvaccinated controls. Only one of nine vaccinated wild boar had a culture-confirmed lung infection, as compared to seven of eight controls. Serum antibody levels were highly variable and did not differ significantly between BCG re-vaccinated wild boar and controls. Gamma IFN levels differed significantly between BCG re-vaccinated wild boar and controls. The mRNA levels for IL-1b, C3 and MUT were significantly higher in vaccinated wild boar when compared to controls after vaccination and decreased after mycobacterial challenge.

Conclusions: Oral re-vaccination of wild boar with BCG yields a strong protective response against challenge with a field strain. Moreover, re-vaccination of wild boar with BCG is not counterproductive. These findings are relevant given that re-vaccination is likely to happen under real (field) conditions.

Keywords: Bacillus Calmette Guerin, *Sus scrofa*, Tuberculosis, Vaccination and challenge, Wildlife disease control

Background

Wildlife vaccination is increasingly being explored as a tuberculosis (TB) control tool in all major wildlife reservoirs. An attenuated mutant of *M. bovis*, *Mycobacterium bovis* BCG (bacille Calmette Guerin [1]), is often the vaccine used in wildlife vaccination trials worldwide [2-8]. In countries such as UK and the Republic of Ireland, Eurasian badger (*Meles meles*) vaccination field trials are becoming part of the TB control strategies [9].

Field vaccination trials are also ongoing in Spain, where the Eurasian wild boar (*Sus scrofa*) is regarded as the main driver of animal TB at the wildlife-livestock interface [10-12].

In Mediterranean Spain, modeling of tuberculosis in wild boar populations predicted that over 70% of the annual cohort of wild boar piglets would be needed to be vaccinated in order to significantly reduce the prevalence of the disease and eventually eradicate the infection [13]. It has been demonstrated that the most efficient oral vaccine bait deployment strategy in this area consists in using selective piglet feeders and deploying vaccine baits both at the start (late June/early July) and the end of summer (late August/early September), in order to gain

* Correspondence: christian.gortazar@uclm.es

†Equal contributors

¹SaBio IREC (CSIC-UCLM-JCCM), Ronda de Toledo s/n, 13005 Ciudad Real, Spain

Full list of author information is available at the end of the article

access to a high proportion of early and late born wild boar piglets respectively, avoiding the hottest period from late July to early August [14,15]. However, with this strategy of a double delivery in summer, it would be probable that some individuals actually gain access to vaccine baits twice in a season. This possibility of some wild boar ingesting vaccine baits in both vaccination campaigns raises the question of the effects of re-vaccination in wild boar. In human beings, revaccination with BCG confers only a modest advantage over single vaccination [16,17]. Among cattle and wildlife MTC hosts, there are controversial studies regarding the use of a single dose or the need of a booster dose of the vaccine to achieve a better vaccination protocol (Table 1). Re-vaccination has the potential to be counter-productive, and such risks needed to be assessed in wild boar as an integral part of vaccine safety. Although adverse effects of re-vaccination or over-dosage of BCG in wild boar were not expected, we were interested in assessing the effects of re-vaccination with a 52 day interval imitating field conditions.

The aim of this study was to assess the response of wild boar to re-vaccination with BCG and to subsequent challenge with an *M. bovis* field strain, under experimental conditions. Based on current knowledge, we expected a protective response similar to those recorded in previous single dose BCG vaccination experiments [2,22].

Methods

Experiment design

Seventeen 3-4-month-old wild boar piglets were bought in a commercial farm known to be free of mycobacterial lesions at slaughter and with a fully negative ELISA test [23]. The animals were housed in class III bio-containment facilities with ad libitum food and water. Wild boar piglets were randomly assigned to treatment (BCG vaccinated animals) or control groups (unvaccinated animals). The *M.*

bovis BCG Danish reference strain (CCUG 27863) was cultured on Coletsos medium (Biomerieux, France), and prepared as described for previous experiments [22]. Vaccine containing 10^6 cfu was administered orally in baits designed for wild boar piglets [24]. For the challenge, 5 ml of a suspension containing 10^5 colony forming units (cfu) of an *M. bovis* field strain were administered by the oropharyngeal route as described in previous experiments [2,22]. The isolate used for challenge was originally isolated from a naturally infected wild boar and identified as *M. bovis* spoligotype profile SB0339 according to the *M. bovis* Spoligotype Database website (www.mbovis.org).

The animals were handled five times during the experiment, including vaccination (T0, day 1), re-vaccination 52 days after the first vaccination (T1, day 52), challenge 74 days after re-vaccination (T2, day 126), one blood sampling two months after challenge (T3, day 185) and necropsy four months after challenge (T4, day 255). Handling procedures and sampling frequency were designed to reduce stress and health risks for subjects, according to European (86/609) and Spanish laws (R.D. 223/1988, R.D. 1021/2005). The protocol was approved by the Committee on the Ethics of Animal Experiments of the Regional Agriculture Authority (Diputación Foral de Vizcaya, Permit Number: BFA10.373 (27/19/2010)).

Sampling, pathology and microbiology

Blood samples were collected at T0-T4 time points for RNA extraction from PBMC and serum preparation. Animals were anesthetized by intramuscular injection of Zoletil (Virbac, Esplugues de Llobregat, Spain) and euthanized by captive bolt. At necropsy, all wild boar were carefully inspected for the presence of macroscopic TB-compatible lesions. Samples for culture were immediately processed and copies frozen at -80°C for mRNA isolation. TB-compatible lesions were classified based on

Table 1 BCG vaccination studies comparing the use of both one single dose or repeated doses of the vaccine in cattle and wildlife

	Dose	Interval	Route	Reference	Effect of boosting
Red deer (<i>Cervus elaphus</i>)	2 doses of 2.5×10^6 cfu BCG Pasteur	8 weeks	Subcutaneous	[5]	Two doses of vaccine were superior to one dose producing protection against infection.
Red deer (<i>Cervus elaphus</i>)	2 doses of 2×10^6 cfu BCG Pasteur	6 weeks	Subcutaneous	[18]	Single-dose vaccines protected against disease. Boosting was required to protect against infection.
White-tailed deer (<i>Odocoileus virginianus</i>)	2 doses of 10^7 cfu BCG Pasteur	6 weeks	Subcutaneous	[19]	Pathology scores were lower in deer receiving 2 doses of <i>M. bovis</i> BCG compared to unvaccinated ones or those who received a single dose.
Brush-tailed possum (<i>Trichosurus vulpecula</i>)	12 doses of 1×10^8 ; 2 doses of 1×10^9 cfu BCG Pasteur	1 Week (12 doses); 6 weeks (2 doses)	Intranasal aerosol and conjunctival instillation	[20]	The group vaccinated 12 times showed the greatest level of protection. Revaccination after a period of 6 weeks had no beneficial and no deleterious effects as compared to the protection induced by a single dose of vaccine.
Cattle	2 doses of 10^6 CFU BCG Pasteur	6 weeks	Subcutaneous	[21]	Significantly less protection than those vaccinated only once. Revaccination reduced the level of protection induced by a single vaccination.

lesion distribution and lesion intensity, and scored as previously described [2]. Briefly, lymph nodes and the oropharyngeal tonsils were scored as 0 (No visible lesion), 1 (1–2 small (<1 cm) caseous foci), 2 (Several small foci), 3 (Same and at least one lesion >1 cm) or 4 (Diffusely affected); lung lobes were scored as 0 (No visible lesion), 1 (Few small lesions), 2 (Numerous or clustered small lesions with some coalescence), 3 (Densely clustered small lesions), 4 (Same and at least one large lesion) or 5 (Two or more large lesions). Visceral organ lesions were scored as 0 (No visible lesion), 1 (1–2 mm foci scattered throughout organ) or 2 (5–10 mm diameter clusters of 1–2 mm foci or single focus >1 cm diameter). Lymphoid tissues and samples of lung tissue were cultured for mycobacteria and scored (total number of culture-positive samples) as described in [22]. All isolates were spoligotyped in order to confirm the strain [25]. Tonsil samples were flash frozen and stored in liquid N until used for gene expression studies.

Serology and interferon tests

Serum samples were tested for anti-PPD immunoglobulin G (IgG) antibodies by means of an in-house ELISA using bovine tuberculin purified protein derivative (bovine PPD; CZ Veterinaria SL, Porriño, Lugo, Spain) as antigen and protein G horseradish peroxidase (Sigma-Aldrich, Madrid, Spain) as a conjugate applying the protocol described by Boadella et al. [23]. Serum samples were also tested by the DPP technology developed by Chembio Diagnostic Systems, Inc. using selected *M. bovis* antigens. Briefly, the presence and intensity of either of the 2 separate test lines (T1, MPB83 antigen; T2, CFP10/ESAT-6 fusion protein) were evaluated by a DPP optical reader (in relative light units, RLU) [23]. Blood samples taken at times 2, 3 and 4 were also used for detection of the IFN-gamma response in presence of avian and bovine PPD, as described previously [22].

RNA isolation and real time RT-PCR

Total RNA was extracted from wild boar PBMC and tonsils using TRI reagent (Sigma, Madrid, Spain) following manufacturer's recommendations. RNA was used for real-time RT-PCR analysis of mRNA levels of selected genes in individual samples. Selected genes are involved in innate immunity (complement component 3, *C3* and interleukin 1-beta, *IL-1b*) and methylmalonyl CoA mutase, *MUT*. Real-time RT-PCR was performed with gene-specific primers (*C3*, SsC3-L: acaaattgaccagcgtagg and SsC3-R: gcacgtcctgtctgtactga; *IL-1b*, SsIL1beta-L: ccaaagaggacatg gagaa and SsIL1beta-R: ttatatcttggcccttg; *MUT*, SsMUT-L: gtttgccaacggtaaaagt and SsMUT-R: aatgagacct caaggcagcat) using the One-Step RT-PCR Kit with SYBR Green and the CFX thermal cycler (Bio-Rad, Hercules, CA, USA) following manufacturer's recommendations.

Control reactions were performed using the same procedures, but without RT to monitor DNA contamination in the RNA preparations and without RNA added to monitor contamination of the PCR reaction. A dissociation curve was run at the end of RT-PCR reaction to ensure that only one amplicon was formed and that the amplicon denatured consistently at the same temperature range for every sample [26]. The mRNA values were normalized against *S. scrofa* cyclophilin (SsCyclophilin-L: agcactggggagaaaggatt and SsCyclophilin-R: ctggcagtgcataatgaaa), β-actin (Ss-BactinF: ggaccgtaccgtaccatca and Ss-BactinR: ggccagtcgtacgtttcat) and GAPDH (Ss-GAPDH: gtcgggtgtggatctgacct and Ss-GAPDH: agtttgacgaagtggctgt) using the genNorm ddCT method [27].

Results

No clinical signs such as wasting or cough were recorded during the experiment. Figure 1 presents the pathology scores and total culture scores. BCG re-vaccinated wild boar showed reductions of 75.8% and 66.9%, respectively, as compared to controls (U tests; $Z > 3.2$, $p < 0.01$). Two BCG re-vaccinated wild boar had no positive culture. The mean (\pm SE) thorax culture scores were 0.22 (\pm 0.3) for vaccinated and 2.25 (\pm 0.3) for control wild boar (90.2% reduction; U test; $Z = 3.1$, $p < 0.01$). Only one of nine vaccinated wild boar had a culture-confirmed lung infection, as compared to seven of eight controls (Fisher's test; $p = 0.003$).

Serum antibody levels were highly variable and did not differ significantly between BCG re-vaccinated wild boar and controls (ANOVA; Treatment effect $F_{1, 45} < 2.71$, $p > 0.05$). Gamma IFN levels differed significantly between BCG re-vaccinated wild boar and controls (ANOVA; Treatment effect $F_{1, 39} = 6.08$, $p < 0.05$). The gamma IFN response to bPPD was undetectable at T2. A peak was recorded at T3 both in BCG re-vaccinated wild boar (mean OD \pm SE 6.9 ± 1.7) and in controls (11.4 ± 1.8), slightly declining in both groups thereafter (Figure 2).

The mRNA levels for IL-1b, *C3* and *MUT* were significantly higher in vaccinated wild boar when compared to controls at T2 and decreased after mycobacterial challenge (Figure 3A). Only IL-1b mRNA levels remained higher in vaccinated animals when compared to controls until the end of the experiments (Figure 3A). In tonsils, only *MUT* mRNA levels were significantly higher in vaccinated than in control animals at the end of the experiment (Figure 3B).

Discussion

This experiment confirmed that oral BCG re-vaccination of wild boar induces a strong protective response against challenge with an *M. bovis* field strain. This response includes lower lesion and culture scores; lower gamma




Figure 1 Wild boar TB lesion scores (left) and culture scores (right) at necropsy. The solid lines show the median values.

IFN levels and higher IL-1 β , C3 and MUT mRNA levels in vaccinated wild boar.

The main difference between this experiment and the previous vaccine and challenge trials in wild boar was re-vaccination. Although no single vaccination controls were included, the protective response recorded after re-vaccination (76% lesion score, 67% culture score), was higher than those recorded in two previous single dose BCG vaccination experiments, i.e. 56% reduction in

lesion score, 50% in culture score [2]; 52% reduction in lesion score, 9% in culture score [22]. This animal experiment was run along with another one testing a new, heat-inactivated vaccine, and the number of available experimental wild boar and the available housing space did not allow including a single-vaccination BCG group. Serum antibody and gamma IFN responses were essentially similar to those recorded in these single vaccination experiments. These levels of protection were better




Figure 2 Mean optical density (OD) readings of the gamma interferon ELISA in BCG re-vaccinated (solid line) and control (dashed line) wild boar. Vertical bars represent 95% confidence intervals.




Figure 3 Gene expression in response to BCG vaccination in wild boar. The mRNA levels of selected genes were analyzed by real-time RT-PCR in (A) PBMC of vaccinated and control wild boar collected at T0 (before vaccination), T2 (before challenge) and T4 (at necropsy) or (B) tonsils at T4. The mRNA levels were normalized against *S. scrofa* cyclophilin, β-actin and GAPDH and normalized Ct values were represented as Ave + S.D. in arbitrary units and compared between groups by Student's t-test with unequal variance (*P ≤ 0.05).

than those documented for BCG-Pasteur re-vaccinated African buffaloes (*Syncerus caffer*), in which revaccination gave no differences in the protection for vaccinated animals compared to unvaccinated animals [4]. In ferrets (*Mustela furo*), two doses of 5×10^8 cfu of BCG-Pasteur given orally in a 4 week interval produced a significant reduction in pathology after oral challenge with virulent *M. bovis*, decreasing the mean bacterial count for retropharyngeal LN tissue of culture-positive vaccinated animals to 70% [7].

It is known that factors such as the variation of the period between the administration of the two doses of the vaccine could affect the protective efficacy of the vaccination. For example, the reduction of this period from 8 to 4 weeks in vaccinated red deer did not affect the protection induced by the vaccine, whereas increasing this period to 43 weeks largely ablated protection [28]. In our study, wild boar vaccination took place 52 days after the first vaccination (between 7-8 weeks), which imitates field conditions.

Another important factor that could affect the protection of the vaccine is the dose. Low and medium booster doses (10^4 - 10^7 cfu; 6-8 weeks interval) of BCG-Pasteur produced significant protection against infection and disease in vaccinated red deer, but higher booster doses (5×10^8 cfu) provided protection against disease but not against infection [5,18]. In our case, a dose of 10^6 cfu was used, again imitating field protocols.

Molecular characterization of host-pathogen interactions identified wild boar genes such as MUT, C3 and other innate and adaptive immune response genes involved in

resistance to mycobacterial infection [29-33]. As in previous experiments, C3 and MUT mRNA levels rose after vaccination and decreased after challenge [22]. The up-regulation of genes encoding pro-inflammatory cytokines such as pro-IL-1b probably stimulates the production of C3 which may contribute to the protective response to BCG vaccination in wild boar.

Conclusions

Oral re-vaccination of wild boar with BCG yields a strong protective response against challenge with a field strain. One important conclusion of this study is that re-vaccination of wild boar with BCG is not counterproductive as suggested for cattle [34]. This finding is relevant given that wild boar TB is increasingly reported [35,36] and that re-vaccination is likely to happen in the ongoing field vaccination trials.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

BB-B, JMG, AA, LD, RJ, JdLF and CG: contributed to the conception, design, and data collection, laboratory work, drafting and writing of the manuscript. IAS, MB, KPL, RCG and VM: contributed to laboratory work, data analysis and drafting of the manuscript. All authors have read and approved the final manuscript.

Acknowledgements

Carlos Martín, Universidad de Zaragoza, kindly provided the vaccine BCG strain. Many colleagues at SaBio IREC, NEIKER and VISAVENT granted valuable technical help. This is a contribution to EU FP7 grant 613779 WildTBVac and to MINECO Plan Nacional grant AGL2011-30041 and FEDER.

Author details

¹SaBio IREC (CSIC-UCLM-JCCM), Ronda de Toledo s/n, 13005 Ciudad Real, Spain. ²NEIKER-Tecnalia, Animal Health Department, C/Berreaga 1, E-48160 Derio, Bizkaia, Spain. ³Departamento de Sanidad Animal, Facultad de Veterinaria, Universidad Complutense de Madrid, Avenida de Puerta de Hierro s/n, 28040 Madrid, Spain. ⁴Chembio Diagnostic Systems Inc., Medford, NY 11763, USA. ⁵Centro de Vigilancia Sanitaria Veterinaria (VISAVET), Facultad de Veterinaria, Universidad Complutense de Madrid, Avenida de Puerta de Hierro s/n, 28040 Madrid, Spain. ⁶Department of Veterinary Pathobiology, Center for Veterinary Health Sciences, Oklahoma State University, Stillwater, OK 74078, USA.

Received: 5 February 2014 Accepted: 22 April 2014

Published: 26 April 2014

References

- Calmette A: *La vaccination preventive contre la tuberculose*. Paris: Masson; 1927.
- Ballesteros C, Garrido JM, Vicente J, Romero B, Galindo RC, Minguijón E, Villar M, Martín-Hernando MP, Sevilla I, Juste R, Aranaz A, de la Fuente J, Gortázar C: First data on Eurasian wild boar response to oral immunization with BCG and challenge with a *Mycobacterium bovis* field strain. *Vaccine* 2009, **27**:6662–6668.
- Corner LAL, Costello E, O'Meara D, Lesellier S, Aldwell FE, Singh M, Hewinson RG, Chambers MA, Gormley E: Oral vaccination of badgers (*Meles meles*) with BCG and protective immunity against endobronchial challenge with *Mycobacterium bovis*. *Vaccine* 2010, **28**:6265–6272.
- de Klerk LM, Michel AL, Bengis RG, Kriek NPJ, Godfroid J: BCG vaccination failed to protect yearling African buffaloes (*Synacerus caffer*) against experimental intratonsilar challenge with *Mycobacterium bovis*. *Vet Immunol Immunopathol* 2010, **137**:84–92.
- Griffin JF, Mackintosh CG, Slobbe L, Thomson AJ, Buchan GS: Vaccine protocols to optimise the protective efficacy of BCG. *Tuber Lung Dis* 1999, **79**:135–143.
- Nol P, Palmer MV, Waters WR, Aldwell FE, Buddle BM, Triantis JM, Linke LM, Phillips GE, Thacker TC, Rhyau JC, Dunbar MR, Salman MD: Efficacy of oral and parenteral routes of *Mycobacterium bovis* bacilli Calmette-Guerin vaccination against experimental bovine tuberculosis in white-tailed deer (*Odocoileus virginianus*): A feasibility study. *J Wildl Dis* 2008, **44**:247–259.
- Qureshi T, Labes RE, Cross ML, Griffin JFT, Mackintosh CG: Partial protection against oral challenge with *Mycobacterium bovis* in ferrets (*Mustela furo*) following oral vaccination with BCG. *Int J Tuberc Lung Dis* 1999, **3**:1025–1033.
- Tompkins DM, Ramsey DSL, Cross ML, Aldwell FE, De Lisle GW, Buddle BM: Oral vaccination reduces the incidence of tuberculosis in free-living brushtail possums. *Proc R Soc Lond B Biol Sci* 2009, **276**:2987–2995.
- Gormley E, Corner LAL: Control strategies for wildlife tuberculosis in Ireland. *Transbound Emerg Dis* 2013, **60**:128–135.
- Gortázar C, Delahay RJ, McDonald RA, Boadella M, Wilson GJ, Gavier-Widen D, Acevedo P: The status of tuberculosis in European wild mammals. *Mammal Rev* 2012, **42**:193–206.
- Gortázar C, Vicente J, Boadella M, Ballesteros C, Galindo RC, Garrido J, Aranaz A, de la Fuente J: Progress in the control of bovine tuberculosis in Spanish wildlife. *Vet Microbiol* 2011, **151**:170–178.
- Naranjo V, Gortázar C, Vicente J, de la Fuente J: Evidence of the role of European wild boar as a reservoir of *Mycobacterium tuberculosis* complex. *Vet Microbiol* 2008, **127**:1–9.
- Anderson LG, Gortázar C, Vicente J, Hutchings MR, White PCL: Modelling the effectiveness of vaccination in controlling bovine tuberculosis in wild boar. *Wildlife Res* 2013, **40**:367–376.
- Ballesteros C, Vicente J, Carrasco-García R, Mateo R, de la Fuente J, Gortázar C: Specificity and success of oral-bait delivery to Eurasian wild boar in Mediterranean woodland habitats. *E J Wildlife Res* 2011, **57**:749–757.
- Beltrán-Beck B, Ballesteros C, Vicente J, De La Fuente J, Gortázar C: Progress in oral vaccination against tuberculosis in its main wildlife reservoir in Iberia, the Eurasian wild boar. *Vet Med Int* 2012, doi:10.1155/2012/978501.
- Barreto ML, Pereira SM, Pilger D, Cruz AA, Cunha SS, Sant'Anna C, Ichihara MY, Genser B, Rodrigues LC: Evidence of an effect of BCG revaccination on incidence of tuberculosis in school-aged children in Brazil: Second report of the BCG-REVAC cluster-randomised trial. *Vaccine* 2011, **29**:4875–4877.
- Whelan KT, Pathan AA, Sander CR, Fletcher HA, Poultney I, Alder NC, Hill AVS, McShane H: Safety and immunogenicity of boosting BCG vaccinated subjects with BCG: Comparison with boosting with a new TB vaccine, MVA85A. *PLoS One* 2009, **4**:e5934. doi:10.1371/journal.pone.0005934.
- Griffin JF: Veterinary tuberculosis vaccine development. *Clin Infect Dis* 2000, **30**:S223–S228.
- Palmer MV, Thacker TC, Waters WR: Vaccination of white-tailed deer (*Odocoileus virginianus*) with *Mycobacterium bovis* bacillus Calmette Guérin. *Vaccine* 2007, **25**:6589–6597.
- Corner LAL, Buddle BM, Pfeiffer DU, Morris RS: Vaccination of the brushtail possum (*Trichosurus vulpecula*) against *Mycobacterium bovis* infection with bacille Calmette-Guerin: the response to multiple doses. *Vet Microbiol* 2002, **84**:327–336.
- Buddle BM, Wedlock DN, Parlame NA, Corner LAL, de Lisle GW, Skinner MA: Revaccination of Neonatal Calves with *Mycobacterium bovis* BCG reduces the level of protection against bovine tuberculosis induced by a single vaccination. *Infect Immun* 2003, **71**:6411–6419.
- Garrido JM, Sevilla IA, Beltrán-Beck B, Minguijón E, Ballesteros C, Galindo RC, Boadella M, Lyashchenko KP, Romero B, Geijo MV, Ruiz-Fons F, Aranaz A, Juste RA, Vicente J, de la Fuente J, Gortázar C: Protection against Tuberculosis in Eurasian Wild Boar Vaccinated with Heat-inactivated *Mycobacterium bovis*. *PLoS One* 2011, **6**:e24905. doi:10.1371/journal.pone.0024905.
- Boadella M, Lyashchenko K, Greenwald R, Esfandiari J, Jaroso R, Carta T, Garrido JM, Vicente J, de la Fuente J, Gortázar C: Serological tests for detecting antibodies against *Mycobacterium bovis* and *Mycobacterium avium* subspecies paratuberculosis in Eurasian wild boar (*Sus scrofa scrofa*). *J Vet Diagn Invest* 2011, **23**:77–83.
- Ballesteros C, Gortázar C, Canales M, Vicente J, Lasagna A, Gamarr JA, Carrasco-García R, Fuente J: Evaluation of baits for oral vaccination of European wild boar piglets. *Res Vet Sci* 2009, **86**:388–393.
- Kamerbeek J, Schouls L, Kolk A, van Agterveld M, van Soolingen D, Kuijper S, Bunschoten A, Molhuizen H, Shaw R, Goyal M, van Embden J: Simultaneous detection and strain differentiation of *Mycobacterium tuberculosis* for diagnosis and epidemiology. *J Clin Microbiol* 1997, **35**:907–914.
- Ririe KM, Rasmussen RP, Wittwer CT: Product differentiation by analysis of DNA melting curves during the polymerase chain reaction. *Anal Biochem* 1997, **245**:154–160.
- Livak KJ, Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-ΔΔCT} method. *Methods* 2001, **25**:402–408.
- Griffin JF, Mackintosh CG, Rodgers CR: Factors influencing the protective efficacy of a BCG homologous prime-boost vaccination regime against tuberculosis. *Vaccine* 2006, **24**:835–845.
- Naranjo V, Ayoubi P, Vicente J, Ruiz-Fons F, Gortázar C, Kocan KM, de la Fuente J: Characterization of selected genes upregulated in non-tuberculous European wild boar as possible correlates of resistance to *Mycobacterium bovis* infection. *Vet Microbiol* 2006, **116**:224–231.
- Naranjo V, Höfle U, Vicente J, Martin MP, Ruiz-Fons F, Gortázar C, Kocan KM, de la Fuente J: Genes differentially expressed in oropharyngeal tonsils and mandibular lymph nodes of tuberculous and non-tuberculous European wild boars naturally exposed to *Mycobacterium bovis*. *FEMS Immunol Med Microbiol* 2006, **98**:77–85.
- Naranjo V, Villar M, Martín-Hernando MP, Vidal D, Höfle U, Gortázar C, Kocan KM, Vázquez J, de la Fuente J: Proteomic and transcriptomic analyses of differential stress/inflammatory responses in mandibular lymph nodes and oropharyngeal tonsils of European wild boars naturally infected with *Mycobacterium bovis*. *Proteomics* 2007, **7**:220–231.
- Galindo RC, Ayoubi P, Naranjo V, Gortázar C, de la Fuente J: Gene expression profiles of European wild boar naturally infected with *Mycobacterium bovis*. *Vet Immunol Immunopathol* 2009, **129**:119–125.
- de la Lastra JM P, Galindo RC, Gortázar C, Ruiz-Fons F, Aranaz A, de la Fuente J: Expression of immunoregulatory genes in peripheral blood mononuclear cells of European wild boar immunized with BCG. *Vet Microbiol* 2009, **134**:334–339.
- Buddle BM, Pollock JM, Skinner MA, Wedlock DN: Development of vaccines to control bovine tuberculosis in cattle and relationship to vaccine development for other intracellular pathogens. *Int J Parasitol* 2003, **33**:555–566.

35. Muñoz-Mendoza M, Marreros N, Boadella M, Gortázar C, Menéndez S, de Juan L, Bezos J, Romero B, Copano MF, Amado J, Sáez JL, Mourelo J, Balseiro A: Wild boar tuberculosis in Iberian Atlantic Spain: a different picture from Mediterranean habitats. *BMC Vet Res* 2013, **9**:176.
36. Richomme C, Boadella M, Courcoul A, Durand B, Drapeau A, Corde Y, Hars J, Payne A, Fediaevsky A, Boschioli ML: Exposure of Wild Boar to *Mycobacterium tuberculosis* Complex in France since 2000 Is Consistent with the Distribution of Bovine Tuberculosis Outbreaks in Cattle. *PLoS One* 2013, **8**:e77842. doi:10.1371/journal.pone.0077842.

doi:10.1186/1746-6148-10-96

Cite this article as: Gortazar et al.: Oral re-vaccination of Eurasian wild boar with *Mycobacterium bovis* BCG yields a strong protective response against challenge with a field strain. *BMC Veterinary Research* 2014 **10**:96.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

