



Current Report

EXTENSION

Oklahoma Cooperative Extension Fact Sheets are also available on our website at:
extension.okstate.edu

Management of Insect and Mite Pests in Sunflowers

Tom A. Royer
 Extension Entomologist

Unmanaged sunflower pests can reduce yield and quality of seed and oil. Pesticides should not be used as a substitute for good agronomic practices or as “preventative insurance” because this approach can cause pest resurgence issues and is rarely economically or environmentally justifiable. Many sunflower pest problems can be avoided by developing an Integrated Pest Management (IPM) plan that includes preventive pest management practices, such as planting high-quality, vigorous, Oklahoma-proven hybrid seed, planting it at the proper time for optimal health and yield, providing proper fertilization and weed control, and using crop rotations.

The information herein is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Cooperative Extension Service is implied.

Pesticide recommendations in this publication were correct as of the “Modified Date” but always check the label that came with the purchased insecticide for the most current rates and restrictions

The first name listed is the trade name of a product

registered for use in corn for the listed pest. The name in (parentheses) listed below the trade name is the name of the active ingredient. The active ingredient name is provided because in many cases, there are other registered products containing the same active ingredient that may cost less, so producers should compare prices.

The number [in brackets] following a product is its Mode of Action number [MOA]. The more frequently insecticides with the same MOA are used, the more likely resistance will occur. This number provides an easy way to select different modes of action to avoid selecting for pests that are resistant to a certain mode of action.

Refer to the following publications for additional information on sunflower pest management.

- EPP-7196 Grasshopper Management in Rangeland, Pastures and Crops (OSU)
- MF-2384 High Plains Sunflower Production Handbook (Kansas State) <http://www.bookstorew.ksre.ksu.edu/pubs/mf2384.pdf>

Management of Insect and Mite Pests in Sunflowers

Pest, Damage and Treatment Threshold	Insecticide, Formulation, [MOA Group] & (Active Ingredient)	Rate of Product per Acre	Comments
Cutworms (black, granulate, sandhill) Striped or solid colored, robust caterpillars that “roll” up when disturbed, and prefer to live under ground. Damage: Cutworms generally feed at night and live under the soil during the day. Plants will be cut at or slightly above the soil level, causing stand reductions. Threshold: Scout fields at seedling emergence. Threshold is one cutworm per square foot	Asana XL [3] (esfenvalerate)	5.8 to 9.6 fl oz (0.03 to 0.05 lb ai/A)	28 day waiting period for harvest; do not graze
	Baythroid XL [3] (beta-cyfluthrin)	0.8 to 1.6 fl oz (0.007 to 0.013 lb ai/A)	30 day waiting period for harvest; do not graze
	Besiege [28,3] (lambda-cyhalothrin + chlorantraniliprole)	5.0 to 8.0 fl oz	45 waiting period for harvest. Do not use adjuvant with application. Follow drift precautions to protect pollinators
	Delta Gold [3] (deltamethrin)	1.0 to 1.5 fl oz (0.012 to 0.018 lb ai/A)	21 day waiting period for harvest; do not graze
	Karate w Zeon [3] (lambda-cyhalothrin)	0.96 to 1.60 fl oz (0.015 to 0.025 lb ai/A)	45 day waiting period for harvest

combined with a 25% stand reduction. Treat when worms are less than ½ inch long.	Mustang MAXX EC [3] (zeta-cypermethrin)	1.28 to 4 fl oz (0.008 to 0.025 lb ai/A)	30 day waiting period for harvest; do not graze
	Proaxis 0.5 CS [3] (gamma-cyhalothrin)	1.92 to 3.2 fl oz (0.0075 to 0.0125 lb ai/A)	45 day waiting period for harvest
	Sevin XLR [1A] (carbaryl)	1.5 quarts (1.5 lb ai/A)	30 day wait for grazing, 60 days for harvest
	Tombstone [3] (cyfluthrin)	0.80 to 1.60 fl oz (0.013 to 0.025 lb ai/A)	30-day waiting period for har- vest.
	Warrior II w Zeon [3] (lambda-cyhalothrin)	0.96 to 1.60 fl oz (0.015 to 0.025 lb ai/A)	45 day waiting period for harvest
Grasshopper 1-2 inches long, outer wings leathery, inner wings clear or colored. Enlarged hind legs designed for jumping. Damage: Chew leaves, leaving ragged edges or completely chewing leaf blade. Damage developing seed heads, causing yield loss. Threshold: See EPP-7196: Grasshopper Management in Rangeland, Pastures, and Crops	Asana XL [3] (esfenvalerate)	5.8 to 9.6 fl oz (0.03 to 0.05 lb ai/A)	28 day waiting period for harvest for harvest; do not graze
	Baythroid XL [3] (beta-cyfluthrin)	2.0 to 2.8 fl oz (0.016 to 0.022 lb ai/A)	30 day waiting period for harvest for harvest; do not graze
	Besiege [28,3] (lambda-cyhalothrin + chlorantra- niliprole)	6.0-10.0 fl oz	45 waiting period for harvest. Do not use adjuvant with application. Follow drift precautions to protect pollinators
	Delta Gold [3] (deltamethrin)	1.0 to 1.5 fl oz (0.012 to 0.018 lb ai/A)	21 day waiting period for harvest for harvest; do not graze
	Karate w Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	45 day waiting period for harvest
	Mustang MAXX EC [3] (zeta-cypermethrin)	2.6 to 4 fl oz (0.016 to 0.025 lb ai/A)	30 day waiting period for harvest for harvest; do not graze
	Proaxis 0.5 CS [3] (gamma-cyhalothrin)	2.56 to 3.84 fl oz (0.01 to 0.015 lb ai/A)	45 day waiting period for harvest
	Vantacor [28] (chlorantraniliprole)	0.7 to 1.7 fl oz (0.026 to 0.065 lb ai/A)	1 day PHI
	Warrior II w Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	45 day waiting period for harvest
Foliar-feeding caterpillars (painted lady, woolly bear) Various caterpillars, painted lady and woolly bear caterpillars have hairy bodies. Damage: Feed on leaves Threshold: Treat when defolia- tion exceeds 25% and caterpil- lars are still present.	Besiege [28,3] (lambda-cyhalothrin + chlorantra- niliprole)	6.0 to 10.0 fl oz	45 waiting period for harvest. Do not use adjuvant with application. Follow drift precautions to protect pollinators
	Karate w Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	45 day waiting period for harvest
	Mustang MAXX EC [3] (zeta-cypermethrin)	1.28 to 4 fl oz (0.008 to 0.025 lb ai/A)	30 day waiting period for harvest Check label for species labeled rate
	Proaxis 0.5 CS [3] (gamma-cyhalothrin)	2.56 to 3.84 fl oz (0.01 to 0.015 lb ai/A)	45 day waiting period for harvest
	Warrior II w Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	45 day waiting period for harvest
Seed weevils (Red and Gray) Reddish weevil about 1/8-inch- long, and grey weevil about 1/4-inch long. Larvae are white, about 1/6-inch long when mature. Damage: Larvae feed inside seed, cut exit hole when mature, and burrow into ground. Threshold: Scout for red weevil when 85% of plants are past R-4	Asana XL [3] (esfenvalerate)	5.8 to 9.6 fl oz (0.03 to 0.05 lb ai/A)	28 day waiting period for harvest for harvest; do not graze
	Baythroid XL [3] (beta-cyfluthrin)	2.0 to 2.8 fl oz (0.016 to 0.022 lb ai/A)	30 day waiting period for harvest; do not graze
	Besiege [28,3] (lambda-cyhalothrin + chloran- traniliprole)	6.0 to 10.0 fl oz	45 waiting period for harvest. Do not use adjuvant with application. Follow drift precautions to protect pollinators
	Delta Gold [3] (deltamethrin)	1.0 to 1.5 fl oz (0.012 to 0.018 lb ai/A)	21 day waiting period for harvest

growth stage. Treat when counts exceed 10 weevils per head. Continue to scout to determine if second spray is needed.	Karate w Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	45 day waiting period for harvest
	Mustang MAXX EC [3] (zeta-cypermethrin)	2.6 to 4 fl oz (0.016 to 0.025 lb ai/A)	30 day waiting period for harvest
	Proaxis 0.5 CS [3] (gamma-cyhalothrin)	2.56 to 3.84 fl oz (0.01 to 0.015 lb ai/A)	45 day waiting period for harvest
	Tombstone [3] (cyfluthrin)	2.0 to 2.8 fl oz (0.031 to 0.044 lb ai/A)	30 day waiting period for harvest
	Warrior II w Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	45 day waiting period for harvest
Stem weevil 1/8 inch long, grayish-brown with varying white spots on wing covers. Adults emerge in mid-late June. Damage: Adults insert eggs in stalks. Larval feeding causes weakening of stalk, easily lodged heads. Threshold: Begin scouting in mid-June. Treat when counts reach 1 weevil per three plants. In areas with history of problem, treat when plants reach 8-10 leaf stage if planted before June 1.			
	Asana XL [3] (esfenvalerate)	5.8 to 9.6 fl oz (0.03 to 0.05 lb ai/A)	28 day waiting period for harvest for harvest; do not graze
	Baythroid XL [3] (beta-cyfluthrin)	1.6 to 2.4 fl oz (0.013 to 0.019 lb ai/A)	30 day waiting period for harvest; do not graze
	Besiege [28,3] (lambda-cyhalothrin + chlorantraniliprole)	6.0 to 10.0 fl oz	45 waiting period for harvest. Do not use adjuvant with application. Follow drift precautions to protect pollinators
	Delta Gold [3] (deltamethrin)	1.0 to 1.5 fl oz (0.012 to 0.018 lb ai/A)	21 day waiting period for harvest; do not graze
	Karate w Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	45 day waiting period for harvest
	Mustang MAXX EC [3] (zeta-cypermethrin)	2.6 to 4 fl oz (0.016 to 0.025 lb ai/A)	30 day waiting period for harvest, do not graze
	Proaxis 0.5 CS [3] (gamma-cyhalothrin)	2.56 to 3.84 fl oz (0.01 to 0.015 lb ai/A)	45 day waiting period for harvest
	Sevin XLR [1A] (carbaryl)	1 to 1.5 quarts (1 to 1.5 lb ai/A)	30 day wait for grazing, 60 days for harvest
	Tombstone [3] (cyfluthrin)	1.6 to 2.4 fl oz (0.025 to 0.038 lb ai/A)	30-day waiting period for harvest
Warrior II w Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	45 day waiting period for harvest	
Sunflower beetle Similar to Colorado potato beetle, light yellow with dark brown stripes and measures about 3/4 inch long. Larvae are yellow and humpbacked. Damage: Feed on foliage, chewing holes in leaves. Thresholds: Seedlings: 1 adult per plant. Larger plants: 10 to 15 larvae + 25% defoliation.			
	Asana XL [3] (esfenvalerate)	1.45 to 5.8 fl oz (0.0075 to 0.03 lb ai/A)	28 day waiting period for harvest for harvest, do not graze
	Besiege [28,3] (lambda-cyhalothrin + (chlorantraniliprole)	5.0 to 8.0 fl oz	45 waiting period for harvest. Do not use adjuvant with application. Follow drift precautions to protect pollinators
	Delta Gold [3] (deltamethrin)	1.0 to 1.5 fl oz (0.012 to 0.018 lb ai/A)	21 day waiting period for harvest; do not graze
	Karate w Zeon [3] (lambda cyhalothrin)	0.96 to 1.60 fl oz (0.015 to 0.025 lb ai/A)	45 day waiting period for harvest
	Mustang MAXX EC [3] (zeta-cypermethrin)	2.6 to 4 fl oz (0.016 to 0.025 lb ai/A)	30 day waiting period for harvest; do not graze
	Proaxis 0.5 CS [3] (gamma-cyhalothrin)	1.92 to 3.2 fl oz (0.0075 to 0.0125 lb ai/A)	45 day waiting period for harvest
	Sevin XLR [1A] (carbaryl)	1 to 1.5 quarts (1 to 1.5 lb ai/A)	30 day waiting period for harvest; do not graze
	Tombstone [3] (cyfluthrin)	0.8 to 1.6 fl oz (0.013 to 0.025 lb ai/A)	30-day waiting period for harvest
Warrior II w Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	45 day waiting period for harvest	
Sunflower (Head) moth Adult is small white moth, 3/8	Asana XL [3] (esfenvalerate)	5.8 to 9.6 fl oz (0.03 to 0.05 lb ai/A)	28 day waiting period for harvest for harvest; do not graze

<p>inch long that folds wings around body when resting. Larvae are brown/purple with longitudinal white stripes.</p> <p>Damage: Young larvae feed on pollen and florets. Older larvae burrow into head and feed on developing seed. Larvae spin webbing on surface of flower head. Damage enables head rots to develop.</p> <p>Threshold: Begin scouting when flowers first open and scout every few days. It is best to scout in evening with flashlight. Treat when moth numbers reach 1-2 moths per five plants at 20% bloom.</p>	Baythroid XL [3] (beta-cyfluthrin)	2.0 to 2.8 fl oz (0.016 to 0.022 lb ai/A)	30 day waiting period for harvest; do not graze
	Besiege [28,3] (lambda-cyhalothrin + chlorantraniliprole)	6.0 to 10.0 fl oz	45 waiting period for harvest. Do not use adjuvant with application. Follow drift precautions to protect pollinators
	Delta Gold [3] (deltamethrin)	1.0 to 1.5 fl oz (0.012 to 0.018 lb ai/A)	21 day waiting period for harvest; do not graze
	Karate w Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	45 day waiting period for harvest
	Mustang MAXX EC [3] (zeta-cypermethrin)	2.6 to 4 fl oz (0.016 to 0.025 lb ai/A)	30 day waiting period for harvest; do not graze
	Proaxis 0.5 CS [3] (gamma-cyhalothrin)	2.56 to 3.84 fl oz (0.01 to 0.015 lb ai/A)	45 day waiting period for harvest
	Sevin XLR [1A] (carbaryl)	1.5 quarts (1.5 lb ai/A)	30 day waiting period for harvest; do not graze
	Tombstone [3] (cyfluthrin)	2.0 to 2.8 fl oz (0.031 to 0.044 lb ai/A)	30-day waiting period for harvest.
	Vantacor [28] (chlorantraniliprole)	1.2 to 2.5 fl oz (0.047 to 0.098 lb ai/A)	1 day PHI
	Warrior II w Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	45 day waiting period for harvest

Pre-Harvest Intervals

Azadirachtin (neem)	0-day PHI for harvest
Bacillus thuringiensis	0-day PHI for harvest
Besiege	21-day PHI
Brigade	35-day PHI for harvest.
Carbine	7-day PHI for harvest
Delta Gold	7-day PHI for harvest
DyndaShield	Harvested seed can only be used for industrial purposes, not for edible oil.
Exirel	7-day PHI for harvest
Fortenza	Wait 30 days before applying a foliar application of cyantraniliprole or other group 28 product
Vantacor	1-day PHI for harvest

* MOA group numbers in brackets [#] following the insecticide name are used to designate the mode of action of the insecticide according to the classification system developed by the Insecticide Resistance Action Committee (IRAC) in 2011. It is intended to help in the selection of insecticides for preventative resistance management. If you make multiple applications for a specific pest during a growing season, simply select a registered insecticide with a different number for each application. To further delay resistance from developing, integrate other control methods into your pest management programs.

** The first name listed is a commercial trade name of a product. The chemical name in parentheses refers to the name of the active ingredient and is included because there are a number of registered products that contain the same active ingredient. Such products may be less expensive to purchase, so producers should compare prices.

The information given herein is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Cooperative Extension Service is implied.

The pesticide information presented in this publication was current with federal and state regulations at the time of printing. The user is responsible for determining that the intended use is consistent with the label of the product being used. Use pesticides safely. Read and follow label directions. The information given herein is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Cooperative Extension Service is implied.

Oklahoma State University, as an equal opportunity employer, complies with all applicable federal and state laws regarding non-discrimination and affirmative action. Oklahoma State University is committed to a policy of equal opportunity for all individuals and does not discriminate based on race, religion, age, sex, color, national origin, marital status, sexual orientation, gender identity/ expression, disability, or veteran status with regard to employment, educational programs and activities, and/or admissions. For more information, visit <https://eoo.okstate.edu>.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of Oklahoma Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President for Agricultural Programs and has been prepared and distributed at a cost of 20 cents per copy. October 2022. AF.