

# **COW/CALF CORNER**

## **The Newsletter**

**From the Oklahoma Cooperative Extension Service**

**July 25, 2011**

### **In this Issue:**

#### **Drought Accelerates Cattle Liquidation in Southern Plains**

Derrell S. Peel, Oklahoma State University Extension Livestock Marketing Specialist

#### **Drought Management Meetings Scheduled**

Greg Highfill, Oklahoma State University Area Extension Animal Scientist

#### **Handling Heat**

Dave Sparks D.V.M., Oklahoma State University Food Animal Quality and Health Extension Specialist

#### **Drought Information and Resources Available**

Glenn Selk, Oklahoma State University Emeritus Extension Animal Scientist

## **Drought Accelerates Cattle Liquidation in Southern Plains**

Derrell S. Peel, Oklahoma State University Extension Livestock Marketing Specialist

The expanding extreme drought in the Southern Plains is causing a significant acceleration of cattle liquidation in the region. In Oklahoma, the combined total for federally reported auctions the past two weeks has shown a 56 percent increase in feeder cattle sales and a 205 percent increase in cow and bull sales compared to the same period one year ago. The auction totals include significant numbers of double-stocked summer stockers from the Osage country that are typically marketed this time of year.

However, there are large numbers of cows and lightweight feeder cattle that are not typically marketed this time of year. Most likely we are seeing a second wave of cow liquidation made up of cows with spring born calves that are just now big enough to early wean and sell. We are receiving many anecdotal stories from auctions, both large and small of excessive numbers of feeder cattle and cows being marketed. Livestock haulers are booked and it is difficult to arrange shipping at this time.

Prices for slaughter cows, bred cows and cow-calf pairs have dropped sharply in the past two weeks. This is likely a temporary situation due, in part, to the bottlenecks of selling and shipping

so many animals in a short period of time. Producers are selling because they have no other alternatives but those with the ability to postpone sales for a couple of weeks may find the logistics as well as the price better. It is hard to say how long the current bulge in cow liquidation will last but most likely it will be a matter of no more than another 2-4 weeks.

Unfortunately, many of the cows are going to slaughter, contributing to additional cow herd liquidation. Beef cow slaughter in Federal slaughter Region 6, which overlays the drought area, is 16 percent higher for the year to date compared to last year. In the most recent two weeks of data, beef cow slaughter in Region 6 is up 35 percent compared to the same period last year. Total U.S. beef cow slaughter for the year to date is down 2.7 percent but the gap is closing due to large slaughter totals in the drought area. For most weeks of the year, the national total beef cow slaughter has been down compared to last year. However, in the last four weeks of slaughter data, the week to week totals for beef cow slaughter have exceeded year ago levels.

The drought is not only having devastating impacts for producers in the drought region, but is also impacting the beef cattle industry nationwide and will have impacts for several years. By pushing beef cow slaughter close to last year's record levels, the drought insures additional herd liquidation that deepens the hole from which the industry must start to rebuild. The July Cattle report confirmed that the July 1 beef cow herd was down 1 percent from last year but this survey value likely does not reflect the accelerated liquidation that has occurred so far in the month of July.

## **Drought Management Meetings Scheduled**

Greg Highfill, Oklahoma State University Area Extension Animal Scientist

Cattle and forage producers looking to make best use of management practices for fighting the drought should register now to attend the Oklahoma Cooperative Extension Service Northwest Oklahoma Drought Meetings taking place on August 4<sup>th</sup> and 5<sup>th</sup>. To address key issues producers are facing in the current drought situation, regional meetings are being held to provide cattle producers with production information and decision making tools.

The first meeting will take place from noon to 3 p.m. on Thursday, Aug. 4, at the High Plains Area Vo-Tech, located at 3921 34<sup>th</sup> St. in Woodward. Lunch is free to participants who pre-register. To register, contact the Woodward County Extension Office by email at [dana.bay@okstate.edu](mailto:dana.bay@okstate.edu) or by phone at 580-254-3391.

The second meeting will take place from 9:30 a.m. to noon on Friday, Aug. 5, at the Northwest Stockyards, located at 11802 W Owen K Garriott, seven miles west of Enid. Lunch is free to participants who pre-register. To register, contact the Garfield County Extension Office by email at [jeff.bedwell@okstate.edu](mailto:jeff.bedwell@okstate.edu) or by phone at 580-237-1228.

Topics to be discussed include: Agronomic Considerations of Drought; Hay Feed Value, Availability & Cost; Cow Feeding Options & Cost; Nitrate, Prussic Acid & Aflatoxin Issues in Forages; Cow Herd – Keeping & Feeding vs. Selling & Buying the cows back; and Tax

Implications of Selling Your Cow Herd. Participants will also be able to interact with speakers and ask questions.

For your convenience, Extension Educators from the Northwest District will be providing a quick nitrate test at no cost, and if needed, send forage samples in for lab testing at a cost of \$6.00/sample. The program is free to the public and with your reservation a meal will be provided to you by local sponsors.

## **Handling Heat**

Dave Sparks D.V.M., Oklahoma State University Food Animal Quality and Health Extension Specialist

The widespread use of air-conditioning has made life much better for people, but little has changed for our livestock. Summer is still a high stress time and your livestock can have serious issues when the thermometer climbs. Males are more affected by extreme heat than females; fat animals more than thin animals; and older animals more than young. Animals with any compromise of the immune or respiratory systems are at serious risk. Some advance planning and observation of your livestock can make a big difference in their comfort and in your profits.

Unlike horses and humans, cattle, sheep, goats, and pigs do not sweat, at least not in amounts sufficient to be beneficial for body cooling. They maintain their body temperature at or near a constant, normal, level by panting. This moves air across the highly vascular and moist mucous membranes of the mouth, tongue, and nasal passages, thus cooling the blood passing through these tissues much like the water in an engine is cooled as it passes through the radiator. For this to occur they need a lower environmental humidity and adequate water for evaporation on the surface of the membranes. If livestock are not able to maintain their normal body temperature they start to show signs of reproductive compromise first, followed by heat exhaustion at about 105 F, and cell breakdown and death at about 107 F.

In extremely hot weather it is normal for body temperatures to rise moderately above normal during the heat of the day and to cool off at night when environmental temperatures are less. It takes several hours, however for this to occur. Although air temperatures often decline in the late afternoon or evening the animal's body temperature may not fully recover its normal level until 2 or 3 o'clock in the morning after several hours of cooler temperatures. Because of this, taking the temperature to determine if an animal is sick is best done early in the morning to get a true indication. If you must "work" or handle livestock during hot weather, do it as early as possible in the morning and be finished before their body temperature starts to rise. The digestion of grain generates a lot of heat so in hot weather it is best not to feed high levels of grain and to feed grain early in the morning.

In hot weather the first thing to suffer in your herd is reproductive efficiency. Reproductive problems can range from poor fertility to no fertility. In some males high core body temperature causes suppression of libido, but that is only the beginning of the problems. In male mammals the testicles cannot produce or maintain sperm cells at body temperature. The scrotum is designed to keep the testicles several degrees cooler than the body's core temperature by means of special muscles that lower the testicles away from the body as air temperature rises and pull

them back closer as air temperature decreases. Also, the pampiniform plexus is a heat exchange unit that cools the blood entering the testicles. When these mechanisms are overcome by the environmental temperature problems occur. Sperm cell formation, or spermatogenesis, starts to decrease when the testicular temperature rises as little as 1/2 degree and sperm cells start to die if the testicular temperature rises as much as 2 degrees above optimum. This can be significant because if extremely hot weather causes the death or deformation of sperm in the male system it can take as long as 6 weeks for new cells to be formed and mature. This can result in a temporary sterility. Research has shown that in females, high body temperatures can result in lowered conception rates and embryonic death. Excessive heat affects embryo survival and fetal development most markedly during the first 21 to 30 days after breeding.

Charles Dudley Warner, a friend of Mark Twain, is famous for saying “Everyone complains about the weather, but no one does anything about it.” So what can we do? The simplest answer is shade and the simplest shade is the shade tree. Closed in spaces are not very helpful because they restrict air movement. If you use a barn for shade, utilize a breezeway or fans. If natural shade is not available a little creativity and simple materials can provide permanent or temporary shade. On an extremely hot day a shade structure can cause a drop of 12 to 15 degrees F in environmental temperature. The accompanying photos were taken at the same time, one in the sun and one under a temporary shade. Sprinklers and misters are often used for cooling. Wetting the ground, can help to reduce temperature by evaporative cooling and also helps to keep dust irritation down.

A good rule of thumb for drinking water is if you wouldn't drink it and enjoy it, neither will your livestock! In cool temperatures mature livestock drink about 10% of their body weight per day in water. This is about 1.25 gallon per day for each 100 lbs of body weight. In summer this doubles or even triples in order to meet body cooling requirements. Cleaning water tanks and supplying fresh water becomes even more critical. The question is not “does my animal drink”, but “does my animal drink enough?” Summertime temperatures also necessitate more frequent tank cleaning because as temperature rises, bacteria and algae grow much faster. Fencing animals out of ponds and providing drinking access only keeps the water clean. Ponds surrounded by vegetation stay significantly cooler than those surrounded by bare ground and this enhances water consumption. Young animals have even more critical requirements. They drink smaller amounts at a time, have a higher metabolic rate, and are more likely to be finicky drinkers.

Excessive heat can affect your livestock and your profits in several ways. Hot animals have poor appetite and growing animals that don't eat don't grow. Females that are trying to pick up body condition prior to breeding season or calving need to eat to capacity as well. While air conditioning is not practical, adequate shade, plenty of clean water, and a moratorium on handling in the afternoon and evening can limit health problems and help ensure good reproduction rates in your herd.

## **Drought Information and Resources Available**

Glenn Selk, Oklahoma State University Emeritus Extension Animal Scientist

Ranchers impacted by the drought and heat of 2011 now have an important resource available to them. The Oklahoma State University Extension Beef Team has put together a tremendous listing of drought management tools that can be of value to producers. The website contains maps, weather forecast information, forage management suggestions, hay listings, cattle supplementation information, other cattle management suggestions, financial services, tax considerations, on-going government policies resulting from drought, and much more. Go to [www.beefextension.com](http://www.beefextension.com) and click on “drought resources”.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, Title IX of the Education Amendments of 1972, Americans with Disabilities Act of 1990, and other federal laws and regulations, does not discriminate on the basis of race, color, national origin, sex, age, religion, disability, or status as a veteran in any of its policies, practices or procedures. This includes but is not limited to admissions, employment, financial aid, and educational services. References within this publication to any specific commercial product, process, or service by trade name, trademark, service mark, manufacturer, or otherwise does not constitute or imply endorsement by Oklahoma Cooperative Extension Service.