



# Propagating Peaches by "T"-Budding

Dean McCraw  
Extension Horticulturist  
Pecans and Tree Fruits

Oklahoma Cooperative Extension Fact Sheets  
are also available on our website at:  
<http://osufacts.okstate.edu>

The "T" or shield-bud method of propagation may be used on many kinds of plants: apples, pears, peaches, and a large number of ornamentals. The "T" bud is considered the easiest method of plant propagation.

Peaches are usually propagated during the seedling stage of growth when they are at least as large in diameter as a pencil. If this size is attained in June, they may be budded and forced into growth in a short time. When seedling development is delayed, budding is usually performed in August and early September and the buds are not forced into growth until the next spring.


Bud wood is secured from desirable varieties at the time the budding is done. Select strong, vigorous, new shoot growths. Bud wood is big enough to use when a reddish color develops on the twig and the buds show brown-colored spots.

Immediately remove the leaves from the bud stick. Use a knife or hand shears to clip off the leaves, leaving a short stub of the petiole of the leaf (on the stock) 1/4 to 3/8 inch long.


This stub can serve to protect the bud adjacent to it and also serve as a handle to hold the bud while it is being cut from the stick and being inserted into the seedling.

The immature wood at the terminal end of the shoot should be removed. Ordinarily, bud sticks should be 10 to 15 inches in length. Keep the bud sticks wrapped in moist burlap sacks or suitable material to prevent drying.


Material prepared by E.L. Whitehead, former Extension Horticulturist.



**1. PREPARING STOCK**  
A young seedling as large or larger in diameter as a pencil is preferable. In preparing the stock for budding, remove all the growth on the lower 6 to 10 inches of the tree.



**2. MAKING FIRST CUT**  
The first cut is made lengthwise of the stem near the ground line, preferably on the north side of the stock. Next, make the cross cut by a rolling movement of the knife, which lifts the corners of the bark where the two cuts cross each other.



**3. OPEN THE MATRIX**  
The same area with the matrix opened out to receive the bud.



#### 4. CUTTING THE BUD

Cut the bud from the stick by making the cut upward just underneath the bud. This cut should be from 3/4 to 1 inch in length, extending below and above the bud.



#### 5. REMOVE THE BUD

Remove the bud from the bud stick by making a cut across the top through the bark and peel the bark and bud from the stick, leaving the wood attached to the stock. The bud may be held on the blade of the knife with the thumb on the leaf stub while it is being inserted in the stock. If the bark does not slip, leave the wood attached to the bark.



#### 6. INSERT UNDER BARK

Insert the bud in the matrix under the bark and push it down until it is firmly in place. If a part of the tissue extends above the top of the "T" cut, it should be removed.




#### 7. WRAPPING THE BUD

Secure the bud to the stock by wrapping with 1/8-inch rubber budding strips, making at least two wraps below the bud and two above. The band maintains constant pressure, but expands with the growth of the tree.




#### 8. FORCING BUDS

When June budding, to force the bud, break the top over about one-half the distance from the ground to the top of the seedling, leaving it attached to shade the row of seedling trees. When "T"-budding is performed during late August or early September, the buds should callus well during the remainder of the growing season and be forced into growth the following spring.




#### 9. NEW GROWTH STARTS

Shows the new growth from the inserted bud.




#### 10. REMOVE THE TOP

When the forced bud starts to grow, remove the top of the seedling just above the new shoot.



#### 11. REMOVE NEW GROWTH

Keep the new growth removed from the base of the seedling below the bud. Some of the shoots on the new top growth may be removed near the base to force more length of growth into the new tree. In June budding, the old top should be removed as soon as the bud starts to grow. Keep the surplus growth removed in order to produce a larger tree during the growing season.



#### 12. THE TREE IS READY

The converted tree, the new variety, after a season's growth. The tree is ready for transplanting to its permanent location.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, Title IX of the Education Amendments of 1972, Americans with Disabilities Act of 1990, and other federal laws and regulations, does not discriminate on the basis of race, color, national origin, gender, age, religion, disability, or status as a veteran in any of its policies, practices, or procedures. This includes but is not limited to admissions, employment, financial aid, and educational services.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Robert E. Whitson, Director of Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources and has been prepared and distributed at a cost of 20 cents per copy. 0607