UNIVERSITY OF OKLAHOMA

GRADUATE COLLEGE

ON THE NASH PROBLEM OVER 3-FOLD TERMINAL SINGULARITIES

A DISSERTATION

SUBMITTED TO THE GRADUATE FACULTY

in partial fulfillment of the requirements for the Degree of

DOCTOR OF PHILOSOPHY

STEVEN LIN

Norman, Oklahoma

2025

Downloaded from SHAREOK. Persistent Handle URL: https://hdl.handle.net/11244/341608

ON THE NASH PROBLEM OVER 3-FOLD TERMINAL SINGULARITIES

A DISSERTATION APPROVED FOR THE

DEPARTMENT OF MATHEMATICS

BY THE COMMITTEE CONSISTING OF

Dr. Roi Docampo Álvarez, Chair

Dr. Erin Maher

Dr. Travis Mandel

Dr. Gregory Muller

Dr. Murad Özaydın

©Copyright by STEVEN LIN 2025

All Rights Reserved.

Contents

Contents iv

Abstract vi

Acknowledgments vii

1 Introduction 1

2 Preliminaries 6

2.1 Singularities of a Variety 6

2.2 Divisors of a Variety 8

3 Concepts and Tools in Birational Geometry 13

3.1 An Overview of the MMP 14

3.2 Terminal Singularities and Minimal Models 17

3.3 3-Fold Terminal Singularities 19

4 Arc Spaces and the Nash Problem 24

4.1 Arc Spaces of a Variety 24

4.2 The Nash Problem: Formulation 27

4.3 The Nash Problem: Known Results 33

4.4 The Nash Problem: Further Developments 34

5 Some Toric Geometry **37**

5.1 Definitions, Results, and Notations 37

5.2 Some Birational Geometry of Toric Varieties 39

6 Nash Valuations over Toric Terminal Singularities **41**

6.1 Discrepancy bounded by 1 induces Nash Valuations 42

6.2 Nash Valuations have Discrepancy bounded by 1 44

6.3 Main Result and Conjectures 46

7 Nash Valuations over 3-Fold Terminal Singularities of Type $\mathbf{cAx/2}$ **48**

7.1 Candidates for Nash Valuations 49

7.2 Setup for the Proof 49

7.3 Weighted Blowing-Ups 52

7.4 Proof of Theorem 7.0.1 55

8 Nash Valuations over 3-Fold Gorenstein Terminal Singularities **62**

8.1 Main Theorem 62

8.2 Examples 64

Bibliography **66**

Abstract

In this dissertation, we study the *Nash problem* over terminal singularities in dimension 3, or *3-fold terminal singularities*. In particular, for 3-fold terminal singularities, we propose two conjectures that aim to characterize their induced *Nash valuations*.

(A) Any prime divisor with discrepancy bounded by 1 induces a Nash valuation.

(B) Any prime divisor with minimal discrepancy induces a Nash valuation.

We prove that Conjecture A holds for toric terminal singularities in arbitrary dimension, and Conjecture B holds for 3-fold terminal singularities of type cAx/2. In the Gorenstein cases, we provide a partial result with examples for both of the conjectures.

Acknowledgments

This dissertation has been completed with a lot of effort, support and encouragement from many people, including myself.

First of all, I would like to express my sincere gratitude to my academic advisor, Prof. Roi Docampo Álvarez, for his guidance throughout my graduate studies at OU. I came to OU with very limited knowledge of algebraic geometry, as my original background was in differential geometry and geometric analysis. Roi devoted a great deal of time and effort to teaching me most of the fundamental materials (Hartshorne's Algebraic Geometry, for example) and patiently answering my various (sometimes subtle) questions in algebraic geometry, either in official courses or personal discussions. These included essential topics such as commutative algebra, birational geometry, jet schemes, arc spaces, and non-Archimedean analytic geometry. I have been deeply inspired by his philosophy of presenting mathematical ideas and techniques with clear reasoning and depth. This dissertation would not have been possible without countless discussions with him, his careful and rigorous check of the details in my proofs, and many ideas and potential research directions for the research topics he generously provided.

I am thankful to my graduate committee members Prof. Gregory Muller, Prof. Travis Mandel, Prof. Murad Özaydın, Prof. Jonathan Kujawa (Oregon State U.), and Prof. Erin Maher for spending time serving on my graduate committee, and

providing feedback and suggestions during my general exam and dissertation defense, helping me improve my dissertation's quality and make the whole Ph.D. program process run smoothly.

Besides Roi's courses, there are many professors in the math department at OU who offer nice courses or lectures, and these courses have influenced my point of view in mathematics, including the research and teaching perspective. Prof. Gregory Muller provided most of the graduate level courses related to algebraic geometry, including Lie theory, introduction to moduli spaces, quiver representations, algebraic geometry, and homological algebra. Prof. Travis Mandel gives us a very accessible and moderate toric algebraic geometry course. As an extension of this course, Travis organizes a small reading seminar on Fulton's intersection theory and Gromov-Witten invariants, and I also learn a lot from these events. Another amazing course is the representation theory delivered by Prof. Janathan Kujawa. His course helped me grasp most of the fundamental concepts that can help me understand the context of representation theory. Prof. Murad Özaydın has provided moderate-paced and warm courses in algebraic topology , non-commutative geometry, and category theory. In particular, algebraic topology is the first non-qualify exam course that I took after joining the math department at OU. I also enjoy the short but intense D-modules topics course taught by Prof. András Cristian Lőrincz. Prof. Miroslav Kramar offered a topological data analysis (TDA) course, which provided me a good overview for not only the theoretical but also the programming perspective of this topic.

Additionally, I want to thank Prof. Shihshu Walter Wei for sharing many interesting intuitions and ideas about differential geometry and geometric analysis. Especially I learned a lot of interesting properties and philosophy about harmonic maps. Also, I thank Prof. Meijun Zhu for his hospitality and suggestions when I

was in my first year of the Ph.D. program.

There are many people in the math department at OU who have made my Ph.D. journey more enjoyable and memorable. I would like to thank Chih-Kuang Lee, Shravan Saoji, and Prof. Mario Morán Cañón (U. Autónoma de Madrid) from Roi's research team. It is nice that we have many interesting discussions and conversations about math and non-math. Especially, Chih-Kuang and I have known each other for over 15 years as friends studying mathematics from undergraduate and master programs at National Taiwan University, and we came to OU to study the same Ph.D. program. Our friendship began with a shared academic interest, which is a connection that I deeply cherish.

We successfully held *Student Algebra Symposium* (SAS) during the 2023 fall. As an organizer, I would like to express my gratitude to the other organizers Ergın Süer, Ahmet Uzun, and Arman Yagci for their efforts to organize this wonderful event. I also thank the Department of Mathematics at OU (special thanks to Prof. Nikola Petrov, Mike Donlon, and Cristin Sloan) for providing financial support to the participants. Especially, Ergın Süer, Chih-Kuang Lee, and I joined OU in the same year, and we have shared many engaging conversations and interactions about mathematics and various other topics throughout our time in the program. It has been a true pleasure to meet you at OU, and I hope to visit you in Istanbul someday after we graduate.

I also want to mention Chary Addanki, James Beyer, Jeremiah Isu, Jay Lane, Siddharth Ramakrishnan, Martin Carlson, Arghadeep Roy, Sayantan Santra, Lucas Smith, Tomoya Tatsuno, Bach Tran, Noah Torgerson, Maya Verma, Jeremy West, James White, Samuel Lin, Anuradha Ekanayake, José Lozano, Yunchuan Zhou, Oshadi Samarathunge, Javier Loya Chavira, Sajal Halder, Pratit Goswami, Emmanuel Elima, Henry Carnick, Emelia Boye, Lucas Yong, Tatiana Abdelnaim, Carlos Vil-

lanueva, Dustin Gaskins, Ryan Reynolds, Elizabeth Pacheco, Wenwen Li, Haley Bahn, James Ramsey, Nathan Jones, Patrick Dynes, Erin Hausmann, Mingshi Yao, Joseph Randich, Andrew Lutz, Yilin Jiang, Ling Jin, Connor Davis, Ben Johnson, Fahed Mokdad Reyes, and Shaoyun Yi for their academic and non-academic interactions when studying at OU. It is really great to have fun with them.

For my teaching experience at OU, I would like to express my gratitude to Ashley Berger, Carye Chapman, Ashley Taylor, Rachel Wright, Prof. Deborah Moore-Russo, Prof. Peter Patzt, Prof. Justin Malestein, and Prof. Tomasz Przebinda for their guidance and coordination in the courses I taught as a Graduate Teaching Assistant (GTA). Their support helped me navigate various teaching responsibilities smoothly and significantly contributed to the development of my teaching skills. These experiences have laid a solid foundation for my future work in education.

Besides the Department of Mathematics at OU, I also express my sincere gratitude to Prof. Dimitris Diochnos from the School of Computer Science at OU. In his course, he introduced me to the area of computational learning theory, specifically Probably Approximately Correct (PAC) learning, which greatly broadened my academic horizons. I also thank Prof. Xuguang Wang from the School of Meteorology at OU for offering a great course on advanced data assimilation. Her course sparked my interest in the mathematical techniques used in weather forecasting, particularly the ensemble Kalman filter (EnKF).

In addition to my academic life, the Taiwanese community in Norman has also brought me great joy and support during my time at OU. I am especially grateful to the members of the OU Taiwanese Student Association for organizing numerous events and preparing delicious meals. Their warmth and hospitality created a strong sense of belonging that truly enriched my experience in Oklahoma.

Apart from the life in the U.S.A., many connections among my friends and

professors in Taiwan before coming to OU are still ongoing.

Mathematically, I would like to thank Prof. Jung-Kai Alfred Chen (NTU) for inviting him to give a talk about my current research in algebraic geometry seminar of the National Center for Theoretical Science (NCTS), which is the institute where I worked before I came to OU. He discusses some of the related topics and potential directions with me. I also thank Prof. Hsueh-Yung Lin (NTU) for providing me with some questions and ideas during the seminar for my research. Furthermore, my academic advisor of the master program Prof. Shu-Cheng Chang (NTU), still continues to make contacts to me with encouragements, hospitality, a lot of Cauchy-Riemann geometry, and Ricci flows. It is quite nice to keep in touch with them even though I have graduated from the National Taiwan University for several years. I also thank Prof. Chuan-Tsung Chan (Tunghai U.) from the Applied Physics Department for inviting me to give seminar talks/discussions about jet shcemes/arc spaces at Tunghai University, and have various conversations between mathematics and theoretical physics.

I would like to thank Kui-Yo Chen (NCKU), Prof. Tsung-Ju Lee (NCKU), and Wei-Chung Chen (U.Tokyo) for their useful conversations about combinatorics and birational geometry, which directly provides reputations to my work. Many friends in the Taiwanese mathematical community still provide me with inspiration. I would like to thank Zhong-Tang Wu (Tunghai U.) and Ting-Hung Yu (UI) for providing me with further potential directions and ideas, including math, physics, statistics, and life. Also, I would like to thank Tzu-Mo Kuo (NCTS), Chun-Wei Lee (NTU), Yang-Zhi Lin, Yi-Ting Liu, Hsin-I Guo, I-Shing Hu (NTNU), Kuo-Hsin Wang, Chin-Yi Lin, Sergei Hung-Yu Hsiao (Tamkang U.), Jin-Han Liou (NTU), Chia-Jung Hsu (Chalmers U. of Technology), Yu-Tang Wu (NTU), Prof. Nobuo Sato (NTU), Prof. Hsin-Lun Li (NSYSU), and Prof. Chih-Wei Chen (NSYSU) for

their many math or non-math chatting, assistance, suggestions, and fun.

I have also benefited greatly from attending various conferences and summer schools, which have not only inspired me academically but also provided valuable opportunities to connect with fellow mathematicians.

First, I want to thank Simons Laufer Mathematical Sciences Institute (SLMath, or MSRI) for providing me with a nice opportunity to attend summer school *Multigraded and Differential Graded Methods in Commutative Algebra* held at St. Mary College, California, in 2024. I learn a lot from the instructors and origanizers, Prof. Michael Brown (Auburn U.) and Prof. Claudia Miller (Syracuse U.), and other participants (Shravan Saoji was also in this school). It is a wonderful summer to learn math with you all and visit the SLMath Center at the University of California, Berkeley and the city center of San Francisco.

I would like to thank Le Centre Henri Lebesgue for providing me with the opportunity to present my results at *Spring School/Conference: Singularities in Algebraic Geometry*, held in 2025 at the Université de Rennes, France. I am especially grateful to the organizers - Prof. David Bourqui, Prof. Goulwen Fichou, and Prof. Christophe Mourougane (all from Université de Rennes), and other participants around the world - for their kind support and excellent organization. This event significantly broadened my perspective on singularity theory, particularly through the lens of jet schemes and arc spaces. I also thank all the lecturers at the spring school: Prof. Lorenzo Fantini (École Polytechnique), Prof. Shihoko Ishii (U.Tokyo/Tokyo Institute of Technology), Prof. Hussein Mourtada (U. Paris-Cité), Prof. Tamara Servi (U. Paris-Cité), Prof. Enrica Floris (U. de Poitiers), Prof. Anne Moreau (U. Paris-Saclay), and Prof. Mircea Mustață (U. of Michigan) for offering many nice courses and providing me many ideas in my research. I would also like to express my appreciation once again to Prof. Mario Morán Cañón (U. Autónoma

de Madrid) for introducing me to Rennes, a beautiful french city with its charming architecture, furnished histories, cultures, inviting bookstores, shops, restaurants, and delicious galettes, crêpes, and kouign-amanns.

My dear family in Taiwan has been a constant source of support throughout my graduate studies. I would like to thank my father, sister, aunts, uncles, cousins, grandfather, and other relatives for their warm wishes, love, encouragement, and unwavering support during my Ph.D. journey in mathematics. Whenever I felt exhausted and decided to return to Tainan for a break, they always welcomed me with open arms and helped me recharge mentally, giving me the strength to return to Norman and continue my work. I am especially grateful to those who traveled all the way from Taiwan to attend my graduation ceremony-a 14-hour flight-for which I am deeply touched and grateful. Besides, I appreciate Chih-Kuang Lee's family, including his father, mother, and brother, for their hospitality when I visit the city of Taipei, including introducing the Luzhou District, the place where Chih-Kaung grew up.

In addition, I am deeply grateful to my dear cousin Josephine Min-Tsen Tseng and her husband Shane Jui-Pin Tseng in Houston, who have provided me with invaluable information and support on how to adapt to life in the United States, especially in Texas and Oklahoma. Their help was especially well supported when I first arrived in Norman. I will always cherish the wonderful and memorable times I visited them in Houston, along with their lovely children and dogs.

To my beloved mother: I am deeply saddened and sorry that you are not here to witness and celebrate the moment I complete my Ph.D. degree, which was one of the dreams that you wanted to realize before you passed away. Your love, encouragement and strength have always been with me, and I will carry them with me to start my new journey after I graduate. Although we often have many misun-

derstandings about each other in all aspects, I am very sure that you will be proud of my achievement at OU. Don't worry, I promise that I will try my best to take good care of myself, father, and sister.

Here I would like to express my special thanks to Shu-Yu Lin, who consistently offered warm encouragement, thoughtful suggestions, and kind words as I approached the completion of my dissertation. It has been a great pleasure to know her since my master's studies. I look forward to sharing new moments and experiences together after completing my Ph.D. degree.

Finally, I would like to express my respect and gratitude to Martin Gardner (1914–2010). His book *Aha! Insight* offered me a completely new perspective on mathematics when I was a junior high school student-a time when I actually disliked the subject. That book became the starting point of my journey into mathematics from my junior high school period. Even today, it remains a beacon that guides me whenever I feel stuck in my research or studying.

There are many others who have helped me along the way, but are not mentioned here by name. I am deeply grateful for their support and kindness.

This may mark the end of my graduate studies in mathematics, but it is also the beginning of my journey as a researcher. Even if I eventually choose to leave academia, this invaluable period has undoubtedly shaped my life and personal philosophy. I would also like to thank myself for enduring and growing through this challenging but meaningful time.

I would like to conclude with my favorite sentence that appears in the novel *Kino no Tabi-the Beautiful World*, written by Keiichi Sigsawa.

The world is not beautiful. Therefore, it is.

Keng-Hung Steven Lin, in Fort. Collins, Colorado

Chapter 1

Introduction

The aim of this dissertation is to study Nash valuations over 3-fold terminal singularities.

Singularities play a crucial role in the study of algebraic varieties [Ish18], and they often provide useful characterizations of the varieties under consideration.

In birational geometry, singularities can tell us some important information in birational classifications of algebraic varieties. By applying certain birational maps, one can study how singularities behave and transform. A natural and fundamental tool in this context is the resolution of singularities.

Definition 1.0.1 (Resolution of Singularities)**.** Let X be an algebraic variety over a field k. A *resolution of singularities* over X, or a *resolution of* X for short, is a proper birational morphism $f : Y \to X$ such that Y is non-singular.

If the ground field k is of characteristic 0, the existence is confirmed in [Hir64] by H. Hironaka,

Theorem 1.0.2 (Existence of Resolution of Singularities, [Hir64])**.** *Let X be a variety over a field k of characteristic 0, then X has a resolution of singularities.*

For a resolution $f : Y \to X$, the preimage of the singular locus $f^{-1}(X_{\text{sing}})$ is called the *exceptional locus* of f. Since an exceptional locus arises from resolving singularities, it is expected to encode some information about the singularities we study. In particular, we are interested in the *exceptional divisors* with the associated *valuations* induced from the exceptional locus of a resolution. However, resolutions of singularities are generally not unique, so it is natural to seek a birational invariant that connects exceptional loci among all resolutions. This induces the concept of *essential exceptional divisors*.

Definition 1.0.3 (Essential Exceptional Divisors)**.** Let X be a variety over k. An exceptional prime divisor is *essential* over X if for any resolution of singularities $f : Y \to X$, its center is an irreducible component of $f^{-1}(X_{\text{sing}}) \subseteq Y$

The valuation associated with an essential divisor over X is called an *essential valuation* of X. Such valuations can be defined independently from the choice of resolutions, making them natural candidates for birational invariants associated with the given singularities.

However, it is technically difficult to check whether a given valuation of X is essential over X or not, so we need some intrinsic ways to detect it. In [Nas95], J.F. Nash proposed that we can use the irreducible families of arcs based on the singularities in X to characterize essential valuations. In other words, the correspondence

$$\mathcal{N}_X : \{\text{Nash valuations of } X\} \to \{\text{essential valuations of } X\}$$

is a well-defined map and an injection.

The map $\mathcal{N}_X$ is called the *Nash map* of X. It is natural to ask if a Nash map is surjective, i.e., can any essential valuation of X be obtained from a valuation induced from some irreducible family of arcs based at X_{sing}?

Because of this, we are interested in characterizing the images of Nash maps. In [dFD16], de Fernex and Docampo use the *relatively minimal model program (rMMP)* over X to characterize part of the image of a Nash map. More precisely, they introduce the concept of *terminal valuations* of X and prove that they are Nash valautions of X.

$$\{\text{terminal valuations of } X\} \subseteq \{\text{Nash valuations of } X\}.$$

However, this characterization becomes useless when X has only terminal singularities. In particular, there will be no terminal valuations of X when X has only terminal singularities, so we need other characterizations for these cases.

In this dissertation, we try to use *discrepancies* to characterize Nash valuations of X when X has only terminal singularities. We obtain the following two main theorems.

Theorem 1.0.4. *Let $X(\Delta)$ be a $\mathbb{Q}$-Gorenstein toric variety associated with a fan $\Delta \subseteq N_{\mathbb{R}}$ of dimension at least 3 having only terminal singularities, then any exceptional prime divisor E over $X(\Delta)$ with discrepancy $a(E, X(\Delta)) \leq 1$ induces a Nash valuation of $X(\Delta)$.*

Motivated by this result and the ones of [IK03, Che23], we are led to propose the following conjectures.

Conjecture A. *Let (X, p) be a germ of 3-fold terminal singularity with the Gorenstein index $m \geq 1$, then every exceptional prime divisor E with $a(E, X) \leq 1$ induces a Nash valuation of X.*

Conjecture B. *Let (X, p) be a germ of 3-fold terminal singularity with the Gorenstein index $m \geq 1$, then every exceptional prime divisor E with minimal discrepancy (we have $a(E, X) = \frac{1}{m}$ by [Mar96, Kaw93]) induces a Nash valuation of X.*

We prove that Conjecture B is true for a 3-fold terminal singularity of type cAx/2.

Theorem 1.0.5. *Let (X, p) be a germ of 3-fold terminal singularity of type cAx/2, then any exceptional prime divisor E over X with minimal discrepancy ($a(E, X) = \frac{1}{2}$ in this case) induces a Nash valuation of X.*

In other words, our results show that Conjecture A holds for toric terminal singularities in any dimension at least 3, and Conjecture B holds for 3-fold terminal singularities of type cAx/2.

The structure of this dissertation is organized as follows.

Chapter 2 reviews some fundamental notions and tools in algebraic geometry, including singularities, resolutions, and divisors (especially the relationships between Weil divisors and Cartier divisors with the concept of a divisor is $\mathbb{Q}$-Cartier), and canonical divisors.

Chapter 3 offers a brief introduction to birational geometry, providing the background for the development of our theory. We begin with a short overview of the minimal model program (MMP), then define terminal singularities and minimal models. In particular, our focus is on the 3-fold cases. To study these singularities explicitly, we introduce their classifications along with the classical tool used for their characterization: divisorial contraction with minimal discrepancy.

Chapter 4 introduces the main objects of this dissertation: arc spaces and the Nash problem. We go through some concepts of the arc space of a variety, including its definition and examples. Then the Nash problem is formulated in the language of valuations, and we review some known results and progress that motivate our work.

Chapter 5 briefly recalls the tools from toric algebraic geometry that are essential for the development of the next chapter. This chapter does not provide any detailed proof or explanation.

Chapters 6 through 8 present our results on the Nash problem over 3-fold terminal singularities. In particular, Chapter 6 focuses on Nash valuations over toric terminal singularities of dimension at least 3, and introduces the two conjectures motivated by the observation in this chapter. Chapter 7 deals with Nash valuations over 3-fold terminal singularities of type $\mathrm{cAx}/2$. Finally, Chapter 8 provides a partial result on the Nash problem over 3-fold Gorenstein terminal singularities with some examples.

Chapter 2

Preliminaries

In this chapter, we recall and introduce some fundamental definitions, concepts, notation, and results in algebraic geometry. More details about them can be found in [Har77, Laz04a, Laz04b, Cut18].

In this dissertation, a *variety* X means an integral (reduced and irreducible) and separated scheme of finite type over $\mathbb{C}$. We put the Zariski topology on X.

Remark 2.0.1. The ground field $\mathbb{C}$ can be replaced by any field k that is algebraically closed and of characteristic 0 (for example, let $k = \bar{\mathbb{Q}}$, the field of algebraic numbers).

2.1 Singularities of a Variety

For a variety X, we can ask what is the algebraic condition for characterizing if it can be parametrized by local coordinates (or the condition that X can be a manifold). The following definition tells us that this can be described by the localization of its structure sheaf $\mathcal{O}_X$.

Definition 2.1.1. Let X be a variety. A point $p \in X$ is called a *non-singular point* (or a *regular point*) if the localization of the structure sheaf $\mathcal{O}_{X,p}$ is a regular local ring. Otherwise, p is called a *singularity* of X.

X is called *non-singular* (or *regular*) if all points on X are non-singular.

Remark 2.1.2. Notice that for the varieties in our setting, smooth points and non-singular points are the same concepts (see [Har77, Chapter III, Corollary 10.7]). We will use the terminology *smooth points* to refer to non-singular points, and *smooth variety* for non-singular variety.

Remark 2.1.3. If $X = V(f_1, \ldots, f_m) \subseteq \mathbb{C}^n$ is an affine variety, then the definition 2.1.1 for $p \in X$ being smooth is equivalent to the Jacobian matrix

$$\begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \cdots & \frac{\partial f_1}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1} & \cdots & \frac{\partial f_m}{\partial x_n} \end{bmatrix}$$

has rank equal to $\operatorname{codim}(X, \mathbb{C}^n)$ at p. This definition agrees with the notion that X is a manifold at p in differential geometry, which follows from the implicit function theorem (see [Har77, p. 32]). In particular, when X is a complete intersection, this condition is equivalent to its Jacobian matrix being of full rank at p.

The set of all singular points on X is denoted by X_{sing}, and it is called the *singular locus* of X. It is a closed set in X (see [Har77, Theorem 5.3, p. 33]). Its complement $X_{\text{sm}} := X \setminus X_{\text{sing}}$ is called the *smooth locus* of X.

We begin by recalling the notion of *birational equivalence* between two varieties. Roughly speaking, two varieties are birational if they are isomorphic on dense open subsets.

Definition 2.1.4 (Birational Varieties/Birational Morphisms)**.** Let X, Y be varieties. We say X is *birational* to Y if there are Zariski open sets U and V, with $U \subseteq X$ and

$V \subseteq Y$, such that U is isomorphic to V. Furthermore, a *birational morphism* $f : X \to Y$ is a morphism such that the restriction $f|_U : U \to V$ is an isomorphism for some Zariski open sets U and V, with $U \subseteq X$ and $V \subseteq Y$.

In birational geometry, it is interesting to understand the behaviors of the singular loci of varieties under birational morphisms. Among such morphisms, one of the most intuitive and important candidates is *resolution of singularities*.

Definition 2.1.5 (Resolution of Singularities)**.** Let X be a variety. A *resolution of singularities* of X, or a *resolution of* X for short, is a proper birational morphism $f : Y \to X$ such that Y is smooth. A resolution $f : Y \to X$ is called a *Hironaka resolution* if, furthermore, f induces an isomorphism between $Y \setminus f^{-1}(X_{\text{sing}})$ and $X \setminus X_{\text{sing}}$. A Hironaka resolution $f : Y \to X$ is called *good* if $f : Y \to X$ can be obtained by finitely many blowing-ups along smooth centers.

The result of existence for characteristic 0 is proved by H. Hironaka.

Theorem 2.1.6 (Existence of Resolution of Singularities [Hir64])**.** *Good Hironaka resolutions of singularities exist for varieties over any field of characteristic 0. In particular, resolutions of singularities exist.*

2.2 Divisors of a Variety

A fundamental tool for the algebraic study of varieties is the concept of *divisors*, which encodes some geometric information of a given variety.

Definition 2.2.1. Let X be a variety. X is *regular in codimension one* if every local ring $\mathcal{O}_{X,p}$ with $\dim(\mathcal{O}_{X,p}) = 1$ is regular.

Example 2.2.2.

- Smooth varieties are regular in codimension one.
- Normal varieties are regular in codimension one.

Assumption 2.2.3. We assume that our varieties are normal (for a variety X, every local ring $\mathcal{O}_{X,p}$ for every $p \in X$ is integrally closed in the field of fractions) in the rest of this dissertation. In particular, they are regular in codimension one.

Definition 2.2.4 (Divisors/$\mathbb{Q}$-Divisors)**.** Let X be a variety. A *(Weil) prime divisor* E is a subvariety of X with codimension $\operatorname{codim}(E, X) = 1$. A *(Weil) divisor* D is a finite formal sum $D = \sum_{i \in I} a_i E_i$ where the E_i are prime divisors and $a_i \in \mathbb{Z}$. We denote the free abelian group generated by prime divisors of X by $\operatorname{WDiv}(X)$. That is,

$$\operatorname{WDiv}(X) := \bigoplus_{E\,:\,\text{prime divisor of } X} \mathbb{Z}E.$$

A $\mathbb{Q}$-*divisor* is an element $D \in \operatorname{WDiv}(X) \otimes_{\mathbb{Z}} \mathbb{Q}$. An *effective* divisor (or $\mathbb{Q}$-divisor) is a divisor $D = \sum_{i=1}^{m} a_i E_i$ with all $a_i \geq 0$.

Let E be a prime divisor on X, and let η be the generic point of E. Then the local ring $\mathcal{O}_{X,\eta}$ is a discrete valuation ring, which induces a discrete valuation on the function field $\mathbb{C}(X)$. We call it the *valuation of* E, and denote it by val_E.

Remark 2.2.5. Notice that for a variety X, E is uniquely determined by its valuation since X is separated (see [Har77, Exercise 4.5, p. 106]).

Definition 2.2.6 (Divisors over X)**.** Let X be a variety. A prime divisor E is called a *divisor over* X if there is a proper birational morphism $f : Y \to X$ such that $E \subseteq Y$ is a prime divisor. Such E is called an *exceptional divisor over* X if f is not an isomorphism at the generic point of E. The Zariski closure of $f(E) \subseteq X$ is called *the center of* E in X, and it is denoted by c_X(E).

Definition 2.2.7 (Principal Divisors)**.** Let X be a variety, and let $f \in \mathbb{C}(X)^\times$ be a unit element in the function field of X. The *principal divisor* induced by f in X is defined to be

$$\mathrm{div}_X(f) := \sum \mathrm{val}_E(f)E,$$

where E runs over all prime divisors of X.

Remark 2.2.8. Notice that we have $\mathrm{val}_E(f) = 0$ for all but finitely many prime divisors in X (see [Har77, Lemma 6.1, p. 131]). Therefore, we have $\mathrm{div}_X(f) \in \mathrm{WDiv}(X)$.

It is common to use divisors on X to describe the geometry of X.

By comparing to the usual (Weil) divisors, there is another type of divisors that comes from invertible sheaves, which is called *Cartier divisors*.

Definition 2.2.9 (Cartier Divisors)**.** Let X be a variety. A *Cartier divisor* D consists of the data $\{(U_i, \phi_i)\}_{i\in I}$, where $\{U_i\}_{i\in I}$ is an open cover of X and $\phi_i \in \Gamma(U_i, \mathcal{K}_X^\times)$ which satisfies $\frac{\phi_i}{\phi_j} \in \Gamma(U_i \cap U_j, \mathcal{O}_X^\times)$ for all $i, j \in I$. Let $\{(U_i, \phi_i)\}_{i\in I}$ and $\{(V_j, \psi_j)\}_{j\in J}$ be two Cartier divisors. We define their sum as

$$[\{(U_i, \phi_i)\}_{i\in I}] + [\{(V_j, \psi_j)\}_{j\in J}] := \{(U_i \cap V_j, \phi_i\psi_j)\}_{(i,j)\in I\times J}.$$

The family of Cartier divisors of X with this sum forms an abelian group, and we denote it by $\mathrm{CDiv(X)}$.

Now we relate the Cartier divisors to the (Weil) divisors in the following way. Let $\Phi = \{(U_i, \phi_i)\}_{i\in I}$ be a Cartier divisor. Let E be a prime divisor in X, and we define

$$\mathrm{val}_E(\Phi) := \mathrm{val}_E(\phi_i)$$

for any $i \in I$ such that $c_X(E) \cap U_i \neq \varnothing$ (notice that $\mathrm{val}_E(\phi_i)$ is independent of U_i). Consider the following correspondence

$$\begin{aligned} \mathrm{cyc} : \mathrm{CDiv}(X) &\longrightarrow \mathrm{WDiv}(X) \\ \Phi = \{(U_i, \phi_i)\}_{i\in I} &\mapsto \sum_{E\text{: prime divisor of X}} \mathrm{val}_E(\Phi)E. \end{aligned}$$

This correspondence is injective because X is normal (see [Ful98, Example 2.1.1]). Therefore, we can identify a Cartier divisor of X with the corresponding (Weil) divisor of X via cyc.

Definition 2.2.10. A (Weil) $\mathbb{Q}$-divisor D is $\mathbb{Q}$-*Cartier* if rD is Cartier for some $r \in \mathbb{Z}_{>0}$.

Remark 2.2.11. For a smooth variety, divisors and Cartier divisors coincide. For a singular variety, a divisor is not necessarily Cartier. See [Ful98, Example 2.1.1].

Recall that we can define the intersection product $D.C$ for a Cartier divisor D and a curve C of X (see [Har77, Ful98]). For a $\mathbb{Q}$-Cariter $\mathbb{Q}$-divisor D and a curve C of X, we define $D.C := \frac{1}{m}(mD.C)$ where $m \in \mathbb{Z}_{>0}$ makes mD be Cartier.

Definition 2.2.12 (Nef Divisors)**.** Let D be a $\mathbb{Q}$-Cartier $\mathbb{Q}$-divisor. D is *nef* if $D.C \geq 0$ for all curves $C \subseteq X$.

Among all divisor of X, we usually use a *canonical divisor* to characterize geometric properties of a variety in birational geometry.

Let X be a smooth variety, and let $\Omega_{X/k}$ be the sheaf of differential on X. Consider its canonical sheaf (the sheaf of top forms) $\omega_X := \bigwedge^{\dim X} \Omega_{X/k}$, which is an invertible sheaf (line bundle) on X. Hence, $\omega_X \cong \mathcal{O}_X(K_X)$ for some Cartier divisor K_X. This K_X is not unique, but any two such divisors are linearly equivalent (i.e., their difference is a principal divisor).

Definition 2.2.13 (Canonical Divisor)**.** Such K_X is called a *canonical divisor* of X.

Remark 2.2.14. For a normal variety, we can still have a concept of canonical divisors, which is generalized from the smooth one. We will introduce it in the next chapter. Furthermore, we will use canonical divisors to define *terminal singularities*.

Chapter 3

Concepts and Tools in Birational Geometry

In this section, we recall some important results and tools in birational geometry which will be used throughout our theorems and arguments. More precisely, we will introduce the concepts of *terminal singularities* and the tool used to study them in 3-dimensional cases, which is *divisorial contraction with minimal discrepancy*.

Most of the results that we discuss are treated in detail in [KM98, Mat02, Laz04a, Laz04b, Hay99, Hay00]. We work over the field of complex numbers $\mathbb{C}$, unless indicated otherwise.

In mathematics, a common approach to understanding a certain geometry is to classify the geometric objects which we care about via some suitable equivalence. For example, in differential geometry, manifolds are studied through classifications via homeomorphisms, diffeomorphisms, isometries, conformal maps, homotopies, and other morphisms, depending on the context.

In algebraic geometry, we are interested in studying algebraic varieties up to *birational equivalence* rather than bi-regular isomorphism, and it forms the topic

called *birational geometry*. The principal goal of birational geometry is to birationally classify algebraic varieties with suitable representatives, and to study their associated birational invariants.

The initial step of birational classification is to find suitable representatives in a given birational equivalence class, and then study their properties. Recently, people have tried developing a certain algorithmic procedure to find such representatives by treating algebraic varieties as input, and this leads to the formulation of the so-called *minimal model program (MMP)*.

Generally speaking, MMP aims to produce a suitable representative for a fixed birational equivalence class as output by iteratively simplifying geometric structures within each fixed birational class.

3.1 An Overview of the MMP

MMP for Algebraic Curves (1-Dimensional Cases)

Let $[C]$ be the birational equivalence of algebraic curves represented by some curve C. Notice that there are many birational algebraic curves, either singular or smooth, in $[C]$. However, we can consider the normalization $\tilde{C} \to C$. Notice that $\tilde{C}$ is smooth and unique in $[C]$, so we can take $\tilde{C}$ as our suitable representative for $[C]$ (see [Ful89, Section 7.5] for details).

MMP for Algebraic Surfaces (2-Dimensional Cases)

For algebraic surfaces, the scenario becomes more interesting. Let $[S]$ be a birational equivalence of surfaces represented by some surface S. Similarly to the ap-

proach for curves, we can take a resolution of singularities of S, say $\tilde{S} \to S$, and $\tilde{S}$ is still in the same class $[S]$ since resolutions are birational morphisms.

Potentially, we could select $\tilde{S}$ as our suitable representative for $[S]$. However, resolutions of singularities are generally not unique for a fixed surface S, which means that there are many choices for $\tilde{S}$. In fact, given a smooth surface T we can apply a *blowing-up* along a point on T and obtain a birational morphism $T_1 \to T$, then T_1 is still smooth. Therefore, we can obtain infinitely many smooth models from resolutions of S by repeating this process, which are far from being our candidates for a suitable representative of $[S]$.

Nevertheless, we are still able to select one as our suitable representative among these smooth models. Notice that we can apply blowing-ups at points on a smooth surface to produce another smooth one, so it is natural to consider something that performs an inverse operation of blowing-ups, or called a *blowing-down*.

A blowing-down exists due to *Castelnuovo's Contractibility Criterion*.

Theorem 3.1.1 (Castelnuovo's Contractibility Criterion)**.** *Let S be a smooth projective surface, and let $C \subset S$ be a (-1)-curve on S (that is, an algebraic curve isomorphic to $\mathbb{P}^1$ with self intersection $C.C = -1$). Then there is a morphism $\pi_C : S \to S_1$ for some smooth projective surface S_1 such that*

- *$\pi_C(C) = p \in S_1$ is a point on S_1,*
- *The restriction $\pi_C|_{(S \setminus C)} : S \setminus C \to S_1 \setminus \{p\}$ is an isomorphism.*

Here we have an algorithm: Take a resolution $\tilde{S}$ of S, then there could be some (-1)-curves on $\tilde{S}$, so we can start applying a blowing-down $\tilde{S} \to S_1$ to contract them to points and obtain a "smaller" smooth surface with fewer (-1)-curves. We continue this process to obtain a sequence of morphisms $\tilde{S} \to S_1 \to S_2 \to \dots$. After several blowing-downs, this process terminates, and we can obtain a smooth sur-

face that is either a *minimal surface* or a *ruled surface* as an output. They will be the suitable representative we want for a given birational equivalence (see [Mat02]).

MMP for 3-Folds

For 3-folds, the situation becomes more complicated, but the directions are similar to the surface cases. Given a birational equivalence class $[X]$ for 3-folds, we apply a resolution of singularities of X, say $\pi : \tilde{X} \to X$. Then like the case for surfaces, we apply the *cone theorem* and the *contraction theorem* to obtain a *divisorial contraction* of $\tilde{X}$ to eliminate K_X-negative curves (i.e., the curves with $K_X.C < 0$) on $\tilde{X}$ and get a new 3-fold with fewer K_X-negative curves. In some steps, we need to apply *flipping contractions* if we obtain a small contraction, then we do *flips* to obtain a new variety to continue doing contractions. However, the new 3-fold after the contraction might no longer be smooth, and the singularities on it are called *terminal singularities*.

By adding terminal singularities to the MMP for 3-folds, the whole algorithm still works as the surface cases and produces an output 3-fold which is either a *minimal model* or a *Mori fiber space*.

Unlike surface case, we have many minimal models in a fixed birational equivalence class. However, it has been proved that these minimal models are connected by finite numbers of *flops* in the same birational equivalence class. On the other hand, Mori fiber spaces are also classified in a fixed birational equivalence class. The whole MMP works for 3-folds (see [KM98, Mat02] for details).

3.2 Terminal Singularities and Minimal Models

In this section, we introduce some tools and concepts for developing our theory. In particular, we will rigorously define terminal singularities and minimal models.

First, let us define the concept of canonical divisors for singular varieties (normal varieties, more precisely). Let X be a normal variety, not necessarily smooth. Consider its smooth locus X_{sm}, and we can define a canonical divisor $K_{X_{\mathrm{sm}}}$ of X_{sm} as in the previous chapter. Recall that such $K_{X_{\mathrm{sm}}}$ is not unique and that any two of them are linearly equivalent.

Definition 3.2.1 (Canonical Divisors for Normal Varieties)**.** Let X be a normal variety. A *canonical divisor* of X is defined to be the Zariski closure of $K_{X_{\mathrm{sm}}}$. More precisely, if $K_{X_{\mathrm{sm}}} = \sum_{i\in I} b_i E_i^\circ$ where the E_i° are prime divisors in X_{sm}, then $K_X := \sum_{i\in I} b_i E_i$ where each E_i is the Zariski closure of E_i°.

Remark 3.2.2. Like in the smooth case, the K_X defined here is also not unique, and any two of them are linearly equivalent.

Remark 3.2.3. It immediately follows from the definition that if $f\colon Y \to X$ is a proper birational map between normal varieties, and K_Y is a canonical divisor on Y, then $f_*(K_Y)$ is a canonical divisor on X.

Now we can use resolutions to compare a canonical divisor in X and a canonical divisor on a smooth model of X. Let X be a normal variety such that K_X is $\mathbb{Q}$-Cartier (we will need the condition that K_X is $\mathbb{Q}$-Cartier to do a pullback of it). Take a resolution of singularities $f : Y \to X$ and choose K_X and K_Y such that $f_*(K_Y) = K_X$. Define the *relatively canonical divisor* as $K_{Y/X} := K_Y - f^*(K_X)$ on Y where $f^*(K_X) := \frac{1}{r} f^*(rK_X)$ for any $r \in \mathbb{Z}_{>0}$ such that rK_X is Cartier. Notice that such $K_{Y/X}$ is independent of the choice of K_Y. Also, $K_{Y/X}$ is supported on the

exceptional locus of f. Let $\{E_1, \ldots, E_m\}$ be the exceptional prime divisors of f for some $m \in \mathbb{Z}_{>0}$, so we have

$$K_{Y/X} = \sum_{i=1}^{m} a_i E_i$$

for some $a_i \in \mathbb{Q}$. We denote $a(E_i, X) := a_i$.

Definition 3.2.4 (Discrepancies)**.** Let X be a normal variety such that K_X is $\mathbb{Q}$-Cartier (or X is called $\mathbb{Q}$-Gorenstein), and let E be a prime divisor over X. The number $a(E, X) \in \mathbb{Q}$ is called the *discrepancy* of E over X. The *minimal discrepancy* of X is defined as the infimum of the discrepancies $a(E, X)$, where E runs through all exceptional divisors over X.

Remark 3.2.5. Recall that a prime divisor E over X induces a valuation val_E. If two divisors E and E' over X induce the same valuation ($\mathrm{val}_E = \mathrm{val}_{E'}$), then we have $a(E, X) = \mathrm{val}_E(K_{Y/X}) = \mathrm{val}_{E'}(K_{Y'/X}) = a(E', X)$ where Y and Y' are the corresponding models. Therefore, in this sense, the discrepancy $a(E, X)$ is independent of the choice of resolution.

Discrepancies in birational geometry encode the information of X_{sing}. In particular, the minimal discrepancy is an important birational invariant of X_{sing}.

Here, we define the concept of terminal singularities.

Definition 3.2.6 (Terminal Singularities)**.** Let X be a $\mathbb{Q}$-Gorenstein normal variety. We say X has only *terminal singularities* if $a(E, X) > 0$ for all exceptional prime divisors E over X.

With the concept of terminal singularities, we can define minimal models / relatively minimal models in higher dimensions.

Definition 3.2.7 (Minimal Model)**.** Let X be a $\mathbb{Q}$-Gorenstein normal variety. X is called a *minimal model* if

- X has only terminal singularities,
- A canonical divisor K_X is nef.

Definition 3.2.8 (Relatively Minimal Model)**.** Let X be a variety, and let Y be a $\mathbb{Q}$-Gorenstein normal variety . Y is called a *relatively minimal model* over X if

- there is a projective birational morphism $f : Y \to X$,
- Y has only terminal singularities,
- A canonical divisor K_Y is f-nef. That is, $K_Y.C \geq 0$ for all f-exceptional curves $C \subseteq Y$.

Here is an important theorem that applies to varieties of general type (we will not define this term here).

Theorem 3.2.9 ([BCHM10])**.** *Let X be a variety of general type, then X is always birational to a minimal model.*

Terminal singularities appear in many research topics in algebraic geometry. In particular, people have tried classifying these kinds of singularities. For 3-folds, this has been done in [Dan82, MS84, Mor85, Rei83]. Notice that for X being terminal, its singular locus X_{sing} has the following lower bound for its codimension (see [KM98, Corollary 5.18, p. 159]).

$$\operatorname{codim}(X_{\text{sing}}, X) \geq 3.$$

Therefore, the singular locus X_{sing} of a terminal 3 -fold X is a set of isolated points.

3.3 3-Fold Terminal Singularities

Definition 3.3.1 (Gorenstein index)**.** Let X be a $\mathbb{Q}$-Gorenstein normal variety. The *Gorenstein index* of X is the smallest $m \in \mathbb{Z}_{>0}$ such that mK_X is Cartier.

For a germ of 3-dimensional terminal singularities (X, p), the classification for the Gorenstein index equal to 1 (or *Gorenstein terminal*, shortly) are done by M. Reid in [Rei83].

Theorem 3.3.2 ([Rei83])**.** *A germ of 3-fold Gorenstein singularity is terminal if and only if it is an isolated cDV point.*

For the cases of Gorenstein index greater than 1, the complete classification is treated in [Dan82, MS84, Mor85].

Let us introduce the classic notation to describe the classification. Denote the coordinates of the complex space $\mathbb{C}^4$ by (x, y, z, u). Let $\mathbb{Z}_m$ be a cyclic group of order m, and consider the action $\tau : \mathbb{Z}_m \times \mathbb{C}^4 \to \mathbb{C}^4$ by $\tau(x) = \zeta^{\alpha_1} x, \tau(y) = \zeta^{\alpha_2} y, \tau(z) = \zeta^{\alpha_3} z, \tau(u) = \zeta^{\alpha_4} u$ where ζ is a primitive m-root of unity and $\alpha_i \in \mathbb{Z}$ for $i = 1, 2, 3, 4$. The quotient variety of $\mathbb{C}^4$ with respect to this action is denoted by

$$\mathbb{C}^4/\frac{1}{m}(\alpha_1, \alpha_2, \alpha_3, \alpha_4).$$

Its ring of regular functions agrees with the ring of invariant regular functions with respect to $\frac{1}{m}(\alpha_1, \alpha_2, \alpha_3, \alpha_4)$ (see [CLS11, Chapter 5]).

Denote the ring of the complex convergent power series in the variables x, y, z, u by $\mathbb{C}\{x, y, z, u\}$. Let $\phi \in \mathbb{C}\{x, y, z, u\}$ be a $\mathbb{Z}_m$-semi-invariant with respect to the previous action determined by the vector $\frac{1}{m}(\alpha_1, \alpha_2, \alpha_3, \alpha_4)$. Then we obtain an induced action on the germ of hypersurface $(\phi = 0) \subseteq \mathbb{C}^4$. It induces a germ of subvariety of $\mathbb{C}^4/\frac{1}{m}(\alpha_1, \alpha_2, \alpha_3, \alpha_4)$, and it is denoted by $(\phi = 0) \subseteq \mathbb{C}^4/\frac{1}{m}(\alpha_1, \alpha_2, \alpha_3, \alpha_4)$. It is called a hyperquotient singularity.

With this notation, we have the following list for the classification:

Theorem 3.3.3. *([Dan82, MS84, Mor85, Hay99, Hay00, Che16]) Let X be a germ of 3-fold terminal singularity with Gorenstein index greater than 1, then it is isomorphic to one*

of the following form:

- *(type cA/m)*

$$(xy + f(z,u) = 0) \subseteq \mathbb{C}^4/\frac{1}{m}(a,-a,1,0)$$

where $a, m \in \mathbb{Z}_{>0}$ *with* $\gcd(a,m) = 1$ *and* $f(z,u) \in \mathbb{C}\{z,u\}$ *is a* $\mathbb{Z}_m$*-invariant.*

- *(type cAx/2)*

$$(x^2 + y^2 + f(z,u) = 0) \subseteq \mathbb{C}^4/\frac{1}{2}(0,1,1,1)$$

where $f(z,u) \in (z,u)^4 \subseteq \mathbb{C}\{z,u\}$ *is a* $\mathbb{Z}_2$*-invariant.*

- *(type cAx/4)*

$$(x^2 + y^2 + f(z,u) = 0) \subseteq \mathbb{C}^4/\frac{1}{4}(1,3,1,2)$$

where $f(z,u) \in \mathbb{C}\{z,u\}$ *is a* $\mathbb{Z}_4$*-semi-invariant such that* $f(z,u)$ *does not contain* u *as a monomial.*

- *(type cD/2)*

$$(\phi = 0) \subseteq \mathbb{C}^4/\frac{1}{2}(1,1,0,1)$$

where ϕ *has one of the following forms:*

  - *(cD/2-1)* $\phi = u^2 + xyz + x^{2a} + y^{2b} + z^c$ *where* $a, b \geq 2$*,* $c \geq 3$*.*
  - *(cD/2-2)* $\phi = u^2 + y^2 z + \lambda y x^{2a+1} + g(x,z)$ *where* $\lambda \in \mathbb{C}$*,* $a \geq 1$*,*

    $g(x,z) \in (x^4, x^2z^2, z^3) \subseteq \mathbb{C}\{x,z\}$*.*

- *(type cD/3)*

$$(\phi = 0) \subseteq \mathbb{C}^4/\frac{1}{3}(1,2,2,0)$$

where ϕ *has one of the following forms:*

  - *(cD/3-1)* $\phi = x^3 + u^2 + yz(y+z)$*.*
  - *(cD/3-2)* $\phi = x^3 + u^2 + yz^2 + xy^4\alpha(y^2) + y^6\beta(y^3)$

    where $\alpha(y^3), \beta(y^3) \in \mathbb{C}\{y^3\}$ *with* $4\alpha^3 + 27\beta^2 \neq 0$*.*
  - *(cD/3-3)* $\phi = x^3 + u^2 + y^3 + xyz^3\alpha(z^3) + xz^4\beta(z^3) + yz^5\gamma(z^3) + z^6\delta(z^3)$

    where $\alpha(z^3), \beta(z^3), \gamma(z^3), \delta(z^3) \in \mathbb{C}\{z^3\}$*.*

- *(type cE/2)*

$$(x^3 + u^2 + g(y,z) + h(y,z) = 0) \subseteq \mathbb{C}^4/\frac{1}{2}(0,1,1,1)$$

where $g(y,z) \in (y,z)^4 \subseteq \mathbb{C}\{y,z\}$, $h(y,z) \in (y,z)^4 \setminus (y,z)^5 \subseteq \mathbb{C}\{y,z\}$.

In particular, three-fold toric terminal singularities are studied in [Dan82] and [MS84]. The classification is completed by the following result in [KSB88].

Theorem 3.3.4 ([KSB88, Theorem 6.5])**.** *Let*

$$X = (\phi(x,y,z,u) = 0) \subseteq \mathbb{C}^4/\frac{1}{m}(\alpha_1, \alpha_2, \alpha_3, \alpha_4)$$

be one of the hyperquotient singularities listed in Theorem 3.3.3. Assume

- $\phi(x,y,z,u) \in \mathbb{C}[x,y,z,u]$ *defines an isolated singularity at* $(0,0,0,0) \in \mathbb{C}^4$*,*
- *The action* $\tau : \mathbb{Z}_m \times \mathbb{C}^4 \to \mathbb{C}^4$ *defined previously is free outside* $(0,0,0,0) \in \mathbb{C}^4$*.*

Then $(X, (0,0,0,0))$ *defines a 3-fold terminal singularity.*

Three-fold terminal singularities have been studied explicitly in many articles [Hay99, Hay00, Hay05, Che16]. One of the important tools for explicitly studying them is *divisorial contraction with minimal discrepancy* (or *divisorial blowing-ups with minimal discrepancy*).

Theorem 3.3.5 ([Kaw93, Mar96])**.** *For a 3-fold terminal singularity with Gorenstein index* $r \geq 1$*, its minimal discrepancy is exactly* $\frac{1}{r}$*.*

Definition 3.3.6 ([Hay99] and [Hay00])**.** Let X be a germ of 3-fold terminal singularity with Gorenstein index $m \geq 2$. A *divisorial contraction with minimal discrepancy* to X is a projective birational morphism $\pi : \overline{X} \to X$ such that

- $\overline{X}$ has only terminal singularities,
- the exceptional locus of π is a prime divisor E over X,
- and $K_{\overline{X}/X} = \frac{1}{m}E$, i.e. E computes the minimal discrepancy $a(E,X) = \frac{1}{m}$.

Remark 3.3.7. In the definition 3.3.6 of the divisorial contraction, we see that $-K_{\overline{X}}$ is π-ample.

The existence of such morphisms is proved by T. Hayakawa.

Theorem 3.3.8 ([Hay99, Hay00])**.** *Let X be a germ of 3-fold terminal singularity such that its Gorenstein index is greater than 1. Then there is a divisorial contraction with minimal discrepancy to X.*

Here is a rough idea of the proof from T. Hayakawa. Notice that 3-fold terminal singularities with Gorenstein indices $m > 1$ have been classified completely in Theorem 3.3.3. For any such singularity, Hayakawa selects some particular 4-tuple $\sigma = \frac{1}{m}(a, b, c, d) \in \mathbb{C}^4$ and applies the weighted blowing-up to the singularity with respect to such weight σ. Then he shows that such weighted blowing-up provides the divisorial contraction by establishing a correspondence between certain equivalences of maximal weighted pseudo-valuations with the parameter $\frac{1}{m}$ and the set of isomorphism classes of divisorial contraction with minimal discrepancy to X.

Chapter 4

Arc Spaces and the Nash Problem

In this chapter, we introduce some important concepts about arc spaces of a given variety X. Especially, we will formulate the Nash problem over given singularities. The readers can check the details in [IK03, CLNS18, dFD16, dF18].

4.1 Arc Spaces of a Variety

Nash first introduces the concept of arc spaces to study the singularities of a variety X in [Nas95]. Generally speaking, an arc of a variety X is a formal analytic curve in X, and the arc space X_∞ is the space which parametrizes all formal analytic curves in X. Here we rigorously define arcs of X in the language of the functor of points.

Definition 4.1.1. Let X be a scheme over $\mathbb{C}$, and let $\mathbb{K}$ be a $\mathbb{C}$-algebra. A $\mathbb{K}$-*arc* in X is a $\mathbb{K}[\![t]\!]$-valued point of X. In other words, a $\mathbb{K}$-arc on X is a morphism $\alpha \in \mathrm{Hom}_{\mathrm{Schemes}/\mathbb{C}}(\mathrm{Spec}(\mathbb{K}[\![t]\!]), X)$.

Theorem 4.1.2 ([CLNS18, Corollary 3.3.7] and [Bha16, Remark 4.6]). *The functor*

$$X_\infty(-) := \mathrm{Hom}_{\mathrm{Schemes}/\mathbb{C}}(\mathrm{Spec}(-[\![t]\!]), X) : (\mathbb{C}\text{–Algebras}) \to (\mathrm{Sets})$$

is representable by a scheme over $\mathbb{C}$. That is, there is a scheme Y over $\mathbb{C}$ such that

$$\mathrm{Hom}_{\mathrm{Schemes}/\mathbb{C}}(\mathrm{Spec}(\mathbb{K}[\![t]\!]), X) \cong \mathrm{Hom}_{\mathrm{Schemes}/\mathbb{C}}(\mathrm{Spec}(\mathbb{K}), Y).$$

By the Yoneda lemma, such a representative Y is unique up to isomorphism, and we denote it by X_∞.

Definition 4.1.3. This representative X_∞ is called the *arc space* of X.

Remark 4.1.4. Notice that X_∞ is just a scheme, not a variety.

Remark 4.1.5. In fact, this is not the original definition for the arc space of X. Originally, it is defined as the inverse limit of the jet schemes X_m of X for $m \in \mathbb{Z}_{\geq 0}$.

$$X_\infty := \varprojlim X_m.$$

From [Bha16], our definition is equivalent to the original one.

In many scenarios, we simply call a $\mathbb{K}$-arc an *arc* of X if it is unnecessary to specify the base $\mathbb{K}$.

Notice that when $\mathbb{K}$ is a field, $\mathbb{K}[\![t]\!]$ is a discrete valuation ring (DVR), so it has a unique maximal ideal $(t) \subset \mathbb{K}[\![t]\!]$, associated with the unique closed point in $\mathrm{Spec}(\mathbb{K}[\![t]\!])$. Denote this closed point of $\mathrm{Spec}(\mathbb{K}[\![t]\!])$ by 0 and the generic point by η. In fact, we have $\mathrm{Spec}(\mathbb{K}[\![t]\!]) = \{0, \eta\}$.

Example 4.1.6. Let $X = \mathrm{Spec}(\mathbb{C}[x,y]/(y^2 - x^3))$, then an arc is a morphsim of schemes $\phi : \mathrm{Spec}(\mathbb{K}[\![t]\!]) \to \mathrm{Spec}(\mathbb{C}[x,y]/(y^2 - x^3))$, which corresponds to a $\mathbb{C}$-algebra homomorphism $\phi^\# : \mathbb{C}[x,y]/(y^2 - x^3) \to \mathbb{K}[\![t]\!]$ and it is determined by $\phi^\#(x) = \alpha(t)$,

$\phi^{\#}(y) = \beta(t)$ for some $\alpha(t), \beta(t) \in \mathbb{K}[[t]]$. Notice that we have $\beta(t)^2 - \alpha(t)^3 = 0$ since $\phi^{\#}$ is a $\mathbb{C}$-algebra homomorphism. Therefore, we have

$$X_\infty(\mathbb{K}) = \{(\alpha(t), \beta(t)) \in \mathbb{K}[[t]] \mid \beta(t)^2 - \alpha(t)^3 = 0\}.$$

For the characterization of X_∞, let $(\alpha(t), \beta(t)) \in X_\infty$ and write $\alpha(t) = a_0 + a_1 t + a_2 t^2 + \dots$ and $\beta(t) = b_0 + b_1 t + b_2 t^2 + \dots$, then we obtain

$$(b_0 + b_1 t + b_2 t^2 + \dots)^2 - (a_0 + a_1 t + a_2 t^2 + \dots)^3 = 0.$$

Comparing the coefficients of t^i for $i = 0, 1, 2 \dots$ in this equation, we have the following relations:

$$\begin{aligned} b_0^2 - a_0^3 &= 0, \\ 2b_0 b_1 - 3a_0^2 a_1 &= 0, \\ b_1^2 + 2b_0 b_2 - 3a_0 (a_1)^2 - 3a_0^2 a_2 &= 0, \\ &\vdots \end{aligned}$$

Therefore, we have the following expression for X_∞.

$$X_\infty = \left\{ (a_0, a_1, a_2, \dots, b_0, b_1, b_2, \dots) \;\middle|\; \begin{array}{c} b_0^2 - a_0^3 = 0, \\ 2b_0 b_1 - 3a_0^2 a_1 = 0, \\ b_1^2 + 2b_0 b_2 - 3a_0 (a_1)^2 - 3a_0^2 a_2 = 0, \\ \vdots \end{array} \right\}.$$

Or equivalently,

$$X_\infty = \operatorname{Spec}\left(\frac{\mathbb{C}[a_0, a_1, a_2, \dots, b_0, b_1, b_2, \dots]}{(\, b_0^2 - a_0^3, 2b_0 b_1 - 3a_0^2 a_1, b_1^2 + 2b_0 b_2 - 3a_0 (a_1)^2 - 3a_0^2 a_2, \dots)} \right).$$

It is not hard to see that X_∞ is a scheme over $\mathbb{C}$, not a variety.

Example 4.1.7. In general, if $X = \mathrm{Spec}(k[x_0, \ldots, x_n]/(f_0, \ldots, f_m))$ is an affine variety, then an arc is a formal power series solution in $\mathbb{K}[[t]]$ of the system of equations $\{f_0 = 0, \ldots, f_m = 0\}$, and the arc space X_∞ collects all such formal power series solutions. In other words,

$$X_\infty(\mathbb{K}) = \{\, \alpha(t) \in (\mathbb{K}[[t]])^{n+1} \,|\, f_0(\alpha(t)) = 0, \ldots, f_m(\alpha(t)) = 0 \,\}.$$

Consider the canonical map $\mathbb{K}[[t]] \to \mathbb{K}$ defined by $t \mapsto 0$, which induces the map $\mathrm{Spec}(\mathbb{K}) \to \mathrm{Spec}(\mathbb{K}[[t]])$. Then we have

$$\mathrm{Hom}_{\mathrm{Schemes}/k}(\mathrm{Spec}(\mathbb{K}[[t]]), X) \to \mathrm{Hom}_{\mathrm{Schemes}/k}(\mathrm{Spec}(\mathbb{K}), X),$$

and it induces the canonical projection.

$$\pi_X : X_\infty \longrightarrow X$$

$$\alpha \longmapsto \alpha(0).$$

Let $S \subseteq X$, then $\pi_X^{-1}(S)$ is the set of all arcs passing through S.

$$\pi_X^{-1}(S) = \{\, \alpha \in X_\infty \,|\, \alpha(0) \in S \,\}.$$

If $p \in X$ is a non-singular point, then we have $\pi_X^{-1}(\{p\}) \cong \{p\} \times \mathbb{A}_k^\infty$. If $p \in X$ is singular, the situation will become complicated. This leads to the point of view of J.F. Nash [Nas95] in the next section:

What information can arcs passing through X_{sing} tell us about resolutions of X?

4.2 The Nash Problem: Formulation

In this section, we will formulate the so-called *Nash problem*. This is an interesting observation proposed by J. F. Nash in [Nas95]. From his point of view, the set of all arcs passing through the singular locus of X (or $\pi_X^{-1}(X_{\mathrm{sing}})$) can help us study resolutions of singularities of X.

Essential Divisors / Essential Valuations

We need some notions and facts to formulate Nash's observation. Let X be a variety, and we introduce the concept that a prime divisor over X "appears" in other models over X.

Definition 4.2.1 (Essential Divisors / Essential Valuations)**.** Let E be an exceptional prime divisor over X. E is called *essential* over X if any resolution $f : Y \to X$ is given, the center of val_E in Y is an irreducible component of $f^{-1}(X_{\mathrm{sing}})$. We say E *appears* in Y if the center of val_E is a divisor in Y. The divisorial valuation induced by an essential exceptional prime divisor of X is called an essential valuation of X.

For examples / non-examples of essential divisors / valuations, see [IK03, Example 2.4,2.5,2.6].

Nash Components / Nash Valuations

For the remainder of this chapter, we let $\mathbb{K}$ denote a field extension of $\mathbb{C}$.

Let X be a topological space, and let $x, y \in X$. We say y is a *generalization* of x (or x is a *specialization* of y) if $x \in \overline{\{y\}}$ in X, where $\overline{\{\ \}}$ denotes the closure in X.

Proposition 4.2.2 ([IK03])**.** *Let X be a variety, and let $\alpha : \mathrm{Spec}(\mathbb{K}[[t]]) \to X$ be a $\mathbb{K}$-arc such that $\alpha(0) \in X_{\mathrm{sing}}$. Then α has a generalization $\beta \in X_\infty$ such that $\beta(0) \in X_{\mathrm{sing}}$ and $\beta(\eta) \notin X_{\mathrm{sing}}$.*

Proof. See [IK03, Lemma 2.12] or [CLNS18, Chapter 3, Proposition 4.3.2]. □

Remark 4.2.3. The proof of this proposition actually needs the condition that X is over a field k of characteristic zero.

Proposition 4.2.4 ([IK03]). *$\pi_X^{-1}(X_{\mathrm{sing}})$ has the following decomposition*

$$\pi_X^{-1}(X_{\mathrm{sing}}) = \bigcup_{i\in I} C_i,$$

where the C_i are the irreducible components of $\pi_X^{-1}(X_{\mathrm{sing}})$, and each C_i contains arcs $\alpha \in X_\infty$ such that $\alpha(\eta) \notin X_{\mathrm{sing}}$.

Proof. See [IK03, Lemma 2.12]. □

Definition 4.2.5 (Nash Components of X)**.** These C_i are called the *Nash components* of X.

A $\mathbb{K}$-arc $\alpha : \mathrm{Spec}(\mathbb{K}[\![t]\!]) \to X$ induces a discrete semi-valuation

$$\mathrm{val}_\alpha(f) := \mathrm{ord}_t(f(\alpha(t)))$$

on the local ring $\mathcal{O}_{X,\pi_X(\alpha)}$, defined by taking composition with the t-adic valuation.

In general, this semi-valuation cannot be extended to a valuation defined on the function field $\mathbb{C}(X)$. This will happen if $\mathbb{C}(X) \subseteq \mathbb{K}((t))$ is a field extension, which is the same as the image of α being dense in X. This leads to the concept of *fat arcs*.

Definition 4.2.6 (Thin / Fat Arcs)**.** An arc $\alpha : \mathrm{Spec}(\mathbb{K}[\![t]\!]) \to X$ is called *thin* (or *algebraically thin*) if the image of α is contained in a proper closed subscheme of X. Otherwise, α is called *fat* (or *algebraically fat*). Equivalently, α is fat if and only if val_α is a valuation on $\mathbb{C}(X)$.

Now we define the semi-valuations of X induced by the Nash components of X, which will be shown that they are actually valuations of X. Let C_i be a Nash component, and let $\alpha_i \in C_i$ be the generic point. By Proposition 4.2.2, we have a generalization $\tilde{\alpha}_i \in \pi_X^{-1}(X_{\mathrm{sing}})$ such that $\tilde{\alpha}_i(\eta) \notin X_{\mathrm{sing}}$ and $\tilde{\alpha}_i(0) \in X_{\mathrm{sing}}$. Hence, we have $C_i = \overline{\{\alpha_i\}} \subseteq \overline{\{\tilde{\alpha}_i\}}$. Since C_i is an irreducible component of $\pi_X^{-1}(X_{\mathrm{sing}})$, we get

$\tilde{\alpha}_i$ is also the generic point of C_i (i.e., $\tilde{\alpha}_i = \alpha_i$, and this also implies $C_i \notin \pi_X^{-1}(X_{\mathrm{sing}})$). We obtain the semi-valuation

$$\begin{aligned} \mathrm{val}_{C_i} : \mathcal{O}_{X,\pi_X(\alpha_i)} &\to \mathbb{Z} \\ f &\mapsto \mathrm{ord}_t[f(\alpha_i(t))] \end{aligned}$$

We now explain how arcs of X with $\alpha(0) \in X_{\mathrm{sing}}$ can be lifted to arcs on a resolution of X.

Lemma 4.2.7. *Let $f : Y \to X$ be a resolution of singularities of X, and assume that it induces an isomorphism between $X \setminus Z$ and $Y \setminus f^{-1}(Z)$ for some proper closed subset $Z \subsetneq X$. Let C_i° denote the open subset of a given Nash component C_i consisting of arcs $\alpha : Spec(\mathbb{K}[\![t]\!]) \to X$ such that $\alpha(\eta) \notin Z$. Then for every $\alpha \in C_i^\circ$, the arc α can be uniquely lifted to an arc $\tilde{\alpha} : \mathrm{Spec}(\mathbb{K}[\![t]\!]) \to Y$.*

Proof. Notice that f is isomorphic outside Z, so we have the following commutative diagram.

$$\begin{array}{ccc} \mathrm{Spec}(\mathbb{K}((t))) & \longrightarrow & Y \\ \downarrow & & \downarrow f \\ \mathrm{Spec}(\mathbb{K}[\![t]\!]) & \xrightarrow{\alpha} & X \end{array}$$

where the downward arrow on the left is induced by the canonical inclusion to the fraction field $\mathbb{K}[\![t]\!] \hookrightarrow \mathbb{K}((t))$, and the top arrow $\mathrm{Spec}(\mathbb{K}((t))) \to Y$ is obtained by the condition $\alpha(\eta) \notin Z$. Notice that f is a proper morphism and $\mathbb{K}[\![t]\!]$ is a valuation ring, so we obtain a unique lifting $\tilde{\alpha} : \mathrm{Spec}(\mathbb{K}[\![t]\!]) \to Y$ by the valuative criterion of properness.

$$\begin{array}{ccc} \mathrm{Spec}(\mathbb{K}((t))) & \longrightarrow & Y \\ \downarrow & \overset{\exists\tilde{\alpha}}{\nearrow} & \downarrow f \\ \mathrm{Spec}(\mathbb{K}[\![t]\!]) & \xrightarrow{\alpha} & X \end{array}$$

□

Remark 4.2.8. If $f : Y \to X$ is a Hironaka resolution of X, we have $Z = X_{\text{sing}}$.

The Nash Maps / The Nash Problem

Theorem 4.2.9 ([Nas95, IK03, CLNS18])**.** *Let X be a singular variety with the singular locus X_{sing}, and let C be a Nash component of X. Then there is an unique essential exceptional prime divisor E over X such that*

$$\text{val}_C = \text{val}_E.$$

In particular, val_C is a valuation and it is essential. Also, there are only finitely many Nash components of X.

Proof. Let $f : Y \to X$ be a resolution of singularities of X. Without loss of generality, we can assume that f is a Hironaka resolution of X. We have a morphism of arc spaces $f_\infty : Y_\infty \to X_\infty$ induced by f (see [CLNS18, 3.1.2 in Chapter 3]). Consider the exceptional locus $f^{-1}(X_{\text{sing}}) = Z_1 \cup \cdots \cup Z_m$ with $\{Z_i\}_{i=1}^m$ be the family of irreducible components of $f^{-1}(X_{\text{sing}})$.

f_∞ induces the following continuous bijection.

$$\tilde{f}_\infty : \pi_Y^{-1}(f^{-1}(X_{\text{sing}})) \setminus (f^{-1}(X_{\text{sing}}))_\infty \to \pi_X^{-1}(X_{\text{sing}}) \setminus (X_{\text{sing}})_\infty$$

where $\tilde{f}_\infty := f_\infty|_{\pi_Y^{-1}(f^{-1}(X_{\text{sing}})) \setminus (f^{-1}(X_{\text{sing}}))_\infty}$ (see [CLNS18, Proposition 4.4.2 in Chapter 3]). As Y is smooth, $\{\pi_Y^{-1}(Z_i)\}_{i=1}^m$ are the irreducible components of $\pi_Y^{-1}(f^{-1}(X_{\text{sing}}))$

Since C is an irreducible component of $\pi_X^{-1}(X_{\text{sing}})$, there is a unique $\pi_Y^{-1}(Z_j)$ for some $j \in \{1, \ldots, m\}$ such that $\tilde{f}_\infty$ restricts to the following continuous bijection.

$$\pi_Y^{-1}(Z_j) \setminus (f^{-1}(X_{\text{sing}}))_\infty \xrightarrow{\tilde{f}_\infty} C \setminus (X_{\text{sing}})_\infty,$$

and we have $\tilde{f}_\infty(\beta) = \alpha$, where α is the generic point of C and β is the generic point of $\pi_Y^{-1}(Z_j)$. Notice that β is a fat arc since Y is smooth, therefore α is also fat, which means that $\text{val}_C = \text{val}_\alpha$ is a valuation of X.

Take the blowing-up $\pi : \tilde{Y} \to Y$ along Z_j and let $E \subseteq \pi^{-1}(Z_j)$ be the unique exceptional prime divisor dominating Z_j, then we have

$$\mathrm{val}_C = \mathrm{val}_E$$

because $(f_\infty \circ \pi_\infty)(\pi_{\tilde{Y}}^{-1}(E)) = C$, which implies that α is the image of the generic point of $\pi_{\tilde{Y}}^{-1}(E)$.

The whole argument does not depend on the choice of resolution $f : Y \to X$, so we have that val_E is an essential valuation of X. □

Definition 4.2.10 (Nash Valuations of X)**.** Let $C_1, \ldots, C_\ell$ be the Nash components of X, then the induced valuations $\mathrm{val}_{C_1}, \ldots, \mathrm{val}_{C_\ell}$ are called the *Nash valuations* of X.

In other words, we have the following correspondence

$$\mathcal{N}_X : \{\text{Nash components of } X\} \to \{\text{essential valuations of } X\}$$
$$C \longmapsto \mathrm{val}_C = \mathrm{val}_E$$

is injective. $\mathcal{N}_X$ is called the *Nash map* of X. Therefore, we have

$$\{\text{Nash valuations of } X\} \subseteq \{\text{essential valuations of } X\}.$$

Consequently, we obtain an intrinsic method to identify an essential valuation of X without requiring to consider all resolutions of singularities of X. It is natural to ask if any essential valuation can be identified in this way:

$$\text{Is the Nash map } \mathcal{N}_X \text{ surjective?}$$

Now we can formulate *the Nash Problem of* X:

$$\text{Understand the image of the Nash map } N_X.$$

4.3 The Nash Problem: Known Results

Unfortunately, it is known that the Nash maps are in general not surjective for $\dim(X) \geq 3$. However, we still have several important cases for which the Nash maps are surjective.

Nash maps are surjective for the following cases:

- Curve Singularities
- Surface Singularities [FdBP12]
- Toric Singularities in arbitrary dimension [IK03]
- Quasi-Ordinary Hypersurface Singularities in arbitrary dimension [GPH09]
- Non-Rational Normal Varieties with Torus Action of Complexity 1 in arbitrary dimension [BLM25]

There are some examples that show that the Nash maps need not be surjective.

Theorem 4.3.1 ([IK03, Proposition 4.5])**.** *Consider the isolated hypersurface singularity given by*

$$X = (x_1^3 + x_2^3 + x_3^3 + x_4^3 + x_5^6 = 0) \subseteq \mathbb{A}_k^5$$

over an algebraically closed field k with $\operatorname{char}(k) \neq 2, 3$. Then the Nash map is not surjective. In fact, it has only one Nash valuation and exactly two essential valuations.

Theorem 4.3.2 ([dF13, Theorem 6.1])**.** *Let*

$$X = ((x_2^2 + x_3^2)x_4 + x_1^3 + x_2^3 + x_3^3 + x_4^5 + x_4^6 = 0) \subseteq \mathbb{A}_{\mathbb{C}}^4.$$

Then the Nash map is not surjective.

4.4 The Nash Problem: Further Developments

We know that a Nash map $\mathcal{N}_X$ is in general not surjective for $\dim(X) \geq 3$, so the next step is to investigate how to characterize the image of $\mathcal{N}_X$.

Here is a result for non-uniruled divisors. Recall that a prime divisor E is *uniruled* if there is a variety E' with $\dim(E') = \dim(X) - 1$ and a dominant rational map $\phi : E' \times \mathbb{P}^1_k \cdots \to E$.

Theorem 4.4.1 ([LJR12, Theorem 3.3])**.** *Let X be a singular variety, and let E be a non-uniruled prime divisor over X. Then its divisorial valuation val_E is a Nash valuation of X.*

Proof. See [LJR12]. □

Recently, T. de Fernex and R. Docampo show that we can use a relatively minimal model to characterize it in [dFD16].

Definition 4.4.2 (Terminal Valuations of X, [dFD16])**.** Let X be a variety. A valuation val_E of X is called a *terminal valuation* of X if E is an exceptional prime divisor on a relatively minimal model over X.

Theorem 4.4.3 ([dFD16, Theorem 1.1])**.** *Let X be a variety, then every terminal valuation of X is a Nash valuation of X.*

Proof. See [dFD16]. □

Remark 4.4.4. Notice that two relatively minimal models over X are connected by sequence of flops ([BCHM10, Corollary 1.1.3]), which are isomorphisms in codimension 1. In particular, this implies that terminal valuations of X appear on every relatively minimal model over X. Thus, it is enough to consider any one such model to identify them.

Based on this result, we have the following relations of valuations.

$$\{\text{terminal vals. of } X\} \subseteq \{\text{Nash vals. of } X\} \subseteq \{\text{essential vals. of } X\}$$

Notice that this theorem alternatively proves that Nash maps are surjective for surface singularities. Indeed, every relatively minimal model is non-singular for surfaces, so we have

$$\{\text{essential vals. of } X\} \subseteq \{\text{terminal vals. of } X\} \qquad \text{when } \dim X \leq 2.$$

This approach is useful for all types of singularities except for the following two cases:

(1) X has only terminal singularities.

(2) X has small resolutions (i.e., resolutions with exceptional locus having codimension at least 2).

There is no terminal valuations in both cases, and this motivates the whole work of this dissertation. In other words,

find an alternative characterization for the image of a Nash map over terminal singularities.

In [JK13], J. Kollár and J.M. Johnson provide some characterizations of the singularities of type cA in arbitrary dimensions. Recently, there is a result proven by H.-K. Chen [Che23] for the 3-fold terminal singularity of type cA/m.

Theorem 4.4.5 ([Che23, Theorem 1.1])**.** *Let (X, p) be a germ of 3-fold terminal singularity of type cA/m for some $m \in \mathbb{Z}_{>0}$, then there is an one-to-one correspondence between Nash valuations over X and exceptional prime divisors over X whose discrepancies are less than or equal to one.*

On the other hand, the example in [IK03, Proposition 4.5] also mentions that its Nash valuation of X has minimal discrepancy, which is 1.

These observations serve as evidence that discrepancies might provide a useful criterion to identify whether a given valuation of X is Nash or not. In particular, we would like to study the valuations val_E with $a(E, X) \leq 1$, and this condition is also motivated by our result in Chapter 6.

Chapter 5

Some Toric Geometry

In this chapter, we will introduce some fundamental notions and tools in toric algebraic geometry that will be used in the next chapter. Most of their details can be found in [CLS11, Mat02].

From a practical perspective, toric varieties serve as important candidates for testing hypothesis or conjectures in algebraic geometry.

5.1 Definitions, Results, and Notations

Definition 5.1.1 (Toric Variety)**.** A *toric variety* X is a variety that contains an algebraic torus $T_N \cong (\mathbb{C}^*)^n$ as a Zariski open subset such that the action of T_N on itself extends to an algebraic action of T_N on X (i.e., an action $T_N \times X \to X$ given by a morphism).

It is well-known that toric varieties admit a combinatorial construction via fans and lattices, which makes them suitable objects to proceed computations. Here we briefly recall the construction.

Let $N \cong \mathbb{Z}^n$ be a lattice of rank n and $M = \mathrm{Hom}_{\mathbb{Z}}(N, \mathbb{Z})$ be its dual lattice. Denoted $N_{\mathbb{R}} := N \otimes_{\mathbb{Z}} \mathbb{R}$, $N_{\mathbb{Q}} := N \otimes_{\mathbb{Z}} \mathbb{Q}$, $M_{\mathbb{R}} := M \otimes_{\mathbb{Z}} \mathbb{R}$, and $M_{\mathbb{Q}} := M \otimes_{\mathbb{Z}} \mathbb{Q}$.

Definition 5.1.2 (Fans)**.** A *fan* Δ in $N_{\mathbb{R}}$ is a finite collection of cones $\sigma \subseteq N_{\mathbb{R}}$ such that

- Every $\sigma \in \Delta$ is a strongly convex rational polyhedral cone,
- For every $\sigma \in \Delta$, all its faces are also in Δ,
- For every $\sigma_1, \sigma_2 \in \Delta$, their intersection $\sigma_1 \cap \sigma_2$ is a face of σ_1 and σ_2 (this implies $\sigma_1 \cap \sigma_2 \in \Delta$).

A toric variety $X(\Delta)$ associated with a fan Δ is obtained by gluing the affine toric varieties

$$U_\sigma := \mathrm{Spec}(\mathbb{C}[\sigma^\vee \cap M])$$

for all $\sigma \in \Delta$ in the following way: Two affine patches U_{σ_1} and U_{σ_2} are glued together via $U_{\sigma_1 \cap \sigma_2}$, which is an affine toric variety associated with their intersecting face.

Theorem 5.1.3. *Let Δ be a fan in $N_{\mathbb{R}}$, then $X(\Delta)$ is a normal separated toric variety.*

Proof. See [CLS11, Theorem 3.1.5, p. 107]. □

With this characterization, we can express the geometric properties of toric varieties in combinatoric ways. Here we list some of them without providing proofs. The readers can check their proofs in [CLS11, Mat02]

Definition 5.1.4. Let $\sigma \subseteq N_{\mathbb{R}}$ be a strongly convex rational polyhedral cone.

- σ is called *smooth* if its minimal generators form part of a $\mathbb{Z}$-basis of N.
- σ is called *singular* if it is not smooth.
- σ is called *simplicial* if its minimal generators are linearly independent over $\mathbb{R}$.

Definition 5.1.5. Let Δ be a fan in $N_{\mathbb{R}}$.

- Δ is *smooth* (or *regular*) if every cone $\sigma \in \Delta$ is smooth (or regular).
- Δ is simplicial if every cone $\sigma \in N_{\mathbb{R}}$ is simplicial.
- Δ is *complete* if its support $|\Delta| := \bigcup_{\sigma \in \Delta} \sigma = N_{\mathbb{R}}$.

Theorem 5.1.6. *Let $X(\Delta)$ be a toric variety associated with a fan Δ in $N_{\mathbb{R}}$, then*

(1) $X(\Delta)$ is a smooth (or non-singular) variety if and only if the fan Δ is smooth.

(2) $X(\Delta)$ is factorial (every Weil divisor is Cartier) if and only if Δ is smooth.

(3) $X(\Delta)$ is $\mathbb{Q}$-factorial (every Weil divisor is $\mathbb{Q}$-Cartier) if and only if Δ is simplicial.

(4) $X(\Delta)$ is compact in the classical topology if and only if Δ is complete.

Proof. For (1) and (4), see [CLS11, Theorem 3.1.19, p. 113]. For (2), see [CLS11, Proposition 4.2.6, p. 179]. For (3), see [CLS11, Proposition 4.2.7, p. 180]. □

5.2 Some Birational Geometry of Toric Varieties

Here we recall some characterizations for $\mathbb{Q}$-Gorenstein toric varieties with terminal singularities. One can check details from [CLS11] and [Mat02].

Proposition 5.2.1. *Let $X(\Delta)$ be toric variety associtated with a fan Δ.*

(1) Its canonical divisor $K_{X(\Delta)}$ is $\mathbb{Q}$-Cartier (i.e. $X(\Delta)$ is $\mathbb{Q}$-Gorenstein) if and only if for any cone $\sigma \in \Delta$, there exists an $m_\sigma \in M_{\mathbb{Q}}$ such that

$$\langle m_\sigma, v_i \rangle = 1$$

for every primitive element $v_i \in N$ with $\langle v_i \rangle \in \Delta$ and $\langle v_i \rangle \subset \sigma$.

(2) Assume $K_{X(\Delta)}$ is $\mathbb{Q}$-Cartier. For a singular cone $\sigma \in \Delta$, the discrepancy can be expressed as follows.

$$a(E, X(\Delta)) = \langle m_\sigma, u_E \rangle - 1$$

where $u_E \in \sigma \cap N$ is the primitive vector corresponding to the divisor E over X.

(3) Assume $K_{X(\Delta)}$ is $\mathbb{Q}$-Cartier, then $X(\Delta)$ has terminal singularities if and only if

$$\{w \in \sigma \cap N \mid \langle m_\sigma, w\rangle \leq 1\} = \{0\} \cup \{v_i \in N \mid v_i : \textit{primitive}, \langle v_i\rangle \in \Delta \textit{ and } \langle v_i\rangle \subset \sigma\}$$

for every $\sigma \in \Delta$. In particular, we have

$$\langle m_\sigma, u\rangle > 1$$

for a non-zero $u \in (\sigma \cap N) \setminus \{v_i \in N \mid v_i : \textit{primitive}, \langle v_i\rangle \in \Delta \textit{ and } \langle v_i\rangle \subset \sigma\}$.

Proof. See [Mat02, Proposition 14-3-1, p. 442], [CLS11, Lemma 11.4.10, p. 550], and [CLS11, Proposition 11.4.12, p. 550]. □

Chapter 6

Nash Valuations over Toric Terminal Singularities

It is natural to consider the Nash problem over toric terminal singularities as our initial cases. In this chapter, we present our results for the Nash problem over toric terminal singularities in any dimension. In particular, we prove that any exceptional prime divisor E over toric terminal singularities with discrepancy

$$a(E, X) \leq 1$$

will induce a Nash valuation.

Let Δ be a fan in $N_{\mathbb{R}}$. Given a cone $\sigma \in \Delta$, we consider the following partial order in $\sigma \cap N$.

Definition 6.0.1. Let σ be a strongly convex polyhedral cone, and let $u_1, u_2 \in \sigma \cap N$. We define $u_1 \leq u_2$ if $u_2 \in u_1 + \sigma$. i.e., there exists a $u' \in \sigma \cap N$ such that $u_2 = u_1 + u'$. For a subset $A \subset \sigma \cap N$, $u \in A$ is called minimal in A if $v \in A$ with $v \leq u$ implies $u = v$.

Consider

$$S_\sigma := (\bigcup_{\tau \preceq \sigma, \tau:\text{singular}} \overset{\circ}{\tau}) \cap N$$

for a given cone $\sigma \in \Delta$. $\tau \preceq \sigma$ means that τ is a face of σ, and $\overset{\circ}{\tau}$ denotes the relative interior of τ.

6.1 Discrepancy bounded by 1 induces Nash Valuations

Here we provide a sufficient condition for being a Nash valuation over toric terminal singularities in arbitrary dimensions.

Proposition 6.1.1. *Let Δ be a fan associated to a toric variety $X(\Delta)$ of dimension at least 3, and let $\sigma \in \Delta$ be a singular cone which defines a terminal singularity. A primitive element $w \in \overset{\circ}{\sigma} \cap N$ is minimal in S_σ if $\langle m_\sigma, w \rangle \leq 2$.*

Proof. We prove its contrapositive statement. Let $w \in \overset{\circ}{\sigma} \cap N$ which is not minimal in S_σ, so there exists a non-zero element $u \in S_\sigma$ such that $u \leq w$. i.e. $w = u + u'$ for some non-zero $u' \in \sigma \cap N$. Notice that $\langle m_\sigma, v_i \rangle = 1$ because $K_{X(\Delta)}$ is $\mathbb{Q}$-Cartier, and

$$\langle m_\sigma, u \rangle > 1$$

for a non-zero $u \in (\sigma \cap N) \setminus \{v_i \in N \mid v_i : \text{primitive}, \langle v_i \rangle \in \Delta \text{ and } \langle v_i \rangle \subset \sigma\}$ since the U_σ has a terminal singularity.

We have the following two cases for u':

Case 1 If $u' \in (\sigma \cap N) \setminus (\cup_i \langle v_i \rangle)$, then we have $\langle m_\sigma, u' \rangle > 1$, and

$$\langle m_\sigma, w \rangle = \langle m_\sigma, u \rangle + \langle m_\sigma, u' \rangle > 1 + 1 = 2.$$

Case 2 If $u' \in \cup_i \langle v_i \rangle$, then we have $u' = \alpha v_j$ for some v_j and some $\alpha \in \mathbb{Z}_{>0}$, and we get $\langle .m_\sigma, u' \rangle = \alpha \langle m_\sigma, v_j \rangle = \alpha \cdot 1 \geq 1$. So

$$\langle m_\sigma, w \rangle = \langle m_\sigma, u \rangle + \langle m_\sigma, u' \rangle > 1 + 1 = 2.$$

Therefore, we obtain

$$\langle m_\sigma, w \rangle > 2.$$

□

Theorem 6.1.2 ([IK03, 3. The Nash problem for toric singularities])**.** *Let $X(\Delta)$ be an affine toric variety in any dimension, then its Nash map is bijective. Moreover, every Nash valuation of $X(\Delta)$ (also being an essential valuation of $X(\Delta)$) is one-to-one corresponded to a minimal element in S_σ for some cone $\sigma \in \Delta$.*

By combining Proposition 6.1.1 and Theorem 6.1.2, we obtain the following Theorem.

Theorem 6.1.3. *Let $X(\Delta)$ be a toric variety with $\mathbb{Q}$-Cartier $K_{X(\Delta)}$ of dimension at least 3 having only terminal singularities (we have $a(E, X(\Delta)) > 0$), and let E be an exceptional prime divisor over $X(\Delta)$. If its discrepancy $a(E, X(\Delta)) \leq 1$, then the induced valuation val_E is a Nash valuation of $X(\Delta)$.*

Proof. Recall that for every exceptional prime divisor E over a toric variety, its discrepancy is $a(E, X(\Delta)) = \langle m_\sigma, w \rangle - 1$, where $w \in \mathring{\sigma} \cap N$ is the primitive element of a ray $\rho_E := \langle w \rangle \subset \mathring{\sigma}$ corresponding to E.

By assumption, since $a(E, X(\Delta)) = \langle m_\sigma, w \rangle - 1 \leq 1$, we get $\langle m_\sigma, w \rangle \leq 2$, and we obtain $w \in \mathring{\sigma} \cap N$ is minimal in S_σ by Proposition 6.1.1. Therefore, the induced valuation val_E is a Nash valuation of $X(\Delta)$ according to Theorem 6.1.2. □

6.2 Nash Valuations have Discrepancy bounded by 1

Next, we show that the converse of Theorem 6.1.3 is also true if $X(\Delta)$ is a toric terminal 3-fold. To prove this, we need the notion of heights of elements in the Hilbert basis for a cone.

Recall that for a given polyhedral cone σ, there exists a set $\mathcal{H}_N(\sigma) \subset \sigma \cap N$ such that

(1) Every $v \in \sigma \cap N$ can be expressed as a linear combination of elements in $\mathcal{H}_N(\sigma)$ over $\mathbb{Z}_{\geq 0}$, i.e. $v \in \sum_{h \in H_N(\sigma)} \mathbb{Z}_{\geq 0} h$,

(2) $\mathcal{H}_N(\sigma)$ has minimal cardinality with respect to all subsets of $\sigma \cap N$ for which (1) holds.

This set $\mathcal{H}_N(\sigma)$ is called the *Hilbert basis* for σ. Moreover, if σ is a pointed cone (i.e., $\sigma \cap -\sigma = \{0\}$), then $\mathcal{H}_N(\sigma)$ is uniquely determined by the following expression.

$$\mathcal{H}_N(\sigma) = \{v \in (\sigma \cap N) \setminus \{0\} \mid v \text{ is not the sum of two other vectors in } (\sigma \cap N) \setminus \{0\}\}.$$

Definition 6.2.1. Let $\sigma = \sum_{i=1}^{m} \mathbb{R}_{\geq 0} v_i \subset N_{\mathbb{R}}$ be a pointed cone, and let $u \in \mathcal{H}_N(\sigma)$. The number defined to be

$$g_\sigma(u) := \min\{\sum_{i=1}^{m} \lambda_i \mid u = \sum_{i=1}^{m} \lambda_i v_i, \lambda_i \geq 0, 1 \leq i \leq m\}$$

is called the *height* of u.

Proposition 6.2.2 ([HW97, LTZ93, EW91])**.** *Let $\sigma \subset N_{\mathbb{R}} \cong \mathbb{R}^d$ be a pointed cone, and let $u \in \mathcal{H}_N(\sigma)$. We have an upper bound for $g_\sigma(u)$:*

$$g_\sigma(u) < d - 1$$

for $d \geq 2$.

Corollary 6.2.3. *Let $X(\Delta)$ be a toric terminal 3-fold, then a valuation val_E associated to an exceptional prime divisor E over $X(\Delta)$ is a Nash valuation of $X(\Delta)$ if and only if*

$$a(E, X(\Delta)) \leq 1.$$

Proof. For the sufficient condition, apply Theorem 6.1.3. For the necessary condition, let E be an exceptional prime divisor over $X(\Delta)$ inducing a Nash valuation val_E. By [IK03], val_E is corresponded to a $w \in \overset{\circ}{\sigma} \cap N$ which is minimal in S_σ for some $\sigma \in \Delta$ being singular.

We have the following two cases for w:

Case 1. If $w \notin \mathcal{H}_N(\sigma)$, we have $w = v_1 + v_2$ for some primitive elements $v_1, v_2 \in \sigma \cap N$ such that $\langle v_1 \rangle$, $\langle v_2 \rangle \in \Delta$ and $\langle v_1 \rangle$, $\langle v_2 \rangle \subset \sigma$. Then we obtain

$$\langle m_\sigma, w \rangle = \langle m_\sigma, v_1 \rangle + \langle m_\sigma, v_2 \rangle = 1 + 1 = 2.$$

Notice that we use $\langle m_\sigma, v_i \rangle = 1$ for all $i = 1, 2$ since $X(\Delta)$ is $\mathbb{Q}$-Gorenstein ($X(\Delta)$ is terminal).

Case 2. If $w \in \mathcal{H}_N(\sigma)$, then we can express $w = \sum_{i=1}^m \lambda_i v_i$, where $\sigma = \sum_{i=1}^m \mathbb{R}_{\geq 0} v_i$. Then we get

$$\langle m_\sigma, w \rangle = \sum_{i=1}^m \lambda_i \langle m_\sigma, v_i \rangle = \sum_{i=1}^m \lambda_i.$$

Notice that we use $\langle m_\sigma, v_i \rangle = 1$ for all $i = 1 \ldots m$ since $X(\Delta)$ is $\mathbb{Q}$-Gorenstein. Now we consider $w = \sum_{i=1}^m \lambda_i v_i$ minimizing $\sum_{i=1}^m \lambda_i$, then we get

$$\langle m_\sigma, w \rangle = g_\sigma(w).$$

We obtain $\langle m_\sigma, w \rangle = g_\sigma(w) < 3 - 1 = 2$ by Proposition 6.2.2 (notice that $X(\Delta)$ is a 3-fold).

Therefore, we have $\langle m_\sigma, w \rangle \leq 2$ by Case 1 and Case 2, which implies that

$$a(E, X(\Delta)) = \langle m_\sigma, w \rangle - 1 \leq 2 - 1 = 1. \qquad \square$$

Remark 6.2.4. Notice that in the proof of the necessary condition of Corollary 6.2.3, we do not need the assumption $X(\Delta)$ is terminal. In fact, we use the condition that $X(\Delta)$ is $\mathbb{Q}$-Gorenstein.

6.3 Main Result and Conjectures

By combining the previous results, the condition $a(E, X(\Delta)) \leq 1$ fully characterizes Nash valuations over 3-fold toric terminal singularities.

Theorem 6.3.1. *Let $X(\Delta)$ be a toric terminal 3-fold, and val_E be a valuation associated to an exceptional prime divisor E over $X(\Delta)$. Then the following statements are equivalent:*

(1) val_E is essential over $X(\Delta)$.

(2) val_E is Nash over $X(\Delta)$.

(3) $a(E, X(\Delta)) \leq 1$.

Proof. For the equivalence (1)⇔(2) apply Theorem 6.1.2, and for (2)⇔(3) apply Corollary 6.2.3. □

Theorem 6.1.3 (or Theorem 6.3.1) provides a potential approach that we can try using discrepancies to characterize Nash valuations over terminal singularities. In particular, we propose the following conjecture.

Conjecture 6.3.2 (Conjecture A)**.** *Let (X, p) be a germ of 3-fold terminal with the Gorenstein index $m \geq 1$, then every exceptional prime divisor E with discrepancy $a(X, E) \leq 1$ induces a Nash valuation over (X, p).*

We propose a weaker version of Conjecture A.

Conjecture 6.3.3 (Conjecture B)**.** *Let (X, p) be a germ of 3-fold terminal with the Gorenstein index $m \geq 1$, then every exceptional prime divisor E with discrepancy $a(X, E) = \frac{1}{m}$ (i.e. the minimal discrepancy of X) induces a Nash valuation over (X, p).*

In this chapter, we prove that Conjecture A holds for toric terminal singularities for dimensions at least 3. In the next chapter, we will prove that for (X, p) is a germ of 3-fold terminal singularity of type cAx/2, then Conjecture B is true.

Chapter 7

Nash Valuations over 3-Fold Terminal Singularities of Type cAx/2

In this chapter, we will prove that the exceptional prime divisors with minimal discrepancy (it is $\frac{1}{2}$ in this case) will induce Nash valuations over 3-fold terminal singularities X of type cAx/2.

Let (X, p) be a germ of 3-fold terminal singularity of type cAx/2. It admits an embedding

$$(x^2 + y^2 + f(z, u) = 0) \subseteq \mathbb{C}^4/\frac{1}{2}(0, 1, 1, 1)$$

such that $p = (0, 0, 0, 0) \in \mathbb{C}^4/\frac{1}{2}(0, 1, 1, 1)$ and $f(z, u) \in (z, u)^4 \subseteq \mathbb{C}\{z, u\}$ is a $\mathbb{Z}_2$-invariant.

Here is our main Theorem in this chapter.

Theorem 7.0.1. *Let (X, p) be a germ of 3-fold terminal singularity of type cAx/2, then any prime divisor E over (X, p) with the minimal discrepancy (it is $\frac{1}{2}$ in this case) induces a Nash valuation val_E over (X, p).*

In other words, we prove that Conjecture B holds for the case (X, p) being a 3-fold terminal singularity of type cAx/2.

7.1 Candidates for Nash Valuations

Let (X, p) be a germ of terminal singularity. It makes sense to choose a prime divisor E over (X, p) with $a(E, X) \leq 1$ to be a candidate to produce a Nash valuation of X because val_E is, in fact, an essential valuation of X.

Proposition 7.1.1 ([JK13, Proposition 24])**.** *Let (X, p) be a germ of canonical singularity (see [KM98, Mat02]), and let E be an exceptional prime divisor over X centered at p and*

$$a(E, X) < 1 + \textit{the minimal discrepancy of } (X, p).$$

Then val_E is an essential valuation of X.

Corollary 7.1.2. *Let (X, p) be a 3-fold terminal singularity with Gorenstein index $m \geq 1$, then any exceptional prime divisor E over (X, p) such that $a(E, X) \leq 1$ is essential. In particular, any exceptional divisor over (X, p) with minimal discrepancy (in this case it is $\frac{1}{m}$) is essential.*

7.2 Setup for the Proof

Define a partial order among the valuations over (X, p):

$$\mathrm{val}_E \leq_X \mathrm{val}_F \quad \text{if} \quad \mathrm{val}_E(f) \leq \mathrm{val}_F(f) \quad \text{for all} \quad f \in \mathcal{O}_{X,p}$$

Lemma 7.2.1. *Let X be a singular variety, and let E be an exceptional prime divisor over X such that val_E is non-Nash over X. Then there exists another prime divisor F over X*

such that

$$\mathrm{val}_F <_X \mathrm{val}_E.$$

Proof. Define

$$N_G := \overline{\{f_\infty(\pi_Y^{-1}(G))\}} = \overline{\{\alpha \in X_\infty \mid \mathrm{val}_\alpha = \mathrm{val}_G\}}$$

for a prime divisor G in some Y such that $f : Y \to X$ is a resolution of X, and $\overline{\{\quad\}}$ denotes the Zariski closure in X_∞.

Let E be an exceptional prime divisor over X such that val_E is non-Nash over X. Since E is non-Nash, we have that N_E is contained in a Nash component of X. Let F be the prime divisor such that val_F is the Nash valuation of X induced by such the Nash component, then we have

$$N_E \subsetneq N_F,$$

and this implies $\mathrm{val}_E <_X \mathrm{val}_F$. □

Let (X, p) be a 3-fold terminal singularity with the Gorenstein index $r \geq 1$, and let $\pi : \overline{X} \to X$ be a divisorial contraction with minimal discrepancy in Definition 3.3.6. Denote $E = \pi^{-1}(p)$ to be the $\mathbb{Q}$-Cartier exceptional prime divisor in $\overline{X}$ obtained by π. Let F be an essential exceptional prime divisor over X centered at p. Here we derive an equation relating val_E and val_F.

Since $\pi : \overline{X} \to X$ is a divisorial contraction, $\overline{X}$ has only terminal singularities, which are isolated points (notice that $\mathrm{codim}(\overline{X}_{\mathrm{sing}}, \overline{X}) \geq 3$ since $\overline{X}$ is terminal). Let $\phi : Y \to \overline{X}$ be a resolution. Notice that val_F is essential over X, so F appears in Y. Hence, F is centered at some terminal singular point $q \in \overline{X}$.

Let $g \in m_{\overline{X},q} \setminus \mathcal{I}_{E,q}$ be irreducible, where $m_{\overline{X},q} \subseteq \mathcal{O}_{\overline{X},q}$ is the maximal ideal, and $\mathcal{I}_E$ is the sheaf of ideals associated with E. Let $G = \mathrm{div}_{\overline{X}}(g) \subseteq \overline{X}$ be the principal divisor induced by g, which is a prime divisor in $\overline{X}$.

Let $h \in \mathcal{I}_{\pi_* G,p}$, where $\pi_* G$ is the push-forward of G via π in X, which is a Weil divisor containing p, and $\mathcal{I}_{\pi_* G}$ is the sheaf of ideals associated with $\pi_* G$. Let $H = \mathrm{div}_X(h) \subseteq X$ be the principal divisor induced by h, which is Cartier. Then

$$\pi^* H = aG + bE + A$$

where $a, b \in \mathbb{Z}_{\geq 1}$ and A is some $\mathbb{Q}$-Cartier divisor in $\overline{X}$ without components from G or E (notice that here E is $\mathbb{Q}$-Cartier, $a \geq 1$ since $h \in \mathcal{I}_{\pi_* G}$, and $b \geq 1$ since $x \in \pi_* G$).

Applying ϕ^* to $\pi^* H$, we get

$$\phi^* \pi^* H = \phi^*(aG + bE + A) = a\tilde{G} + b\tilde{E} + (aa_1 + bc + d)F + (\text{others}) \qquad (\#)$$

where $\phi^* E = \tilde{E} + cF + (\text{others})$, $\phi^* G = \tilde{G} + a_1 F + (\text{others})$, $\phi^* A = dF + \tilde{A} + (\text{others})$, with $a_1 \in \mathbb{Z}_{\geq 1}$, $c \in \mathbb{Q}_{>0}, d \in \mathbb{Q}_{\geq 0}$. "Others" means the divisors whose supports are disjoint from the shown divisors in the corresponding equations. In addition, $\tilde{G}$, $\tilde{E}$, and $\tilde{A}$ denote the strict transforms of G, E, and A to Y.

Here we can see that

$$\mathrm{val}_E(h) = b, \qquad \mathrm{val}_F(h) = aa_1 + bc + d.$$

In other words,

$$\mathrm{val}_F(h) = aa_1 + c\mathrm{val}_E(h) + d.$$

Our goal is to find some particular $h \in \mathcal{I}_{\pi_* G,p}$ such that $\mathrm{val}_E(h) < \mathrm{val}_F(h)$.To obtain this inequality, it suffices to construct an $h \in \mathcal{I}_{\pi_* G,p}$ satisfying either (1) $b = 1$ or (2) $c \geq 1$.

Remark 7.2.2. In the following proof, we will construct an $h \in \mathcal{I}_{\pi_* G,p}$ such that $b = 1$.

7.3 Weighted Blowing-Ups

In [Hay99, Hay00], we know that any divisorial contraction with minimal discrepancy is obtained by a *weighted blowing-up* with some weight $\sigma = \frac{1}{m}(a,b,c,d) \in \frac{1}{m}(\mathbb{Z}_{>0})^4$. In this section, we introduce the notation and tools that will be used in the proof of our main theorem in this chapter. The details can be found in [Hay99, Hay00, Che16, Che23]

The cyclic quotient singularity $Y = \mathbb{C}^4/\frac{1}{m}(\alpha_1,\alpha_2,\alpha_3,\alpha_4)$ is the toric variety corresponding to the lattice $\overline{N} := \mathbb{Z}e_1 + \mathbb{Z}e_2 + \mathbb{Z}e_3 + \mathbb{Z}e_4 + \mathbb{Z}e = N + \mathbb{Z}e$ and the cone $C = \mathbb{R}_{\geq 0}e_1 + \mathbb{R}_{\geq 0}e_2 + \mathbb{R}_{\geq 0}e_3 + \mathbb{R}_{\geq 0}e_4$, where e_i, $i = 1,2,3,4$, are the standard basis vectors for $\mathbb{Z}^4$, and $e = \frac{1}{m}(\alpha_1,\alpha_2,\alpha_3,\alpha_4) \in \frac{1}{m}\mathbb{Z}^4$. $\overline{N}$ and its fan Δ consist of all faces of C. In other words,

$$Y = \mathbb{C}^4/\frac{1}{m}(\alpha_1,\alpha_2,\alpha_3,\alpha_4) \cong \mathrm{Spec}(\mathbb{C}[C^\vee \cap \overline{M}])$$

where $\overline{M} := \mathrm{Hom}_{\mathbb{Z}}(\overline{N},\mathbb{Z})$.

Let us give a description of weighted blowing-ups for $\mathbb{C}^4/\frac{1}{m}(\alpha_1,\alpha_2,\alpha_3,\alpha_4)$. Let $\sigma = \frac{1}{m}(a,b,c,d) \in \overline{N}$ with $a,b,c,d > 0$, and assume e_1,e_2,e_3,e_4,σ generate the lattice $\overline{N}$. This σ will be called a *weight*.

We construct the *weighted blowing-up* $\bar{\pi}_\sigma : \overline{Y} \to Y = \mathbb{C}^4/\frac{1}{m}(\alpha_1,\alpha_2,\alpha_3,\alpha_4)$ with weight σ as follows.

We divide the cone C by adding 1-dimensional cone $\mathbb{R}_{\geq 0}\sigma$, and we get a subdivision of C into 4 cones:

$$C_1 = \mathbb{R}_{\geq 0}\sigma + \mathbb{R}_{\geq 0}e_2 + \mathbb{R}_{\geq 0}e_3 + \mathbb{R}_{\geq 0}e_4,$$

$$C_2 = \mathbb{R}_{\geq 0}e_1 + \mathbb{R}_{\geq 0}\sigma + \mathbb{R}_{\geq 0}e_3 + \mathbb{R}_{\geq 0}e_4,$$

$$C_3 = \mathbb{R}_{\geq 0}e_1 + \mathbb{R}_{\geq 0}e_2 + \mathbb{R}_{\geq 0}\sigma + \mathbb{R}_{\geq 0}e_4,$$

$$C_4 = \mathbb{R}_{\geq 0}e_1 + \mathbb{R}_{\geq 0}e_2 + \mathbb{R}_{\geq 0}e_3 + \mathbb{R}_{\geq 0}\sigma.$$

Let Δ' be the fan consisting of all faces of C_1, C_2, C_3, C_4. Then the *weighted blowing-up* $\overline{Y}$ of Y is defined as the toric variety corresponding to the lattice $\overline{N}$ and the fan Δ', and $\bar{\pi}_\sigma : \overline{Y} \to Y$ is the morphsim induced from the natural map of fans $(\overline{N}, \Delta') \to (\overline{N}, \Delta)$.

The toric variety $\overline{Y}$ is covered by 4 affine open sets $\overline{U}_1, \overline{U}_2, \overline{U}_3, \overline{U}_4$ corresponding to the cones C_1, C_2, C_3, C_4, respectively. These affine open sets are as follows:

$$\begin{aligned}
\overline{U}_1 &= \mathbb{C}^4/\frac{1}{a}(m, -b, -c, -d) \cong \mathrm{Spec}(\mathbb{C}[C_1^\vee \cap \overline{M}]),\\
\overline{U}_2 &= \mathbb{C}^4/\frac{1}{b}(-a, m, -c, -d) \cong \mathrm{Spec}(\mathbb{C}[C_2^\vee \cap \overline{M}]),\\
\overline{U}_3 &= \mathbb{C}^4/\frac{1}{c}(-a, -b, m, -d) \cong \mathrm{Spec}(\mathbb{C}[C_3^\vee \cap \overline{M}]),\\
\overline{U}_4 &= \mathbb{C}^4/\frac{1}{d}(-a, -b, -c, m) \cong \mathrm{Spec}(\mathbb{C}[C_4^\vee \cap \overline{M}]).
\end{aligned}$$

And $\bar{\pi}_\sigma$ are formulated as follows:

$$\begin{aligned}
\bar{\pi}_\sigma|_{\overline{U}_1} &: \overline{U}_1 \longrightarrow Y \qquad (x, y, z, u) \mapsto (x^{\frac{a}{m}}, x^{\frac{b}{m}}y, x^{\frac{c}{m}}z, x^{\frac{d}{m}}u),\\
\bar{\pi}_\sigma|_{\overline{U}_2} &: \overline{U}_2 \longrightarrow Y \qquad (x, y, z, u) \mapsto (y^{\frac{a}{m}}x, y^{\frac{b}{m}}, y^{\frac{c}{m}}z, y^{\frac{d}{m}}u),\\
\bar{\pi}_\sigma|_{\overline{U}_3} &: \overline{U}_3 \longrightarrow Y \qquad (x, y, z, u) \mapsto (z^{\frac{a}{m}}x, z^{\frac{b}{m}}y, z^{\frac{c}{m}}, z^{\frac{d}{m}}u),\\
\bar{\pi}_\sigma|_{\overline{U}_4} &: \overline{U}_4 \longrightarrow Y \qquad (x, y, z, u) \mapsto (u^{\frac{a}{m}}x, u^{\frac{b}{m}}y, u^{\frac{c}{m}}z, u^{\frac{d}{m}}).
\end{aligned}$$

In these expressions, x, y, z, u denote coordinates in the ambient $\mathbb{C}^4$. For example, the map $\bar{\pi}_\sigma|_{\overline{U}_1}$ is determined by a map $\mathbb{C}^4 \longrightarrow \mathbb{C}^4$ which, at the level of coordinate rings, is given by

$$\pi_\sigma^\#(x) = x^{\frac{a}{m}}, \quad \pi_\sigma^\#(y) = x^{\frac{b}{m}}y, \quad \pi_\sigma^\#(z) = x^{\frac{c}{m}}z, \quad \pi_\sigma^\#(u) = x^{\frac{d}{m}}u.$$

The exceptional divisor $\overline{E}$ of $\bar{\pi}_\sigma$ is isomorphic to the weighted projective space $\mathbb{P}_{\mathbb{C}}(a, b, c, d)$.

Let us formulate weighted blowing-ups for a subvariety defined by a semi-invariant in $\mathbb{C}^4/\frac{1}{m}(\alpha_1, \alpha_2, \alpha_3, \alpha_4)$. Let $\phi \in \mathbb{C}\{x, y, z, u\}$ be a $\mathbb{Z}_m$-semi-invariant, and

it defines a hyperquotient singularity $X = (\phi = 0) \subseteq \mathbb{C}^4/\frac{1}{m}(\alpha_1, \alpha_2, \alpha_3, \alpha_4)$. Let $\overline{X} = (\bar{\pi}_\sigma)^{-1}(X)$ be the strict transform of X via $\bar{\pi}_\sigma$ and let $\pi_\sigma = \bar{\pi}_\sigma|_{\overline{X}}$ be the restriction of $\bar{\pi}_\sigma$ in $\overline{X}$. Then $\pi_\sigma : \overline{X} \to X$ us called the *weighted blowing-up with weight* $\sigma = \frac{1}{m}(a, b, c, d)$ (or briefly σ-blowing-up). Let $U_i := \overline{U}_i \cap \overline{X}$ for $i = 1, 2, 3, 4$, then each U_i is a hyperquotient singularity in $\overline{U}_i$ and $\overline{X}$ is covered by 4 charts U_1, U_2, U_3, U_4. We denote $E := \overline{E} \cap \overline{X} \subseteq \mathbb{P}_{\mathbb{C}}(a, b, c, d)$, and notice that E is not necessarily irreducible in $\overline{X}$.

We can explicitly express the defining equations of $\overline{X} \cap U_i$ for $i = 1, 2, 3, 4$. Write a semi-invariant $\phi = \sum_{i,j,k,\ell} a_{i,j,k,\ell} x^{\alpha_i} y^{\beta_j} z^{\gamma_k} u^{\delta_\ell}$, and we define

$$\mathrm{wt}_\sigma(\phi) := \min\{ \frac{a\alpha_i + b\beta_j + c\gamma_k + d\delta_\ell}{m} \mid a_{i,j,k,\ell} \neq 0 \}$$

for a given weight $\sigma = \frac{1}{m}(a, b, c, d) \in \overline{N}$. Notice that $\overline{X} \cap U_i$ for $i = 1, 2, 3, 4$ is defined by

$$\begin{aligned}
\phi_1 &:= \phi(x^{\frac{a}{m}}, x^{\frac{b}{m}} y, x^{\frac{c}{m}} z, x^{\frac{d}{m}} u) x^{-\mathrm{wt}_\sigma(\phi)} \text{ in } U_1, \\
\phi_2 &:= \phi(y^{\frac{a}{m}} x, y^{\frac{b}{m}}, y^{\frac{c}{m}} z, y^{\frac{d}{m}} u) y^{-\mathrm{wt}_\sigma(\phi)} \text{ in } U_2, \\
\phi_3 &:= \phi(z^{\frac{a}{m}} x, z^{\frac{b}{m}} y, z^{\frac{c}{m}}, z^{\frac{d}{m}} u) z^{-\mathrm{wt}_\sigma(\phi)} \text{ in } U_3, \\
\phi_4 &:= \phi(u^{\frac{a}{m}} x, u^{\frac{b}{m}} y, u^{\frac{c}{m}} z, u^{\frac{d}{m}}) u^{-\mathrm{wt}_\sigma(\phi)} \text{ in } U_4.
\end{aligned}$$

Let us characterize the singular locus of $\overline{X}$. Write

$$(\overline{X})_{\mathrm{sing}} = [(\overline{X})_{\mathrm{sing}}]_{\mathrm{ind}=1} \bigcup [(\overline{X})_{\mathrm{sing}}]_{\mathrm{ind}>1}$$

where $[(\overline{X})_{\mathrm{sing}}]_{\mathrm{ind}=1}$ ($[(\overline{X})_{\mathrm{sing}}]_{\mathrm{ind}>1}$, resp.) denotes the singular locus with Gorenstein index $= 1$ (Gorenstein index > 1, resp.). Then we can interpret $(\overline{X})_{\mathrm{sing}}$ as the following.

Proposition 7.3.1 (2.1 in [Che16]). *Keep the notation in this section. Let $\pi_\sigma : \overline{X} \to X$ be a weighted blowing-up of X with respect to the weight σ, then we have*

(1) $[(\overline{X})_{\text{sing}}]_{\text{ind}>1} = \overline{X} \cap (\mathbb{P}_{\mathbb{C}}(a,b,c,d))_{\text{sing}} = E \cap (\mathbb{P}_{\mathbb{C}}(a,b,c,d))_{\text{sing}}$,

(2) If $\overline{U}_i \cong \mathbb{C}^4$ *for some* $i \in \{1,2,3,4\}$, *then* $(\overline{X})_{\text{sing}} \cap U_i \subseteq E_{\text{sing}} \cap U_i$,

(3) If $\overline{X}$ *is a terminal 3-fold, then* $[(\overline{X})_{\text{sing}}]_{\text{ind}=1} \subseteq E_{\text{sing}}$.

7.4 Proof of Theorem 7.0.1

We follow the notation of Section 7.2. Assume val_E is non-Nash over (X,p), so there is a divisor F over (X,p) inducing a Nash valuation val_F such that

$$\text{val}_F <_X \text{val}_E,$$

and F must be centered at some singular point of $\overline{X}$. It suffices to show that for any $q \in \overline{X}_{\text{sing}}$ such that F is centered at q, there is $f \in \mathcal{O}_{X,p}$ such $\text{val}_F(f) > \text{val}_E(f)$, which contradicts $\text{val}_F <_X \text{val}_E$, so we can conclude that val_E is Nash over (X,p).

We follow the arguments in [Che16, Proposition 18] and [Hay05, 5.2]. Recall that the standard form of this type of 3-fold terminal singularity (X,p) is

$$X = (x^2 + y^2 + f(z,u) = 0) \subseteq \mathbb{C}^4/\frac{1}{2}(0,1,1,1)$$

where $f(z,u) \in (z,u)^4 \subseteq \mathbb{C}\{z,u\}$ is a $\mathbb{Z}_2$-invariant with respect to $\frac{1}{2}(0,1,1,1)$, and $p = (0,0,0,0) \in X$. Define

$$\tau_0 := \min\{\, i+j \mid z^i u^j \in \text{supp}(f(z,u)) \text{ as a monomial } \} \geq 4.$$

Notice that $f(z,u)$ is a $\mathbb{Z}_2$-invariant with respect to $\frac{1}{2}(0,1,1,1)$, so this implies $\tau_0 \in \mathbb{Z}_{>0}$ is even.

Let $f_{\tau_0}(z,u) = \sum_{i+j=\tau_0} a_{ij} z^i u^j$, and express it in the form of the following factorization:

$$f_{\tau_0}(z,u) = \sum_{i+j=\tau_0} a_{ij} z^i u^j = \prod_{t \in T} (a_t z + b_t u)^{m_t}$$

where $\sum_{t \in T} m_t = \tau_0$.

Case 1: $f_{\tau_0}(z,u)$ is not a perfect square

From [Hay99, 8.2], consider the weighted blow-up with the following weight

$$\sigma := \begin{cases} \frac{1}{2}(\frac{\tau_0}{2}, \frac{\tau_0}{2}+1, 1, 1), & \text{if } \frac{\tau_0}{2} \text{ is even,} \\ \frac{1}{2}(\frac{\tau_0}{2}+1, \frac{\tau_0}{2}, 1, 1), & \text{if } \frac{\tau_0}{2} \text{ is odd.} \end{cases}$$

We consider the first case $\sigma = \frac{1}{2}(\frac{\tau_0}{2}, \frac{\tau_0}{2}+1, 1, 1)$, and the second case is followed by an analogous argument. The weighted blowing-up with respect to σ induces a divisorial contraction $\pi_\sigma : \overline{X} \to X$ with the exceptional prime divisor $E = \pi_\sigma^{-1}(p) \subseteq \overline{X}$ with minimal discrepancy (in this case $a(E,X) = \frac{1}{2}$) with respect to (X,p) by Theorem 8.4 in [Hay99]. Notice that $\overline{X}$ is covered by four affine open sets U_1, U_2, U_3, and U_4. By [Che16, Proposition 18] and [Hay05, 5.2], we have that $\overline{X}$ is non-singular on U_1, it has one terminal quotient singularity of index $\frac{\tau_0}{2}+1$ in U_2, denoted by Q_2, and several Gorenstein terminal singularities of type cD in U_3 and U_4.

In U_2, there is a terminal quotient singularity $Q_2 = (0,0,0,0) \in U_2$ with index $\frac{\tau_0}{2}+1$ (see [Che16, Proposition 18 Case 1] and [Hay05, 5.2]). We consider $g = x^2 y^{\frac{\tau_0}{2}-1} + y^{\frac{\tau_0}{2}} + z^2 + u^2 \in \mathcal{O}_{\overline{X},Q_2} \setminus \mathcal{I}_{E,Q_2}$ ($g \notin \mathcal{I}_{E,Q_2}$ since $E = (x^2 + f_{\tau_0}(z,u) = 0)$ in U_2), and let $G = \operatorname{div}_{\overline{X}}(g) \subseteq \overline{X}$. Consider $h = x^2 + y^2 + z^2 + u^2 \in \mathcal{O}_{X,p}$, and let $H = \operatorname{div}_X(h)$.

Here we compute $\pi_\sigma^* H$, the pullback of H via π_σ, in U_2. Notice that π_σ is the weighted blowing-up with respect to σ given by

$$(\pi_\sigma^\#(x), \pi_\sigma^\#(y), \pi_\sigma^\#(z), \pi_\sigma^\#(u)) = (xy^{\frac{\tau_0}{4}}, y^{\frac{\tau_0}{4}+\frac{1}{2}}, zy^{\frac{1}{2}}, uy^{\frac{1}{2}}).$$

Hence,

$$\begin{aligned} \pi_\sigma^\#(h) &= \pi_\sigma^\#(x^2+y^2+z^2+u^2) = x^2 y^{\frac{\tau_0}{2}} + y^{\frac{\tau_0}{2}+1} + z^2 y + u^2 y \\ &= y(x^2 y^{\frac{\tau_0}{2}-1} + y^{\frac{\tau_0}{2}} + z^2 + u^2) = yg. \end{aligned}$$

In other words, we obtain

$$\pi_\sigma^* H = E + G.$$

Notice that $(\pi_\sigma)_* G = H$, so we have $h = x^2 + y^2 + z^2 + u^2 \in \mathcal{I}_{(\pi_\sigma)_* G, p}$. Apply the equation (#) to π_σ, we have $b = \mathrm{val}_E(h) = 1$, so we have $\mathrm{val}_E(h) < \mathrm{val}_F(h)$ for any divisor F over X centered at $Q_2 \in \overline{X}$.

We compute the strict transform of the defining equation of $\overline{X}$ in U_3, and it is given by

$$(\pi_\sigma^\#(x), \pi_\sigma^\#(y), \pi_\sigma^\#(z), \pi_\sigma^\#(u)) = (xz^{\frac{\tau_0}{4}}, yz^{\frac{\tau_0}{4}+\frac{1}{2}}, z^{\frac{1}{2}}, uz^{\frac{1}{2}}).$$

Hence, in U_3,

$$\begin{aligned}
\pi_\sigma^\#(x^2 + y^2 + f(z,u)) &= \pi_\sigma^\#(x^2 + y^2 + f_{\tau_0}(z,u) + f_{>\tau_0}(z,u)) \\
&= x^2 z^{\frac{\tau_0}{2}} + y^2 z^{\frac{\tau_0}{2}+1} + f_{\tau_0}(z^{\frac{1}{2}}, uz^{\frac{1}{2}}) + f_{>\tau_0}(z^{\frac{1}{2}}, uz^{\frac{1}{2}}) \\
&= x^2 z^{\frac{\tau_0}{2}} + y^2 z^{\frac{\tau_0}{2}+1} + \prod_{t\in T}(a_t z^{\frac{1}{2}} + b_t u z^{\frac{1}{2}})^{m_t} + f_{>\tau_0}(z^{\frac{1}{2}}, uz^{\frac{1}{2}}) \\
&= z^{\frac{\tau_0}{2}}(x^2 + y^2 z + \prod_{t\in T}(a_t + b_t u)^{m_t} + \frac{f_{>\tau_0}(z^{\frac{1}{2}}, uz^{\frac{1}{2}})}{z^{\frac{\tau_0}{2}}}).
\end{aligned}$$

We obtain the strict transform of $X = (x^2 + y^2 + f(z,u) = 0) \subseteq \mathbb{C}/\frac{1}{2}(0,1,1,1)$ in U_3 is $(x^2 + y^2 z + \prod_{t\in T}(a_t + b_t u)^{m_t} + \frac{f_{>\tau_0}(z^{\frac{1}{2}}, uz^{\frac{1}{2}})}{z^{\frac{\tau_0}{2}}} = 0) \subseteq \mathbb{C}^4$, which contains some Gorenstein terminal singularities of type cD.

Here we compute $(\overline{X})_{\mathrm{sing}}$ in U_3 by applying the Jacobian criterion to $(x^2 + y^2 z + \prod_{t\in T}(a_t + b_t u)^{m_t} + \frac{f_{>\tau_0}(z^{\frac{1}{2}}, uz^{\frac{1}{2}})}{z^{\frac{\tau_0}{2}}} = 0) \subseteq \mathbb{C}^4$. Let $F = x^2 + y^2 z + \prod_{t\in T}(a_t + b_t u)^{m_t} + \frac{f_{>\tau_0}(z^{\frac{1}{2}}, uz^{\frac{1}{2}})}{z^{\frac{\tau_0}{2}}}$, and we solve the equation $\mathrm{grad}(F) = (0,0,0,0)$ for (x,y,z,u).

$$\begin{aligned}
2x &= 0, \\
2yz &= 0, \\
y^2 + \frac{\partial}{\partial z}(\frac{f_{>\tau_0}(z^{\frac{1}{2}}, uz^{\frac{1}{2}})}{z^{\frac{\tau_0}{2}}}) &= 0, \\
\frac{\partial}{\partial u}(\prod_{t\in T}(a_t + b_t u)^{m_t} + \frac{f_{>\tau_0}(z^{\frac{1}{2}}, uz^{\frac{1}{2}})}{z^{\frac{\tau_0}{2}}}) &= 0.
\end{aligned}$$

Notice that we have $z = 0$ in U_3 because $(\overline{X})_{\mathrm{sing}} \cap U_3 \subseteq E_{\mathrm{sing}} \cap U_3$ by Proposition 7.3.1.

Some direct calculations show that $(\overline{X})_{\text{sing}}$ in U_3 is $\{(0, \beta_t, 0, -\frac{a_t}{b_t})\}_{t\in T} \subseteq \mathbb{C}^4$ for some $\beta_t \in \mathbb{C}$.

Consider $h = x^2+y^2-(\frac{a_t}{b_t})^2z^2+u^2$ ($H = \text{div}_X(h)$), and $g = x^2z^{\frac{\tau_0}{2}-1}+y^2z^{\tau_0}+u^2-(\frac{a_t}{b_t})^2$ ($G = \text{div}_{\overline{X}}(g)$). We have

$$\begin{aligned}\pi_\sigma^{\#}(h) &= \pi_\sigma^{\#}(x^2+y^2-(\frac{a_t}{b_t})^2z^2+u^2) = x^2z^{\frac{\tau_0}{2}}+y^2z^{\frac{\tau_0}{2}+1}-(\frac{a_t}{b_t})^2z+u^2z)\\ &= z(x^2z^{\frac{\tau_0}{2}-1}+y^2z^{\tau_0}-(\frac{a_t}{b_t})^2+u^2) = zg.\end{aligned}$$

Hence, we have

$$\pi_\sigma^* H = E + G.$$

Applying equation (#) to π_σ, we have $b = \text{val}_E(h) = 1$, so we have $\text{val}_E(h) < \text{val}_F(h)$ for any divisor F over X centered at any point in $\{(0, \beta_t, 0, -\frac{a_t}{b_t})\}_{t\in T}$.

In U_4, we have an argument similar to the case in U_3. Therefore, val_E induces a Nash valuation of X.

Case 2: $f_{\tau_0}(z,u)$ is a perfect square

If $f_{\tau_0}(z,u)$ is a perfect square, say $f_{\tau_0}(z,u) = (p_{\frac{\tau_0}{2}}(z,u))^2$, we apply the changes of coordinates in [Hay99, 8.6]:

$$x \mapsto x \pm p_{\frac{\tau_0}{2}}(z,u), \quad y \mapsto y, \quad z \mapsto z, \quad u \mapsto u.$$

and (X, p) is rewritten as

$$(x^2 \pm 2xp_{\frac{\tau_0}{2}}(z,u) + y^2 + f_{\geq\tau_0+1}(z,u) = 0) \subseteq \mathbb{C}^4/\frac{1}{2}(0,1,1,1).$$

By [Hay99, Theorem 8.8], we obtain two non-isomorphic divisorial contractions with minimal discrepancy (in this case the minimal discrepancy is $\frac{1}{2}$). We deal with case $X = (x^2 + 2xp_{\frac{\tau_0}{2}}(z,u) + y^2 + f_{\geq\tau_0+1}(z,u) = 0)$ here, and case $X = (x^2 - 2xp_{\frac{\tau_0}{2}}(z,u) + y^2 + f_{\geq\tau_0+1}(z,u) = 0)$ is similar.

From [Hay99, 8.2], consider the weighted blowing-up with respect to the following weight.

$$\sigma := \begin{cases} \frac{1}{2}(\frac{\tau_0}{2}+2, \frac{\tau_0}{2}+1, 1, 1), & \text{if } \frac{\tau_0}{2} \text{ is even,} \\ \frac{1}{2}(\frac{\tau_0}{2}+1, \frac{\tau_0}{2}+2, 1, 1), & \text{if } \frac{\tau_0}{2} \text{ is odd.} \end{cases}$$

We consider the first case $\sigma = \frac{1}{2}(\frac{\tau_0}{2}+2, \frac{\tau_0}{2}+1, 1, 1)$, and the second case $\frac{1}{2}(\frac{\tau_0}{2}+1, \frac{\tau_0}{2}+2, 1, 1)$ will follow to an analogous argument. The weighted blowing-up with respect to σ induces a divisorial contraction $\pi_\sigma : \overline{X} \to X$ with the exceptional prime divisor $E = \pi_\sigma^{-1}(p) \subseteq \overline{X}$ with minimal discrepancy (in this case $a(E, X) = \frac{1}{2}$) of (X, p) by [Hay99, Theorem 8.7].

$\overline{X}$ is non-singular on U_2. In U_1, we have that $\overline{X}$ is singular at $Q_1 \in U_1$ (see [Che16, Proposition 18 Case 2] and [Hay05, 5.2]), which is a terminal quotient singularity of the Gorenstein index $\frac{\tau_0}{2}+2$. We consider $g = x^{\frac{\tau_0}{2}+1} + y^2x^{\frac{\tau_0}{2}} + z^2 + u^2$, and let $G = \mathrm{div}_{\overline{X}}(g) \subseteq \overline{X}$. Consider $h = x^2 + y^2 + z^2 + u^2$, and let $H = \mathrm{div}_X(h) \subseteq X$.

We compute $\pi_\sigma^* H$, and it is given by

$$(\pi_\sigma^\#(x), \pi_\sigma^\#(y), \pi_\sigma^\#(z), \pi_\sigma^\#(u)) = (x^{\frac{\tau_0}{4}+1}, yx^{\frac{\tau_0}{4}+\frac{1}{2}}, zx^{\frac{1}{2}}, ux^{\frac{1}{2}}).$$

Hence, we have

$$\begin{aligned} \pi_\sigma^\#(h) = \pi_\sigma^\#(x^2 + y^2 + z^2 + u^2) &= x^{\frac{\tau_0}{2}+2} + y^2x^{\frac{\tau_0}{2}+1} + z^2x + u^2x \\ &= x(x^{\frac{\tau_0}{2}+1} + y^2x^{\frac{\tau_0}{2}} + z^2 + u^2) = xg. \end{aligned}$$

We obtain

$$\pi_\sigma^* H = E + G.$$

Notice that $(\pi_\sigma)_*(G) = H$ (G is the strict transform of H), so we apply the equation (#) to π_σ and we have $b = \mathrm{val}_E(h) = 1$, so we have $\mathrm{val}_E(h) < \mathrm{val}_F(h)$ for any divisor F over X centered at $Q_1 \in U_1 \subseteq \overline{X}$.

We compute the strict transform of the defining equation of $\overline{X}$ in U_3, and it is given by

$$(\pi_\sigma^\#(x), \pi_\sigma^\#(y), \pi_\sigma^\#(z), \pi_\sigma^\#(u)) = (xz^{\frac{\tau_0}{4}+1}, yz^{\frac{\tau_0}{4}+\frac{1}{2}}, z^{\frac{1}{2}}, uz^{\frac{1}{2}}).$$

Hence, in U_3,

$$\begin{aligned}
&\pi_\sigma^\#(x^2 + 2xp_{\frac{\tau_0}{2}}(z,u) + y^2 + f_{>\tau_0+1}(z,u)) \\
&\quad = x^2 z^{\frac{\tau_0}{2}+2} + 2xz^{\frac{\tau_0}{4}+1} p_{\frac{\tau_0}{2}}(z^{\frac{1}{2}}, uz^{\frac{1}{2}}) + y^2 z^{\frac{\tau_0}{2}+1} + f_{>\tau_0+1}(z^{\frac{1}{2}}, uz^{\frac{1}{2}}) \\
&\quad = z^{\frac{\tau_0}{2}+1}(x^2 z + 2xp_{\frac{\tau_0}{2}}(1,u) + y^2 + \frac{f_{>\tau_0+1}(z^{\frac{1}{2}}, uz^{\frac{1}{2}})}{z^{\frac{\tau_0}{2}+1}}).
\end{aligned}$$

We obtain that the strict transform of the defining equation $(x^2 + y^2 + f(z,u) = 0) \subseteq \mathbb{C}/\frac{1}{2}(0,1,1,1)$ in U_3 is $(x^2 z + 2xp_{\frac{\tau_0}{2}}(1,u) + y^2 + \frac{f_{>\tau_0+1}(z^{\frac{1}{2}}, uz^{\frac{1}{2}})}{z^{\frac{\tau_0}{2}+1}} = 0) \subseteq \mathbb{C}^4$, which contains Gorenstein terminal singularities at worst type cD (it means either non-singular or of singular of type cD).

Here we compute $(\overline{X})_{\text{sing}}$ in U_3 by applying the Jacobian criterion to $(x^2 z + 2xp_{\frac{\tau_0}{2}}(1,u) + y^2 + \frac{f_{>\tau_0+1}(z^{\frac{1}{2}}, uz^{\frac{1}{2}})}{z^{\frac{\tau_0}{2}+1}} = 0) \subseteq \mathbb{C}^4$. Let $F = x^2 z + 2xp_{\frac{\tau_0}{2}}(1,u) + y^2 + \frac{f_{>\tau_0+1}(z^{\frac{1}{2}}, uz^{\frac{1}{2}})}{z^{\frac{\tau_0}{2}+1}}$, and we solve equation $\operatorname{grad}(F) = (0,0,0,0)$ for (x,y,z,u).

$$\begin{aligned}
2xz + 2p_{\frac{\tau_0}{2}}(1,u) &= 0, \\
2y &= 0, \\
x^2 + \frac{\partial}{\partial z}(\frac{f_{>\tau_0+1}(z^{\frac{1}{2}}, uz^{\frac{1}{2}})}{z^{\frac{\tau_0}{2}+1}}) &= 0, \\
\frac{\partial}{\partial u}(2xp_{\frac{\tau_0}{2}}(1,u) + \frac{f_{>\tau_0+1}(z^{\frac{1}{2}}, uz^{\frac{1}{2}})}{z^{\frac{\tau_0}{2}+1}}) &= 0.
\end{aligned}$$

Notice that we have $z = 0$ in U_3 because $(\overline{X})_{\text{sing}} \cap U_3 \subseteq E_{\text{sing}} \cap U_3$ by Proposition 7.3.1. Some direct calculations show that $(\overline{X})_{\text{sing}}$ in U_3 is $\{(a_t, 0, 0, b_t)\}_{t\in T}$ for some $a_t, b_t \in \mathbb{C}$. Consider $h = x^2 + y^2 - b_t^2 z^2 + u^2$ ($H = \operatorname{div}_X(h) \subseteq X$), and $g = x^2 z^{\frac{\tau_0}{2}-1} + y^2 z^{\tau_0} +$

$u^2 - b_t^2 \in \mathcal{O}_{\overline{X},(a_t,0,0,b_t)} \setminus \mathcal{I}_{E,(a_t,0,0,b_t)}$ ($G = \mathrm{div}_{\overline{X}}(g) \subseteq \overline{X}$). We have

$$\begin{aligned}\pi_\sigma^{\#}(h) &= \pi_\sigma^{\#}\left(x^2 + y^2 - b_t^2 z^2 + u^2\right) = x^2 z^{\frac{\tau_0}{2}} + y^2 z^{\frac{\tau_0}{2}+1} - b_t^2 z + u^2 z \\ &= z\left(x^2 z^{\frac{\tau_0}{2}-1} + y^2 z^{\tau_0} - b_t^2 + u^2\right) = zg.\end{aligned}$$

Hence we have

$$\pi_\sigma^* H = E + G.$$

Notice that $(\pi_\sigma)_*(G) = H$ (G is the strict transform of H), so we apply the equation (#) to π_σ and we have $b = \mathrm{val}_E(h) = 1$, so we have $\mathrm{val}_E(h) < \mathrm{val}_F(h)$ for any divisor F over X centered at any point in $\{(a_t, 0, 0, b_t)\}_{t\in T}$.

In U_4, we have an argument similar to the case in U_3. Therefore, val_E induces a Nash valuation.

Therefore, we prove that any exceptional prime divisor E with minimal discrepancy (here $a(E, X) = \frac{1}{2}$) induces a Nash valuation val_E over a 3-fold terminal singularity of type cAx/2.

Chapter 8

Nash Valuations over 3-Fold Gorenstein Terminal Singularities

Let (X, p) be a germ of 3-fold Gorenstein terminal singularity. In this final chapter, we establish a partial result in support of Conjectures A and B, and provide some examples as evidence (notice that Conjectures A and B are the same because the minimal discrepancy of a 3-fold Gorenstein terminal singularity is 1 by [Mar96, Theorem 0.1]).

8.1 Main Theorem

We follow the set-up developed in 7.2, then the following result follows directly.

Proposition 8.1.1. *Let (X, p) be a 3-fold Gorenstein terminal singularity, let $\pi : \overline{X} \to X$ be a divisorial contraction with minimal discrepancy (it is 1 in this case), and let $E = \pi^{-1}(p)$ be its exceptional prime divisor. Assume that the singularities of $\overline{X}$ are terminal cyclic quotient or Gorenstein terminal. Then there is an exceptional prime divisor over X*

with discrepancy 1 *that induces a Nash valuation. In other words,*

$$\{F \text{ exceptional divisor over } X \mid a(F,X) = 1 \text{ and } \mathrm{val}_F \text{ is a Nash valuation of } X\} \neq \varnothing.$$

Proof. If val_E is Nash for X, then we are done. If not, then there is an exceptional prime divisor F over X such that val_F is Nash for X, and $\mathrm{val}_F <_X \mathrm{val}_E$ by Lemma 7.2.1.

Let $\phi : Y \to \overline{X}$ be a resolution (notice that F appears in Y since val_F is Nash, which is essential for X). Consider $\phi^* E = \tilde{E} + cF + \sum c_i F_i$ where $\tilde{E}$ denotes the strict transform of E via ϕ. By direct computations, we have

$$a(F,X) = a(F,\overline{X}) + c.$$

Notice that $a(F,X) \in \mathbb{Z}_{>0}$ since (X,p) is Gorenstein.

We claim that this valuation val_F has $a(F,X) = 1$. Since val_F is Nash for X (so val_F is essential for X), F must be centered at some terminal cyclic quotient singularity or some Gorenstein terminal singularity in $\overline{X}$.

If F is centered at any Gorenstein point $q \in \overline{X}$, we have $a(F,\overline{X}) \in \mathbb{Z}_{>0}$. This implies $c = a(F,X) - a(E,\overline{X}) \in \mathbb{Z}_{>0}$ (notice that $c \in \mathbb{Q}_{>0}$ since F is centered at $q \in \overline{X}$). Applying a similar argument as 7.2 to this scenario, we obtain

$$\mathrm{val}_F(h) = aa_1 + c\mathrm{val}_E(h) + d$$

for some $h \in \mathcal{O}_{X,p}$. Since $c \geq 1$, we get $\mathrm{val}_E(h) < \mathrm{val}_F(h)$, contradicting the fact $\mathrm{val}_E >_X \mathrm{val}_F$. Hence, F must be centered at some terminal cyclic quotient singularity in $\overline{X}$. We have

$$0 < a(F,\overline{X}) < 1$$

because this singularity has a resolution such that all exceptional prime divisors have discrepancies less than 1 (see [Hay99, Proposition 5.1], and the corresponding

morphism is called *economic resolution*). In this case, we must have $a(F, X) = 1$ because $a(F, X) \geq 2$ implies

$$c = a(E, X) - a(E, \overline{X}) > 2 - 1 = 1,$$

which also provides a $h \in \mathcal{O}_{X,p}$ such that $\mathrm{val}_E(h) < \mathrm{val}_F(h)$. Therefore, there is an exceptional prime divisor over (X, p) with discrepancy equal to 1 such that it induces a Nash valuation over X. □

8.2 Examples

Here we select some examples of 3-fold Gorenstein terminal singularities that satisfy the assumptions of the proposition 8.1.1.

Define $\tau(f) := \min\{j + k \mid z^j u^k \in \mathrm{supp}(f(z, u)) \text{ as a monimial } \}$ for $f \in \mathbb{C}\{z, u\}$. Recall for a Gorenstein terminal singularity (X, p), it is called *type* cE is it has the following expression.

$$(x^2 + y^3 + yg(z, u) + h(z, u) = 0) \subseteq \mathbb{C}^4.$$

for some $g, h \in \mathbb{C}\{z, u\}$ with $\tau(g) \geq 3$ and $\tau(h) \geq 4$.

Furthermore,

(1) it is called type cE_6 if $\tau(h) = 4$ and $\tau(g) \geq 3$.

(2) it is called type cE_7 if $\tau(h) \geq 5$ and $\tau(g) = 3$.

(3) it is called type cE_8 if $\tau(h) = 5$ and $\tau(g) \geq 4$.

Example 8.2.1 ([Che15, Che16]). Let (X, p) be the Gorenstein terminal singularity of type cE_6 which satisfies the conditions in [Che16, Subcase 1-2 of Theorem 34]. After applying the change of variables mentioned in [Che16, Subcase 1-2 of Theorem 34] for (X, p), consider the weighted blowing-up $\pi_\sigma : \overline{X} \to X$ with the weight

$\sigma = (3,2,1,1)$. By the argument in [Che16, Lemma 7 and Theorem 5], π_σ is a divisorial contraction with minimal discrepancy to X (in this case it is 1), and we obtain $(\overline{X})_{\text{sing}} = [(\overline{X})_{\text{sing}}]_{\text{ind}>1} = \{Q_1, Q_2\}$ where Q_1, Q_2 are terminal quotient singularities of the Gorenstein index $3, 2$, respectively. Hence, there is a Nash valuation induced by some exceptional prime divisor over (X,p) with minimal discrepancy due to Proposition 8.1.1. In fact, there are exactly two divisors over (X,p) with discrepancy 1 (see [Che15, Level 1 in 5.4]), so we know that one of them should induce a Nash valuation over (X,p).

Example 8.2.2 ([Che16]). Let (X,p) the Gorenstein terminal singularity of type cE_6 which satisfies the conditions in [Che16, Case 3 of Theorem 34]. Consider the weighted blowing-up $\pi_\sigma : \overline{X} \to X$ with the weight $\sigma = (4,3,2,1)$. By the argument in [Che16, Case 3 of Theorem 34], we have $[(\overline{X})_{\text{sing}}]_{\text{ind}=1}$ are at worst of type cE_6 and $[(\overline{X})_{\text{sing}}]_{\text{ind}>1}$ contains three cyclic quotient singularities. Hence, there is a Nash valuation induced by some exceptional prime divisor over (X,p) with minimal discrepancy due to Proposition 8.1.1.

There are other examples of 3-fold Gorenstein terminal singularities of type cE_7 and cE_8 that satisfy the assumptions in Proposition 8.1.1. See [Che16, Theorem 36 and Theorem 37].

Bibliography

[BCHM10] Caucher Birkar, Paolo Cascini, Christopher D. Hacon, and James McKernan. Existence of minimal models for varieties of log general type. *J. Amer. Math. Soc.*, 23(2):405–468, 2010.

[Bha16] Bhargav Bhatt. Algebraization and Tannaka duality. *Camb. J. Math.*, 4(4):403–461, 2016.

[BLM25] David Bourqui, Kevin Langlois, and Hussein Mourtada. The Nash problem for torus actions of complexity one. *J. Reine Angew. Math.*, 818:165–214, 2025.

[Che15] J. A. Chen. Threefold divisorial contractions to singularities of cE type. In *Recent advances in algebraic geometry*, volume 417 of *London Math. Soc. Lecture Note Ser.*, pages 102–122. Cambridge Univ. Press, Cambridge, 2015.

[Che16] Jungkai Alfred Chen. Explicit resolution of three dimensional terminal singularities. In *Minimal models and extremal rays (Kyoto, 2011)*, volume 70 of *Adv. Stud. Pure Math.*, pages 323–360. Math. Soc. Japan, [Tokyo], 2016.

[Che23] Hsin-Ku Chen. On the Nash problem for terminal threefolds of type cA/r. *Internat. J. Math.*, 34(10):Paper No. 2350055, 29, 2023.

[CLNS18] Antoine Chambert-Loir, Johannes Nicaise, and Julien Sebag. *Motivic integration*, volume 325 of *Progress in Mathematics*. Birkhäuser/Springer, New York, 2018.

[CLS11] David A. Cox, John B. Little, and Henry K. Schenck. *Toric varieties*, volume 124 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2011.

[Cut18] Steven Dale Cutkosky. *Introduction to algebraic geometry*, volume 188 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2018.

[Dan82] V. I. Danilov. Birational geometry of three-dimensional toric varieties. *Izv. Akad. Nauk SSSR Ser. Mat.*, 46(5):971–982, 1135, 1982.

[dF13] Tommaso de Fernex. Three-dimensional counter-examples to the Nash problem. *Compos. Math.*, 149(9):1519–1534, 2013.

[dF18] Tommaso de Fernex. The space of arcs of an algebraic variety. In *Algebraic geometry: Salt Lake City 2015*, volume 97.1 of *Proc. Sympos. Pure Math.*, pages 169–197. Amer. Math. Soc., Providence, RI, 2018.

[dFD16] Tommaso de Fernex and Roi Docampo. Terminal valuations and the Nash problem. *Invent. Math.*, 203(1):303–331, 2016.

[EW91] Günter Ewald and Uwe Wessels. On the ampleness of invertible sheaves in complete projective toric varieties. *Results Math.*, 19(3-4):275–278, 1991.

[FdBP12] Javier Fernández de Bobadilla and María Pe Pereira. The Nash problem for surfaces. *Ann. of Math. (2)*, 176(3):2003–2029, 2012.

[Ful89] William Fulton. *Algebraic curves*. Advanced Book Classics. Addison-Wesley Publishing Company, Advanced Book Program, Redwood City, CA, 1989. An introduction to algebraic geometry, Notes written with the collaboration of Richard Weiss, Reprint of 1969 original.

[Ful98] William Fulton. *Intersection theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 1998.

[GPH09] P. D. González Pérez and F. Hernando. Quasi-ordinary singularities, essential divisors and Poincaré series. *J. Lond. Math. Soc. (2)*, 79(3):780–802, 2009.

[Har77] Robin Hartshorne. *Algebraic geometry*, volume No. 52 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Heidelberg, 1977.

[Hay99] Takayuki Hayakawa. Blowing ups of 3-dimensional terminal singularities. *Publ. Res. Inst. Math. Sci.*, 35(3):515–570, 1999.

[Hay00] Takayuki Hayakawa. Blowing ups of 3-dimensional terminal singularities. II. *Publ. Res. Inst. Math. Sci.*, 36(3):423–456, 2000.

[Hay05] Takayuki Hayakawa. Gorenstein resolutions of 3-dimensional terminal singularities. *Nagoya Math. J.*, 178:63–115, 2005.

[Hir64] Heisuke Hironaka. Resolution of singularities of an algebraic variety over a field of characteristic zero. I, II. *Ann. of Math. (2)*, 79:109–203; 79 (1964), 205–326, 1964.

[HW97] Martin Henk and Robert Weismantel. The height of minimal Hilbert bases. *Results Math.*, 32(3-4):298–303, 1997.

[IK03] Shihoko Ishii and János Kollár. The Nash problem on arc families of singularities. *Duke Math. J.*, 120(3):601–620, 2003.

[Ish18] Shihoko Ishii. *Introduction to singularities*. Springer, Tokyo, second edition, 2018.

[JK13] Jennifer M. Johnson and János Kollár. Arc spaces of cA-type singularities. *J. Singul.*, 7:238–252, 2013.

[Kaw93] Yujiro Kawamata. The minimal discrepancy of a 3-fold terminal singularity, 1993. Appendix to [Sho93].

[KM98] János Kollár and Shigefumi Mori. *Birational geometry of algebraic varieties*, volume 134 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.

[KSB88] J. Kollár and N. I. Shepherd-Barron. Threefolds and deformations of surface singularities. *Invent. Math.*, 91(2):299–338, 1988.

[Laz04a] Robert Lazarsfeld. *Positivity in algebraic geometry. I*, volume 48 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Se-*

ries. A Series of Modern Surveys in Mathematics]. Springer-Verlag, Berlin, 2004. Classical setting: line bundles and linear series.

[Laz04b] Robert Lazarsfeld. *Positivity in algebraic geometry. II*, volume 48 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2004. Positivity for Vector Bundles, and Multiplier Ideals.

[LJR12] Monique Lejeune-Jalabert and Ana J. Reguera. Exceptional divisors that are not uniruled belong to the image of the Nash map. *J. Inst. Math. Jussieu*, 11(2):273–287, 2012.

[LTZ93] Ji Yong Liu, Leslie E. Trotter, Jr., and Günter M. Ziegler. On the height of the minimal Hilbert basis. *Results Math.*, 23(3-4):374–376, 1993.

[Mar96] Dimitri Markushevich. Minimal discrepancy for a terminal cdv singularity is 1. *J. Math. Sci. Univ. Tokyo*, 3:445–456, 1996.

[Mat02] Kenji Matsuki. *Introduction to the Mori program*. Universitext. Springer-Verlag, New York, 2002.

[Mor85] Shigefumi Mori. On 3-dimensional terminal singularities. *Nagoya Math. J.*, 98:43–66, 1985.

[MS84] David R. Morrison and Glenn Stevens. Terminal quotient singularities in dimensions three and four. *Proc. Amer. Math. Soc.*, 90(1):15–20, 1984.

[Nas95] John F. Nash, Jr. Arc structure of singularities. *Duke Math. J.*, 81(1):31–38, 1995. A celebration of John F. Nash, Jr.

[Rei83] Miles Reid. Minimal models of canonical 3-folds. In *Algebraic varieties and analytic varieties (Tokyo, 1981)*, volume 1 of *Adv. Stud. Pure Math.*, pages 131–180. North-Holland, Amsterdam, 1983.

[Sho93] V. V. Shokurov. Three-fold log flips. *Russian Acad. Sci. Izv. Math.*, 40(1):95–202, 1993.