CRITICAL REVIEWS IN ORAL BIOLOGY & MEDICINE

Inorganic Chemistry of Defensive Peroxidases in the Human Oral Cavity

M. T. Ashby

Department of Chemistry and Biochemistry, University of Oklahoma, Norman, OK 73019, USA; MAshby@ou.edu

J Dent Res 87(10):900-914, 2008

ABSTRACT

The innate host response system is comprised of various mechanisms for orchestrating host response to microbial infection of the oral cavity. The heterogeneity of the oral cavity and the associated microenvironments that are produced give rise to different chemistries that affect the innate defense system. One focus of this review is on how these spatial differences influence the two major defensive peroxidases of the oral cavity, salivary peroxidase (SPO) and myeloperoxidase (MPO). With hydrogen peroxide (H_2O_2) as an oxidant, the defensive peroxidases use inorganic ions to produce antimicrobials that are generally more effective than H_2O_2 itself. The concentrations of the inorganic substrates are different in saliva *vs*. gingival crevicular fluid (GCF). Thus, in the supragingival regime, SPO and MPO work in unison for the exclusive production of hypothiocyanite ($OSCN^-$, a reactive inorganic species), which constantly bathes nascent plaques. In contrast, MPO is introduced to the GCF during inflammatory response, and in that environment it is capable of producing hypochlorite (OCl^-), a chemically more powerful oxidant that is implicated in host tissue damage. A second focus of this review is on inter-person variation that may contribute to different peroxidase function. Many of these differences are attributed to dietary or smoking practices that alter the concentrations of relevant inorganic species in the oral cavity (*e.g.*: fluoride, F^-; cyanide, CN^-; cyanate, OCN^-; thiocyanate, SCN^-; and nitrate, NO_3^-). Because of the complexity of the host and microflora biology and the associated chemistry, it is difficult to establish the significance of the human peroxidase systems during the pathogenesis of oral diseases. The problem is particularly complex with respect to the gingival sulcus and periodontal pockets (where the very different defensive stratagems of GCF and saliva co-mingle). Despite this complexity, intriguing *in vitro* and *in vivo* studies are reviewed here that reveal the interplay between peroxidase function and associated inorganic chemistry.

KEY WORDS: salivary peroxidase, myeloperoxidase, hypochlorite, hypothiocyanite.

Received April 1, 2008; Last revision May 26, 2008; Accepted June 25, 2008

INTRODUCTION

The oral cavity contains a plethora of specific and non-specific defense factors. The non-specific factors include some mucins, proline-rich proteins, salivary glycoproteins, lactoferrin, lysozyme, histatins, cystatins, and peroxidases. This review focuses on the role of peroxidases in the context of oral health and disease, with an emphasis on the relevant inorganic chemistry. Particular attention is paid to the antimicrobial properties of the inorganic chemicals of the oral cavity that are associated with the peroxidases, and to the inter-person differences in the inorganic chemistry of the oral cavity that may influence peroxidase function. For further information on the structures and origins of human oral peroxidases, the reader is referred to the recent review by Tenovuo and co-workers (Ihalin *et al.*, 2006). There are two principal defensive peroxidase systems in the oral cavity, salivary peroxidase (SPO) and myeloperoxidase (MPO). SPO is structurally and catalytically similar to lactoperoxidase (LPO) (Ihalin *et al.*, 2006). *In vivo*, the SPO and LPO systems essentially use only the pseudohalide SCN^- as a substrate to produce $OSCN^-$ (Pruitt *et al.*, 1988). Such defensive peroxidases are commonly found in regions of the human body that are controlled by the mucosa: *e.g.*, breast milk (Shin *et al.*, 2000), lachrymal fluid (Van Haeringen *et al.*, 1979; Tenovuo *et al.*, 1985), and the mucosal lining of the lungs (Gerson *et al.*, 2000). LPO and SPO are coded for the same gene (Ueda *et al.*, 1997). In contrast to peroxidases that essentially employ only SCN^- as a substrate (*e.g.*, LPO and SPO), the MPO system is also capable of oxidizing Cl^- to produce hypochlorite (OCl^-) (Arnhold *et al.*, 2006). Hypobromite (OBr^-) can also be generated by the MPO system (Thomas *et al.*, 1995), but only in minor amounts in the oral cavity. All of the human defensive peroxidases can also utilize iodide (I^-) as a substrate. However, because of sequestration in the thyroid, the environmentally rare halide I^- is not abundant in most physiologic fluids, including the fluids of the oral cavity (Anttonen and Tenovuo, 1981). Accordingly, the limited bioavailability of I^- precludes its significant involvement in host defense.

SPO is a normal, non-inducible component of the saliva of the parotid and submandibular glands (Riva *et al.*, 1978), whereas MPO is an offensive mechanism of neutrophilic polymorphonuclear leukocytes (PMNs). Leukocytes are not normal components of the saliva of healthy individuals, but rather are introduced to the oral cavity by gingival crevicular fluid (GCF) during inflammatory responses (Kowolik and Grant, 1983). The leukocytes in the GCF are comprised of *ca*. 90% PMNs (Ebersole, 2003), and MPO accounts for about 5% of the total PMN protein (Pullar *et al.*, 2000). PMNs degenerate in saliva due to osmotic lysis, thereby releasing the content of the azurophilic granules (including MPO). It has been estimated that *ca*. 75% of the peroxidase activity in mixed saliva is due to MPO, with the remaining activity attributed to SPO (Thomas *et al.*, 1994a). Most of the SPO activity is associated with the soluble portion of the saliva, whereas most of the MPO activity is associated with the sediment (Thomas *et al.*, 1994a). Note that, in contrast to other regions of the mucosa—for example, the lungs—eosinophils are not usually recruited into

Downloaded from jdr.sagepub.com at UNIV OF OKLAHOMA on January 20, 2016 For personal use only. No other uses without permission.

International and American Associations for Dental Research

Downloaded from SHAREOK. Persistent Handle URL: https://hdl.handle.net/11244/25483




Figure 1. Spatial relationship between the inorganic host defense factors of the oral cavity and the ion gradients that influence their relative abundance. Refer to the text for an explanation of pathways A-E and the meaning of the variable Z.

Table 1. Two-electron Redox Couples for X^- (E^o, pH = 7 *vs.* SHE), Apparent Rate Constants (k) of MPO Compound I (x $10^{-4}M^{-1}s^{-1}$) with X^-,[a] Reference Range Values (RRV) of X^- in Physiologic Fluids (μM or mM), and Specificities (S)[b] for Oxidation of X^- by MPO (consensus substrates in **bold**)

X^-	E^o	k	RRV in GCF	GCF S	RRV in Saliva	Saliva S
Cl^-	1.08	2.5	**90 mM**	**6**	25 mM	1
SCN^-	0.77	960	40 μM	1	**1 mM**	**15**

[a] Furtmueller *et al.* (1998).
[b] $S = k^{X^-}_{maj}[X^-_{maj}]/k^{X^-}_{min}[X^-_{min}]$.

the oral cavity, although they can be introduced to saliva *via* the sputum of individuals who have asthma eosinophilia (Spahn, 2007) and from eosinophilic ulcers (rare lesions of the oral mucosa) (Mezei *et al.*, 1995; Hirshberg *et al.*, 2006). Consequently, there is no evidence that eosinophil peroxidase (another common defensive peroxidase that has properties somewhat different from those of LPO, SPO, and MPO) plays a significant role in oral fluids. Accordingly, this review focuses on SPO and MPO.

PEROXIDASES: ORAL HEALTH AND DISEASE

The two disease states of the oral cavity that we consider here are caries and periodontal diseases. The etiology of caries is clear: Acidogenic bacteria cause damage to tooth enamel in the presence of fermentable carbohydrates (*e.g.*, sucrose, fructose, and glucose) (Featherstone, 2000). When the pH at the surface of the tooth falls below 5.5, demineralization proceeds faster than remineralization, and decay ensues. The role of inorganic chemistry in this process is multifaceted: *e.g.*, the (de)mineralization process largely involves the inorganic mineral hydroxyapatite [*ca.* 96% for enamel and 70% for dentin (with some amorphous calcium phosphate)], and the aforementioned peroxidase-derived reactive inorganic species are involved in controlling the microbial growth ($OSCN^-$ in particular, although other inorganic chemical species have been proposed to be significant, *vide infra*).

Like caries, periodontal diseases are also caused by microbial infection (Smalley, 1994; Genco, 1996; Mombelli, 2003). Although the primary cause of periodontal diseases is the accumulation of dental plaque at the gingival margin and the consequential host response (Azuma, 2006), numerous factors affect the severity of the diseases, include smoking (Bergström, 2004), poorly controlled diabetes (Mealey and Oates, 2006), and genetic susceptibility (Baker and Roopenian, 2002; Shapira *et al.*, 2005). Both soft tissues (gingival and periodontal ligaments) and hard tissues (alveolar bone and cementum, which are both largely hydroxyapatite) are affected, but the cause of this tissue damage is a complex and as-yet-unresolved matter. It appears likely that inflammatory agents (including OCl^-) produced by the host (Pullar *et al.*, 2000; Klebanoff, 2005) and virulence factors produced by the infectious agents (Graves *et al.*, 2000) are both responsible for the tissue damage. The supragingival environment in which caries develops and the subgingival environment of periodontal diseases exhibit different chemistries that have a marked influence on the functions and activities of the human defensive peroxidases, which will be discussed next (Fig. 1, Table 1).

SPATIAL NATURE OF PEROXIDASE SYSTEMS AND THEIR CONSENSUS SUBSTRATES

The oxidation reactions that are catalyzed by the peroxidase systems of the oral cavity are governed by the amount of available hydrogen peroxide (H_2O_2), the limiting chemical reagent. A dual-oxidase system from the salivary glands is an endogenous source of H_2O_2 (Geiszt *et al.*, 2003; Donko *et al.*, 2005; Ris-Stalpers, 2006). Oral bacteria also produce H_2O_2 during anaerobic glycolysis (Carlsson *et al.*, 1983). A third source of H_2O_2 is derived from activated neutrophils during oxidative bursts (Dahlgren and Karlsson, 1999; Quinn, 2005). The amounts of OCl^- and $OSCN^-$ that are produced by the MPO system are related to the relative concentrations of Cl^- and SCN^- (van Dalen *et al.*, 1997; Arnhold *et al.*, 2006). At equal concentrations of (pseudo)halide, MPO catalyzes the oxidation of SCN^- about 1000 times faster than Cl^-, but Cl^- is about 1000 times more abundant in most physiologic fluids [*e.g.*, plasma and GCF (Anttonen and Tenovuo, 1981)]. Consequently, comparable amounts of $OSCN^-$ and OCl^- are produced by the MPO system in such fluids. However, SCN^- is essentially the only substrate of MPO in saliva, where the concentration of SCN^- is higher than in most other extracelluar fluids (Tenovuo and Makinen, 1976), as a consequence of its active transport (Fragoso *et al.*, 2004). While analysis of the data in Table 1 suggests that OCl^- should also be generated in saliva, albeit in a minor amount with respect to $OSCN^-$, it can be estimated that the half-life of the OCl^- in saliva is less than 15 μsec, as a consequence of its very fast non-enzymic reaction with SCN^- (Ashby *et al.*, 2004). The reaction of OCl^- and SCN^- yields $OSCN^-$ (Nagy *et al.*, 2006a). Thus, in effect, the only hypohalite of the SPO and MPO systems in saliva is

Downloaded from jdr.sagepub.com at UNIV OF OKLAHOMA on January 20, 2016 For personal use only. No other uses without permission.

International and American Associations for Dental Research

expected to be $OSCN^-$, but a continuum of products is expected at the gingival margin, where a gradient of concentration of Cl^- and SCN^- exists. The spatial relationship between these gradients and the areas of the oral cavity that are respectively controlled by the OCl^- and $OSCN^-$ defense factors is illustrated in Fig. 1:

- Extraphagosomal OCl^- is cytotoxic to oral bacteria (Path A) (Briseno *et al.*, 1992; Webb *et al.*, 1995; Yesilsoy *et al.*, 1995; Barnard *et al.*, 1996; Winniczuk and Parish, 1997; Calas *et al.*, 1998; D'Arcangelo *et al.*, 1998; D'Arcangelo and Varvara, 1998; Huque *et al.*, 1998; Ferreira *et al.*, 1999; Wunder and Bowen, 1999; Spratt *et al.*, 2001; Mikami *et al.*, 2003; Sassone *et al.*, 2003a,b; Moller *et al.*, 2004; Nagayoshi *et al.*, 2004; Radcliffe *et al.*, 2004; Vianna *et al.*, 2004; Carson *et al.*, 2005; Fang *et al.*, 2006; Sena *et al.*, 2006; Ozok *et al.*, 2007) and gingival tissue (Path B) (Schraufstatter *et al.*, 1990; Vissers *et al.*, 1999; Hidalgo and Dominguez, 2000; Pullar *et al.*, 2000; Vile *et al.*, 2000; Hidalgo *et al.*, 2002). Importantly, nearly all of the investigations of the efficacy of OCl^- on oral bacteria have been carried out for single species in planktonic cultures. However, a recent study has focused on the effects of OCl^- on single- and dual-species biofilms of *Fusobacterium nucleatum* and *Peptostreptococcus micros* (Ozok *et al.*, 2007).
- Alternatively, OCl^- can react with SCN^- to produce HOSCN (Path C) (Ashby *et al.*, 2004). HOSCN is also produced by the SPO-catalyzed oxidation of SCN^- by H_2O_2 (Ihalin *et al.*, 2006 ; Nagy *et al.*, 2006a). HOSCN is antimicrobial toward oral bacteria (Path D) (Clem and Klebanof, 1966; Hoogendoorn, 1976; Pruitt *et al.*, 1979; Carlsson *et al.*, 1983; Thomas *et al.*, 1983, 1994b; Ellen *et al.*, 1988; Lopatin *et al.*, 1991; Lumikari *et al.*, 1991; Courtois *et al.*, 1992; Lenander-Lumikari *et al.*, 1993, 1997; van der Hoeven and Camp, 1993; Kirstila *et al.*, 1994; Jones *et al.*, 1998; Fadel and Courtois, 1999, 2001; Yu *et al.*, 2000; Ihalin *et al.*, 2001, 2003; Korpela *et al.*, 2002; Garcia-Graells *et al.*, 2003; Vannini *et al.*, 2004), but relatively non-injurious to the host (Bjoerck and Claesson, 1980; Marshall and Reiter, 1980; White *et al.*, 1983; Carlsson *et al.*, 1984; Carlsson, 1987).
- In addition to the reaction of OCl^- with SCN^-, it may react with other small molecules (Path E) to produce secondary antimicrobials [*e.g.*, when Z is an amine, a cytotoxic chlora–mine is produced, *vide infra* (Abia *et al.*, 1998; Hawkins and Davies, 1998; Hawkins *et al.*, 2003; Davies, 2005)].

The corresponding relevance of OCl^- *vs.* $OSCN^-$ in the oral cavity is related to the aforementioned spatial heterogeneity of the peroxidase defense systems and the corresponding chemistry. For example, the median concentration of $OSCN^-$ in freshly collected whole saliva is *ca.* 10 μM, although the concentration increases when the saliva is incubated at 37°C (Thomas *et al.*, 1980). However, direct measurement of the concentrations of these hypohalite species is problematic, because they are chemically reactive, and consequently the abundance of free ions does not necessarily reflect their significance *in vivo*. The fluxes of the hypohalites (the rates at which these reactive species are produced and consumed) are difficult to define in the context of the oral cavity. An even more complex issue is the relationship between these fluxes with respect to oral health and disease. This topic will be revisited in the concluding section of this review.

ANTIMICROBIAL PROPERTIES OF INORGANIC COMPOUNDS

Chemical Basis of Cytotoxicity

In contrast to antibiotics that typically target a single chemical step in a biosynthetic pathway, inorganic antimicrobials are generally biocides (they have a propensity to cause wholesale disruption of cellular processes) (Ashby, 2007; Zhu, 2007). Accordingly, these inorganic species tend to be cytotoxic, to greater or lesser degrees, to both eukaryotes and prokaryotes. Thus, any discussion of the antimicrobial properties of inorganic compounds toward infectious agents goes hand-in-hand with a related discussion of host tissue damage. It is fascinating to the author that the human body has found generally effective ways of harnessing the potentially indiscriminant cytotoxic properties of some of these compounds for defensive purposes. The cytotoxic properties of the inorganic compounds that are discussed herein can ultimately be traced to their chemistry, which can be roughly divided into two categories: (1) compounds that engage in one-electron (radical) chemistry, and (2) compounds that engage in two-electron chemistry (generally electrophilic, with eventual oxygen atom transfer). An example of one-electron chemistry is the reduction of O_2 by NADPH oxidase to give $O_2^{\cdot-}$ (a reaction that is carried out by PMN NADPH oxidase):

$$NADPH + 2\ O_2 \rightarrow NADP^+ + 2\ O_2^{\cdot-} + H^+$$

Note that NADPH is a two-electron reductant, so the chemical stoichiometry requires one NADPH to react with two oxygen molecules. However, the chemistry in fact involves one-electron steps, *vis-à-vis* enzyme intermediates. The reaction of HOCl with reduced glutathione (GSH) is an example of a two-electron (O-atom transfer) reaction (GSH is the principal cytoplasmic oxidative defense mechanism of eukaryotes, *vide infra*). The reaction occurs *via* a multistep mechanism, because the intermediate sulfenic acid (GSOH) is unstable (Nagy and Ashby, 2007; Nagy *et al.*, 2007b):

$$GSH + HOCl \rightarrow GSOH + H^+ + Cl^-$$
$$GSOH + GSH \rightarrow GSSG + H_2O$$
$$2\ GSH + HOCl \rightarrow GSSG + H_2O + H^+ + Cl^-$$

Note that the first equivalent of GSH that reacts with HOCl involves a two-electron reaction (O-atom transfer, albeit probably *via* the hydrolysis of a sulfenyl chloride intermediate), as does the second reaction, even though the stoichiometry of the net reaction makes it appear that GSH is a one-electron reductant. In fact, the thiyl radical ($GS^{\cdot}$) is never involved in the reaction. While the distinction between one- and two-electron processes may appear to be a superfluous detail, in fact the difference distinguishes radical processes from non-radical processes. Radicals tend to target unsaturated functional groups in lipids, nucleotides, and aromatic amino acids (Buettner, 1993). In contrast, the hypohalites tend to target the nucleophiles in proteins (Hawkins *et al.*, 2003; Davies, 2005).

Unsaturated organic compounds (*e.g.*, aromatic amino acids and nucleotides) are particularly susceptible to derivation by high-energy radical species; hence, nuclear damage and mutagenesis are frequently the result of one-electron chemistry

Downloaded from jdr.sagepub.com at UNIV OF OKLAHOMA on January 20, 2016 For personal use only. No other uses without permission.

International and American Associations for Dental Research

Table 2. Major Inorganic (Reactive) Oxygen Species in the Oral Cavity

Name	Symbol	Major Sources in the Oral Cavity
Triplet oxygen	$^3O_2$	The atmosphere
Singlet oxygen	$^1O_2$	Peroxidase-catalyzed reactions of H_2O_2
Superoxide	$O_2^{\cdot-}$	NADPH reductase and leakage from peroxidases
Hydrogen peroxide	H_2O_2	Human dual oxidases (Duox) and aerobic metabolism of glucose
Hydroxyl radical	$^{\cdot}OH$	Metal-catalyzed homolysis of H_2O_2 (Fenton chemistry)
Ozone	O_3	Catalytic decomposition of $^1O_2$ by SIgA

Table 3. Major Inorganic (Pseudo) Hypohalites in the Oral Cavity

Name	Symbol	Major Sources in the Oral Cavity
Hypochlorite	OCl^-	Myeloperoxidase
Hypobromite	OBr^-	Myeloperoxidase (eosinophil peroxidase?)
Hypothiocyanite	$OSCN^-$	Myeloperoxidase and salivary peroxidase

(Box *et al.*, 2001; Marnett, 2002; Wang, 2008). Because radical chemistry tends to be very facile and comparatively indiscriminant from a chemical perspective, it is difficult for cells to mount an effective defense against radical species. In contrast, the reactivities of two-electron oxidants are typically related to the nucleophilicities of their reaction partners. Consequently, the chemistry of two-electron oxidants is usually well-defined (*cf.* radical chemistry). Cysteine (Cys) and methionine (Met) are usually the most reactive amino acid residues toward two-electron oxidants (because sulfur-containing compounds tend to be good nucleophiles) and are therefore often the first targets of two-electron oxidants (Hawkins *et al.*, 2003). It is not a coincidence that the cytoplasms of eukaryotes (Meister, 1988; Fernandes *et al.*, 2007) and many prokaryotes (Fahey *et al.*, 1978; Smirnova and Oktyabrsky, 2005) contain high concentrations of GSH (a tripeptide containing Cys, which is used to combat oxidative stress by two-electron oxidants that operate by the aforementioned electrophilic mechanism) (Meister, 1988). It should be noted that glutathione also erects a significant defense against radical species (Sitte and Von Zglinicki, 2003; Djordjevic, 2004) (Table 2).

Oxygen Derivatives

Water (H_2O) and molecular oxygen (O_2) represent limiting extremes in the oxidation state of the element oxygen (O) in an aqueous environment. From a thermodynamic perspective, O exists as O_2 in an overall aerobic (oxidative) environment, whereas it exists as H_2O in an anaerobic (reductive) environment. The oral cavity contains microenvironments that represent these extremes. Molecular oxygen is itself "antimicrobial" toward strict anaerobes (which are routinely found in mature supragingival plaques and are abundant in subgingival plaques). However, ground-state O_2 ($^3O_2$, triplet oxygen, a di-radical) is not generally included among the so-called "reactive oxygen species" (ROS, Table 2). The ROS of Table 2 can be divided into the radical species ($O_2^{\cdot-}$, $^{\cdot}OH$) and the "closed-shell" species ($^1O_2$, H_2O_2, and O_3). Hydroxyl radical ($^{\cdot}OH$) is not produced in large quantities by the defensive peroxidases of the oral cavity, so it will not be further discussed here, but we refer the reader to reviews of the role of ROS in periodontal tissue destruction for more information (Waddington *et al.*, 2000; Chapple and Matthews, 2007). In contrast, $O_2^{\cdot-}$ is pertinent in that it is produced by NADPH oxidase during neutrophilic respiratory bursts (*vide supra*). Despite being a radical, $O_2^{\cdot-}$ is relatively chemically unreactive, and mammalian cells [and some oral bacteria (Amano *et al.*, 1986)] contain superoxide dismutase that catalyzes the disproportionation of $O_2^{\cdot-}$ to give H_2O_2 and O_2 (Packer, 2002). It is noteworthy that the conjugate acid $HOO^{\cdot}$ (hydroperoxyl or perhydroxyl radical) is considerably more reactive than the conjugate base $O_2^{\cdot-}$. It is conceivable that $HOO^{\cdot}$ (pK_a *ca.* 4.8) (Bielski *et al.*, 1985) plays a role in the oral cavity (*e.g.*, during the development of acidogenic plaques). However, since $O_2^{\cdot-}$ and $HOO^{\cdot}$ are in rapid acid/base equilibrium, no distinction will be made between the two species in this review (*cf.* HOCl *vs.* OCl^-, *vide infra*). Of the closed-shell species, only H_2O_2, $^1O_2$, and O_3 are relevant to our discussion.

Hydrogen peroxide is a powerful oxidant that is produced in the oral cavity by the aforementioned mechanisms (Duox, anaerobic glycolysis, disproportionation of $O_2^{\cdot-}$, etc.). Like $O_2^{\cdot-}$, H_2O_2 is relatively chemically inert. For example, it takes more than an hour for H_2O_2 to react with millimolar concentrations of Cys (Ashby and Nagy, 2006a,b). Nonetheless, H_2O_2 is cytotoxic toward mammalian cells (Ward, 1991) and most prokaryotes (Asad *et al.*, 2004). Singlet oxygen has been detected in saliva (Takahama, 1993; Kou and Takahama, 1995; Sun *et al.*, 2006). However, many of the reported measurements of $^1O_2$ remain controversial, because the probes that are used to detect it tend to be insensitive and frequently non-specific (Martinez *et al.*, 2000). Nonetheless, it is believed that human peroxidases produce $^1O_2$ during their decomposition of H_2O_2 (Kanofsky, 1991). In addition to the use of therapeutic O_3 (Azarpazhooh and Limeback, 2008), it has been suggested that $^1O_2$ is converted to O_3 *via* an immunoglobulin-catalyzed reaction (Wentworth *et al.*, 2000, 2002). Secretory IgA (SIgA), the most abundant immunoglobulin in saliva, is also proposed to catalyze the reaction (Uehara *et al.*, 2006). However, the involvement of O_3 remains controversial, because the probes that have been used are not specific (Kettle *et al.*, 2004; Smith, 2004; Kettle and Winterbourn, 2005).

As a consequence of their reactive nature, all of the ROS exhibit cytotoxic properties. The relative importance of ROS as defensive agents in the oral cavity is difficult to assess, because various amounts act on microcosm plaques in diverse environments. There are likely synergisms in multi-species plaques. For example, pure cultures of oral streptococci produce H_2O_2 (they are catalase-negative), but H_2O_2 is not found in dental plaque or salivary sediment, despite streptococci being major components of their mixed bacterial populations. This is presumably due to the fast consumption of free H_2O_2 by catalase-positive species of bacteria (*e.g.*, *Neisseria, Haemophilus, Actinomyces*, and *Staphylococcus* spp.) (Ryan and Kleinberg, 1995). Furthermore, the SPO and MPO systems of the oral cavity may protect H_2O_2-sensitive bacteria (Adamson and Carlsson, 1982). In addition to synergism between bacterial species, there are likely to be additive and/or cooperative effects

Downloaded from jdr.sagepub.com at UNIV OF OKLAHOMA on January 20, 2016 For personal use only. No other uses without permission.

International and American Associations for Dental Research

between defensive agents. For example, there is some evidence that $O_2^{\cdot-}$ can act synergistically with OCl^- (*vide infra*) to induce oxidative damage (Hawkins *et al.*, 2002) (Table 3).

(Pseudo) Hypohalites

The archetypal example of a biocide is hypochlorite (OCl^-, the principal component of household bleach). In sufficient concentrations, it is toxic to all life. Note that the reactive form of OCl^- (and the other hypohalites as well) is the corresponding conjugate acid, hypohalous acid (HOCl, pK_a = 7.4). As a neutral species, HOCl is presumably membrane-permeable, and therefore more cytotoxic. However, since the acid-base equilibrium between HOCl and OCl^- is exceedingly fast, the issue of which species is actually active is academic. Under acidic conditions (and in the presence of excess Cl^-), OCl^- comproportionates to give Cl_2, the corresponding halogen (Adam *et al.*, 1992). However, since the equilibrium between the hypohalites and the halogens is relatively fast, they are not generally treated as different biocides. Instead, the term "total active chlorine" is often used to describe the sum amount of hypohalous acid, hypohalite, and halogen. Furthermore, HOCl/OCl^- are the predominant species at physiologic pH. The relative ease with which the halides are oxidized is $I^- > Br^- > Cl^-$ (note that F^- is never oxidized in an aqueous environment), so the trend in oxidative strengths of the hypohalous acids is HOCl > HOBr >> HOI. As mentioned before, I^- is not abundant in most physiologic fluids, so only Cl^- and Br^- are relevant to this discussion.

HOCl and HOBr exhibit somewhat promiscuous reaction chemistry (the relative reactivities of HOCl toward proteinaceous groups are Cys ≈ Met >> cystine ≈ His ≈ α-amino > Trp > Lys >> Tyr ≈ Arg > backbone amides > Gln ≈ Asn, and a similar trend is observed for HOBr) (Pattison and Davies, 2001, 2004). However, the kinetics of some of the reactions of HOCl, and especially HOBr, approaches the diffusion limit (Nagy *et al.*, 2006b). As a consequence of their facile reaction, HOCl and HOBr probably exhibit poor chemical selectivity in a biological setting. In addition, secondary reactive species are produced during the reactions of HOCl and HOBr that likely contribute significantly to the overall toxicity—for example, haloamines (Grisham *et al.*, 1984; Abia *et al.*, 1998; Hawkins and Davies, 1998, 2003; Davies, 2005). From a chemical perspective, it is appropriate to view the reactions of HOX (X = Cl, Br) as a redox cascade (thermodynamically downhill) with the eventual production of chemically stable derivatives. Because of the labile nature of many of the intermediate species in such chemical cascades, and because of the fact that many of these species exhibit similar chemistries (*e.g.*, HOCl and chloroamines are both electrophilic chlorinating agents, albeit with different reactivities), it is difficult to chart the consequences of the damage that occurs. Accordingly, the complexity of the chemistry precludes a definitive assignment of the antimicrobial mechanism of HOX (X = Cl, Br). Nonetheless, there have been many studies of the effects of HOX (X = Cl, Br) on both eukaryotes (Hawkins and Davies, 2000; Hidalgo and Dominguez, 2000; Hawkins *et al.*, 2001; Hidalgo *et al.*, 2002; Soszynski *et al.*, 2002; King *et al.*, 2004) and prokaryotes (Albrich *et al.*, 1981, 1986; Albrich and Hurst, 1982; Barrette *et al.*, 1987, 1989; Hurst *et al.*, 1991; Palazzolo *et al.*, 2005). Although it is problematic to follow the reaction cascade that begins with the production of HOX (X = Cl, Br) *in vivo*, there is considerable interest in evaluation of the relevance of the HOX (X = Cl, Br) chemistry, particularly in the context of the host tissue damage that occurs with inflammatory disease. One approach to monitoring the chemistry of HOX (X = Cl, Br) *in vivo* is through the use of biomarkers, thermodynamically stable derivatives of HOX (X = Cl, Br). Promising biomarkers that appear to be specific for HOX (X = Cl, Br) include 3-halotyrosines (Winterbourn and Kettle, 2000) and a sulfonamide derivative of GSH (Harwood *et al.*, 2006).

In the field of inorganic chemistry, SCN^- is called a "pseudohalide", because its reaction chemistry frequently mirrors that of the halides (Lappert and Pyszora, 1966). This is reflected in the fact that defensive peroxidases use SCN^- (in addition to the halides) as a substrate. The oxidation potential of SCN^- falls between those of I^- and Br^-. Accordingly, the chemical properties of HOSCN are most similar to those of HOI (Nagy *et al.*, 2007a). In contrast to HOCl and HOBr (which react with a variety of functional groups), the only characterized reactions of HOSCN are with thiol moieties (which are among the most powerful nucleophiles), generally Cys and its derivatives (Ashby and Aneetha, 2004; Nimmo *et al.*, 2007; Lemma and Ashby, 2008). Given that SH groups are apparently the targets of the HOSCN, it is important to note that roughly 40% of all enzymes are rendered ineffective by chemical agents that are reactive toward thiols (Leung-Toung *et al.*, 2002). Thus, the destruction of functional SH moieties by HOSCN is one basis for its cytotoxicity. Importantly, the hypohalites can be interconverted through their reactions with other halides, but the process must be exothermic. For example, HOCl (Ashby *et al.*, 2004) and HOBr (Nagy *et al.*, 2006b) react with SCN^- to give HOSCN and the corresponding halides, but not *vice versa*. *In vivo*, these reactions are important because they restrict the lifetimes of the more powerful (and less discriminant) hypohalites, thereby limiting their propensity to cause collateral host tissue damage. In addition, the antimicrobial $OSCN^-$ is produced in the reaction (Nagy *et al.*, 2006a).

Halides

Fluoride is the only halide that is known to be antimicrobial without oxidation (Hamilton, 1990; Van Loveren, 1990; Marquis, 1995; Jenkins, 1999). Fluoride influences the metabolism of cariogenic and other bacteria *via* multiple mechanisms (Marquis *et al.*, 2003): F^- can bind directly to many enzymes (especially metalloenzymes) (Segal *et al.*, 1968; Wever and Bakkenist, 1980; Zgliczynski *et al.*, 1983; Thibodeau *et al.*, 1985; Ferrari *et al.*, 1997; Suzuki and Ohshima, 2003), thereby affecting their activities; catalase is inhibited by F^- [thereby affecting the ability of H_2O_2 to kill oral bacteria (Phan *et al.*, 2001)]; and some F^- complexes of metals (*e.g.*, AlF_4^- and $BeF_3^-{\cdot}H_2O$) can mimic phosphate, thereby affecting a variety of enzymes and regulatory phosphatases (Thongboonkerd *et al.*, 2002). The weak-acid property of HF (pK_a = 3.15), which is a transmembrane proton conductor, appears to be important for inducing the glycolytic inhibition of oral bacteria that is observed at low pH in dental plaque (Eisenberg and Marquis, 1981) (Table 4).

Nitrogen Derivatives

After many decades of angst about nitrates in our diet and their propensity to form potentially carcinogenic *N*-nitroso derivatives (Eichholzer and Gutzwiller, 2003), there is

Downloaded from jdr.sagepub.com at UNIV OF OKLAHOMA on January 20, 2016 For personal use only. No other uses without permission.

International and American Associations for Dental Research

mounting evidence that NO_3^- is concentrated in saliva for beneficial purposes (McKnight *et al.*, 1999) (although the potential deleterious properties of NO_3^- on systemic health should not be discounted). The concentration of NO_3^- in saliva is proportional to the dietary intake (Eisenbrand *et al.*, 1980), it varies with the salivary flow rate (Granli *et al.*, 1989), and it is influenced by smoking (Tsuchiya *et al.*, 2002). Since NO_3^- is a relatively inert chemical species, the mechanism of antimicrobial action of NO_3^- probably involves a redox cascade. Facultative anaerobic bacteria in the oral cavity use NO_3^- as a terminal electron acceptor (*cf.* O_2 for aerobic bacteria) to produce nitrite (NO_2^-). Acidified NO_2^- inhibits the growth and affects the survival of cariogenic bacteria (*e.g.*, *Streptococcus mutans, Lactobacillus casei*, and *Actinomyces naeslundii*) (Silva Mendez *et al.*, 1999). Acidified NO_2^- has a similar effect on periodontal bacteria (*e.g.*, *Fusobacterium nucleatum, Eikenella corrodens*, and *Porphyromonas gingivalis*) (Allaker *et al.*, 2001). Importantly, while the growth of periodontal bacteria is known to be inhibited by acid in the absence of NO_2^-, there is a dose-dependent decrease in these bacteria in the presence of NO_2^- (Allaker *et al.*, 2001). Nitrous acid (HNO_2, $pK_a = 3.4$) is unstable and will spontaneously disproportionate: $3\ HNO_2 \rightarrow H_3O^+ + NO_3^- + 2\ ^{\cdot}NO$. Some bacteria possess nitrite reductase (*e.g.*, *S. mutans*), an enzyme that is capable of accelerating the disproportionation of NO_2^- (Choudhury *et al.*, 2007). It appears that nitric oxide ($^{\cdot}NO$) is the antimicrobial component of the $NO_3^-/NO_2^-/^{\cdot}NO$ redox cascade (Fang, 1997; Smith *et al.*, 1999; Sato *et al.*, 2008). The mechanism by which $^{\cdot}NO$ induces cell death is the subject of ongoing investigation. Alternative models include "oxidative stress" and "nitrosative stress" [*e.g.*, nitrosylation of proteins without a major alteration in cellular redox state (Eu *et al.*, 2000)]. Nitric oxide also reacts with $O_2^{\cdot-}$ to produce peroxynitrite ($ONOO^-$), which may also contribute to collateral host tissue damage in the oral cavity (Lohinai and Szabo, 1998; Lohinai *et al.*, 2001; Barley *et al.*, 2004).

INTER-PERSON DIFFERENCES

There is considerable variability in the physiological concentrations of many of the chemically stable inorganic ions that have been discussed herein. In most cases, these differences can be attributed to diets or smoking. Given the influence of these ions on the activity and function of the defensive peroxidases, there has been some interest in correlating interperson differences to oral disease. As a caveat, it is important to note that many of the relevant inorganic ions are chemically reactive, and consequently the abundance of free ions may not reflect their relevance *in vivo*. Some ions may exist as transient species (*e.g.*, OCl^-) or as their conjugates with reaction partners (*e.g.*, CN^- and OCN^-). In some cases, steady-state concentrations of reactive species may accumulate (*e.g.*, $OSCN^-$), but measured concentrations may not reflect the time-dependent flux of such species.

Thiocyanate

The main source of SCN^- *in vivo* is CN^-, *vide infra*. Cyanide is principally introduced by the digestion of glucosinolate-containing vegetables (*e.g.*, the *Brassica*) (Weuffen *et al.*, 1984). However, as a consequence of detoxification of hydrogen cyanide (HCN, $pK_a = 9.2$, which is known to be present in microgram amounts *per* cigarette), the level of SCN^- in smokers is considerably elevated relative to that in non-smokers. Indeed, this difference is routinely used as a biomarker for the evaluation of smoking behavior (Morabia *et al.*, 2001). While most of the CN^- that is converted to SCN^- *in vivo* is exogenous in origin, endogenous sources contribute the sulfur *vis-à-vis* a multitude of reactions, some of which are enzyme-catalyzed (Wood, 1975). It has been shown that higher concentrations of SCN^- in saliva can contribute to an enhancement of peroxidase activity (Tenovuo, 1976; Lamberts *et al.*, 1984; Fonteh *et al.*, 2005; Tahboub *et al.*, 2005). Furthermore, as noted earlier, SCN^- is a potent sequestering agent for some reactive oxidants (Ashby *et al.*, 2004; Nagy *et al.*, 2006b). Accordingly, one might conclude that higher SCN^- (either as a consequence of diet or through smoking) should result in a suppression of oral disease. However, while smokers have elevated SCN^- and $OSCN^-$ levels in their saliva, no corresponding correlation with dental caries has been observed (Lamberts *et al.*, 1984). Interestingly, although there is not an association between smoking and caries among adults, there is a positive correlation between environmental (second-hand) tobacco smoke and primary tooth caries in children (Shenkin *et al.*, 2004). In contrast to caries, there is a strong correlation between smoking (and consequently SCN^- levels) and some periodontal diseases (Rivera-Hidalgo, 2003). Since there is also a correlation between smoking and SCN^- levels in GCF, SCN^- levels presumably exhibit a positive correlation with periodontal diseases. However, SCN^- is only one of many inorganic and organic chemicals that are elevated by smoking. Two other inorganic ions are CN^- and cyanate (OCN^-), *vide infra*.

Table 4. Major Inorganic (Reactive) Nitrogen Compounds in the Oral Cavity

Name	Symbol	Major Sources in the Oral cavity
Nitrate	NO_3^-	Diet
Nitrite	NO_2^-	Reduction of nitrate by oral microflora
Nitric oxide	$^{\cdot}NO$	Acidification of nitrite and by enzymic reactions
Peroxynitrite	$ONOO^-$	Reaction of $^{\cdot}NO$ and $O_2^{\cdot-}$

Cyanide

It addition to dietary sources (*e.g.*, cyanogenic glucosides, *vide supra*) and tobacco smoke, other sources of CN^- *in vivo* include micro-organisms (in particular certain pseudomonads) and industrial exposure (*e.g.*, *vis-à-vis* organonitriles) (Wong-Chong *et al.*, 2006). There appear to be no studies that have determined the normal concentration of CN^- in the fluids of the oral cavity. However, there is a statistical correlation between blood and salivary SCN^- levels (Tsuge *et al.*, 2000). As noted earlier, the concentrations of SCN^- in physiological fluids (including GCF and saliva) in smokers are substantially higher than those for non-smokers. Therefore, it follows that the oral cavity is exposed to higher concentrations of CN^- for smokers compared with non-smokers. HCN contributes to the loss of peroxidase activity in saliva upon exposure to cigarette smoke (Klein *et al.*, 2003). This is due to the strong complexation of CN^- to the iron-active sites of the peroxidases (fundamentally the same mechanism that renders CN^- toxic to the respiratory system

Downloaded from jdr.sagepub.com at UNIV OF OKLAHOMA on January 20, 2016 For personal use only. No other uses without permission.

International and American Associations for Dental Research

vis-à-vis the complexation of hemoglobin and myoglobin). It is noteworthy that there is no correlation between CN^- in plasma and the concentration of HCN in breath (Lundquist *et al.*, 1988). Furthermore, the concentrations of HCN measured in breath are higher than expected for blood concentrations, which suggested a local production of HCN in the oropharynx. Under some circumstances, $OSCN^-$ can decompose to give CN^- (Aune and Thomas, 1977). While the major cyanide-derived product of the decomposition of $OSCN^-$ is OCN^-, we have observed the formation of substantial amounts of CN^- during the hydrolysis of thiocyanogen [$(SCN)_2$, analogous to a halogen] at neutral pH (unpublished observations). It is conceivable that the latter reaction is the source of HCN in breath.

Cyanate

Cyanate (OCN^-) is produced by the oxidation of CN^-. Thus, OCN^- levels are higher in smokers. Under some conditions, the decomposition of $OSCN^-$ also produces OCN^- (Oram and Reiter, 1966). MPO is inhibited by OCN^- (Qian *et al.*, 1997). The inhibition could be caused by heme binding of OCN^- (thereby blocking the active site) or by carbamylation of the protein by OCN^-. It is noteworthy that the functional impairment of proteins through carbamylation by OCN^- is thought to promote human inflammatory diseases (Wang *et al.*, 2007). However, the possible relevance of carbamylation in the oral cavity has not been investigated. There is very limited information available regarding the effect of OCN^- on oral bacteria (Morita, 1977).

Fluoride

In the absence of supplementation by fluoride-containing dentrifices, the concentration of F^- in saliva (and presumably GCF) is somewhat lower than the concentration in plasma (Oliveby *et al.*, 1989). The normal concentration of F^- in saliva (*ca.* 1 μM) can increase markedly after the ingestion of F^-, and change dynamically thereafter (Dawes and Weatherell, 1990). An early study that used PMN granules (and not isolated enzymes) suggested that F^- does not influence the activity of MPO (which was determined by measurement of the iodination of zymosan) (Gabler and Leong, 1980). In contrast, the same study reported that the iodination of zymosan was inhibited by F^- for intact PMNs (Gabler and Leong, 1980). However, the pH-dependent competitive inhibition of isolated LPO (Segal *et al.*, 1968; Thibodeau *et al.*, 1985) and MPO (Zgliczynski *et al.*, 1983; van den Abbeele *et al.*, 1992) by F^- has been demonstrated. For example, half of the activity of bovine LPO occurs for < 0.05 mM at pH 4, 0.3 mM at pH 5, 4.0 mM at pH 6, and greater than 10 mM at pH 7, as assayed with 5 mM I^- and 150 μM H_2O_2 (Thibodeau *et al.*, 1985). It is noteworthy that SPO was found to have lower pH optima relative to LPO, but it also was inhibited by F^- at sufficiently low pH (Thibodeau *et al.*, 1985). A similar inhibitory effect of F^- on peroxidase activity has been observed in whole saliva (Thibodeau *et al.*, 1985; Hannuksela *et al.*, 1994; van den Abbeele *et al.*, 1995). These observations suggest that F^- in dental plaque may inhibit the peroxidase defense system. However, when F^- and $OSCN^-$ are added simultaneously at pH 5.0, an additive effect of growth inhibition of *S. mutans* was observed (Lenander-Lumikari *et al.*, 1997). Thus, the small inhibitory effect of F^- on the defensive peroxidase systems of the oral cavity may be offset by the combinatorial antimicrobial effects of F^- and $OSCN^-$.

Nitrate and Nitrite

Basal levels of NO_3^- in saliva are 10-20 times those found in plasma (and presumably GCF) (Duncan *et al.*, 1995; Benjamin and McKnight, 1999; Pannala *et al.*, 2003; Lundberg *et al.*, 2004). For the average diet in the US, *ca.* 80% of dietary NO_3^- is derived from vegetables (White, 1975). Cured meats, for example, typically represent a minor source of NO_3^- in most diets. However, urinary, plasma, and saliva NO_3^- concentrations increase markedly after the consumption of a high-nitrate meal (Pannala *et al.*, 2003). The maximum concentration is achieved within a few hours following the meal, and NO_3^- concentrations return to basal levels within 24 hours. The amount of NO_3^- that is excreted in the urine following the consumption of a high-nitrate meal suggests that the majority of urinary NO_3^- can be accounted for in dietary sources (Pannala *et al.*, 2003). In contrast to NO_3^-, an increase in NO_2^- is observed in saliva only after the consumption of a high-nitrate meal (Pannala *et al.*, 2003), which is consistent with the fact that NO_3^- is metabolized to NO_2^- by bacterial flora on the posterior surface of the tongue in rat models (Duncan *et al.*, 1995). Nitrite has been shown to enhance the reactivity of LPO (Reszka *et al.*, 1997, 1998, 1999; van der Vliet *et al.*, 1997; Gebicka, 1999; Bruck *et al.*, 2001) and MPO (Burner *et al.*, 2000). In addition to an enhancement in the rate of catalysis by NO_2^-, it has been suggested that NO_3^- reduces acidity in the oral cavity (Li *et al.*, 2007). Thus, NO_2^- and NO_3^- may play a beneficial role in the oral cavity. We note that there has apparently been no attempt to correlate oral health with inter-person differences in nitrate-reducing capacity.

PEROXIDE TOXICITY AND PEROXIDASE FUNCTION

At sufficient concentrations, H_2O_2 is cytotoxic to mammalian cell lines, including human epithelial cells (Mattana *et al.*, 1992) and gingival fibroblasts (Tenovuo and Larjava, 1984; Tipton *et al.*, 1995a). While H_2O_2 is a relatively chemically inert molecule, its homolysis to give $^{\cdot}OH$ radicals is catalyzed by transition metals, particularly iron, in a reaction that is referred to as Fenton chemistry (Prousek, 2007). Much of the cytotoxicity of H_2O_2 is attributed to Fenton chemistry (Winterbourn, 1995). In the presence of SCN^-, the LPO system protects cultured mammalian cells against H_2O_2 toxicity (Hänström *et al.*, 1983). This is consistent with the observation that $OSCN^-$ is not toxic toward mammalian cells (Bjoerck and Claesson, 1980; Marshall and Reiter, 1980; White *et al.*, 1983; Carlsson *et al.*, 1984; Carlsson, 1987). It has been previously suggested that one of the important roles of human peroxidases is to detoxify H_2O_2 to prevent host tissue damage (Carlsson, 1987; Tipton *et al.*, 1995b). Since many prokaryotes are also sensitive to H_2O_2, the human peroxidase systems may also protect certain bacteria by sequestering H_2O_2. However, the $OSCN^-$ that is produced is inhibitory toward most bacteria. Consequently, it is difficult to envisage the net effect of H_2O_2 sequestration *vs*. $OSCN^-$ on bacterial growth (Fig. 2).

THE ANTIMICROBIAL DEPLETION MODEL

The efficacy of chemically reactive antimicrobial agents can be diffusion-limited. This is one (but certainly not the only) explanation for the greater sensitivity of planktonic cultures to chemically reactive antimicrobials (Stewart, 1994;

Downloaded from jdr.sagepub.com at UNIV OF OKLAHOMA on January 20, 2016 For personal use only. No other uses without permission.

International and American Associations for Dental Research




Figure 2. Relationship between the reactivities of inorganic defense factors of the oral cavity and maturation of plaque biofilms. The penetrability of the hypohalites and their derivatives is inversely related to their reactivities and the extent of maturation of the biofilm. See the text for a discussion of the possibility that the defensive factors could be generated within biofilms.

Stewart and Raquepas, 1995; Dodds *et al.*, 2000; Stewart *et al.*, 2001, 2004; Hunt *et al.*, 2005). Planktonic cultures are generally agitated, and consequently fluid flows by a convection mechanism that transports solutes rapidly. In contrast, fluid flow in biofilms can be highly restricted. For thick biofilms, such as the plaques of the oral cavity, the primary mechanism of solute transport is diffusion (unagitated flow). For highly reactive molecules such as HOCl, which also exhibit promiscuous reaction chemistry, the biomass of plaques offers a formidable barrier to penetration. While HOCl is an extremely effective antimicrobial toward planktonic cell suspensions, it is a comparatively inefficient killing agent with respect to biofilms. For example, a 600-fold increase in concentration of HOCl (*per* cell) is required to kill certain biofilms of *Staphylococcus aureus*, compared with planktonic cultures of the same species (Luppens *et al.*, 2002). More chemically selective antimicrobials, *e.g.*, $OSCN^-$, are more likely to penetrate thick biofilms before reacting. The relationship between chemical reactivity of the antimicrobial and biofilm penetrability forms the basis of the Antimicrobial Depletion Model (Fig. 2). There is presumably an inverse relationship between penetrability with respect to the antimicrobial reactivity. However, penetrability, of course, is not the only factor that determines the efficacy of the antimicrobial: $OSCN^-$ is generally considered to be bacteriostatic, whereas OCl^- is bactericidal. But, as mentioned before, there is a tradeoff for the higher reactivity/poorer selectivity of OCl^-: the potential of host tissue damage. In the oral cavity, it may be advantageous for the body to use a collection of antimicrobials that exhibit a continuum of reactivity. Thus, complementing HOCl is taurine chloroamine (TauCl), a less reactive (Carr *et al.*, 2001) and more selective (Peskin and Winterbourn, 2006) electrophilic chlorinating agent that may play a role in host defense in the oral cavity (Woldring, 1955; Mainnemare *et al.*, 2004). Similarly, we have discovered a corresponding less-reactive derivative of $OSCN^-$, thiocarbamate-S-oxide [$H_2NC(=O)$-S-O^-] (Nagy *et al.*, 2007c). Thiocarbamate-S-oxide, which is formed by the hydrolysis of $OSCN^-$, is one of the chemical species formed during the redox cascade that results in the decomposition of $OSCN^-$ (Nagy *et al.*, 2007c). Recently, we have learned that $H_2NC(=O)$-S-O^- reacts with thiols *via* a mechanism that is analogous to the one that is observed for $OSCN^-$, albeit with slower kinetics (unpublished observations). While the effect of $H_2NC(=O)$-S-O^- on bacterial physiology remains to be investigated, its discovery demonstrates that there is much to be learned about the inorganic antimicrobials produced by the defensive peroxidases.

The Antimicrobial Depletion Model does not explicitly include the possibility that defensive peroxidases might generate hypohalites within the biofilms, thereby obviating the need for reactive species to diffuse into biofilms. We note that the production of acid in salivary sediment is more effectively inhibited when $OSCN^-$ is produced by sediment-bound peroxidases than when salivary sediment is treated with exogenous $OSCN^-$ (Tenovuo, 1979). However, salivary sediment has a larger buffer capacity and "organic load" (*e.g.*, non-viable cells) than plaque (Singer *et al.*, 1983), so it is not clear whether the penetration of $OSCN^-$ into plaque is comparably diffusion-limited. The production of hypohalites from within plaques requires the transport of the components of peroxidase systems into the biofilm: peroxidase, (pseudo) halide, and H_2O_2. Relevant to the issue of diffusion of peroxidases into plaques, it has been previously observed that LPO binds to *S. mutans* (LPO is a cationic protein, and the outer membranes of Gram-positive bacteria are negatively charged) (Pruitt *et al.*, 1979). While cell-bound LPO remains catalytically active initially, it is eventually inactivated (Pruitt *et al.*, 1979). Thus, the electrostatic attraction of the peroxidases for cells and the subsequent inactivation of the cell-bound enzyme may preclude the generation of hypohalites within a thick plaque (Pruitt *et al.*, 1979). With regard to the availabilities of the other components of the peroxidase defense systems, it is probable that the concentrations of the (pseudo)halides in plaques reflect their concentrations in the physiologic fluids that surround them (*i.e.*, saliva or GCF), but it was noted earlier that H_2O_2 has not been found in dental plaques (Ryan and Kleinberg, 1995). The issue of whether or not the human defensive peroxidase systems are active in thick plaques remains unresolved.

CONCLUSION

The distinctive chemical environments of the supragingival and subgingival regions impose restrictions on the human peroxidase defense strategies of the oral cavity (*e.g.*, *vis-à-vis* substrate bioavailability). Inter-person differences may also influence the function and activity of the enzymes and the chemistry of the reactive species that the enzymes generate. The abilities of the defensive agents produced by the peroxidase systems to differentiate between host tissues and the microbiota are an unsettled issue. Oral diseases are accompanied by microbial shifts of dental plaque, so a more subtle issue is whether these agents differentiate between commensal and pathogenic microbiota. While it seems likely that ecological principles, as in the March ecological hypothesis (Marsh, 2003), can be applied to explain microbial shifts, it is not yet clear what ecological pressures are inducing these microbial shifts, nor is the root cause of host tissue damage completely

Downloaded from jdr.sagepub.com at UNIV OF OKLAHOMA on January 20, 2016 For personal use only. No other uses without permission.

International and American Associations for Dental Research

clear. For dental caries in enamel, damage is clearly a direct effect of the plaque. In dentinal and root caries, some host response may be involved (as in periodontitis). The microbial shift during cariogenesis is toward acidogenic and acid-tolerant Gram-positive bacteria, which displace the acid-sensitive commensal microbiota that are associated with intact tooth tissues. Since supragingivally generated $OSCN^-$ targets Gram-positive bacteria (*S. mutans* may be particularly affected), it is possible that $OSCN^-$ may be important in modifying plaque cariogenicity. Other important health-maintaining functions of $OSCN^-$ may also occur, such as restricting supragingival intra-oral cross-infection by periodontal pathogens. During the development of periodontal diseases, an increase in disease-associated anaerobes occurs in conjunction with increased inflammation. The inflammatory response is primarily due to antigens that have been introduced by the bacteria. It is unclear whether the inflammation is due to changes in the antigens introduced by the microbial shift (Lamster and Novak, 1992), or whether the microbial shift has been induced by the inflammatory response, or both (Dalwai *et al.*, 2006). Is the host tissue damage of periodontal diseases solely the consequence of host mediators (*e.g.*, HOCl), or is there direct attack by bacterial virulence factors such as proteolytic enzymes? Definitive answers to these questions are critical for a rational approach to combating oral disease—Does one treat the inflammation, or does one target pathogenic bacteria, or is it necessary to treat both (Van Dyke, 2007)? Other risk factors being similar, is inter-individual variation in peroxidase activity a key factor in determining why some people develop oral disease and others do not? Whether inter-person differences between the inorganic chemistry of the oral cavity contribute to oral diseases is a topic that deserves further attention.

ACKNOWLEDGMENT

The author is supported by Public Health Service grant R21 DE016889-01A2 from the National Institute of Dental and Craniofacial Research.

REFERENCES

Abia L, Armesto XL, Canle L M, Garcia MV, Santaballa JA (1998). Oxidation of aliphatic amines by aqueous chlorine. *Tetrahedron* 54:521-530.

Adam LC, Fabian I, Suzuki K, Gordon G (1992). Hypochlorous acid decomposition in the pH 5-8 region. *Inorg Chem* 31:3534-3541.

Adamson M, Carlsson J (1982). Lactoperoxidase and thiocyanate protect bacteria from hydrogen peroxide. *Infect Immun* 35:20-24.

Albrich JM, Hurst JK (1982). Oxidative inactivation of *Escherichia coli* by hypochlorous acid. Rates and differentiation of respiratory from other reaction sites. *FEBS Lett* 144:157-161.

Albrich JM, McCarthy CA, Hurst JK (1981). Biological reactivity of hypochlorous acid: implications for microbicidal mechanisms of leukocyte myeloperoxidase. *Proc Natl Acad Sci USA* 78:210-214.

Albrich JM, Gilbaugh JH 3rd, Callahan KB, Hurst JK (1986). Effects of the putative neutrophil-generated toxin, hypochlorous acid, on membrane permeability and transport systems of *Escherichia coli*. *J Clin Invest* 78:177-184.

Allaker RP, Mendez LSS, Hardie JM, Benjamin N (2001). Antimicrobial effect of acidified nitrite on periodontal bacteria. *Oral Microbiol Immunol* 16:253-256.

Amano A, Tamagawa H, Shizukuishi S, Tsunemitsu A (1986). Superoxide dismutase, catalase and peroxidases in oral anaerobic bacteria. *J Osaka Univ Dent Sch* 26:187-192.

Anttonen T, Tenovuo J (1981). Crevicular thiocyanate and iodide ions: cofactors of the antimicrobial peroxidase system in leucocytes. *Proc Finn Dent Soc* 77:318-323.

Arnhold J, Monzani E, Furtmueller PG, Zederbauer M, Casella L, Obinger C (2006). Kinetics and thermodynamics of halide and nitrite oxidation by mammalian heme peroxidases. *Eur J Inorg Chem* 19:3801-3811.

Asad NR, Asad LMBO, Bonacossa de Almeida CE, Felzenszwalb I, Cabral-Neto JB, Leitão AC (2004). Several pathways of hydrogen peroxide action that damage the *E. coli* genome. *Genet Mol Biol* 27:291-303.

Ashby MT (2007). Where old biocides meet new biocides: hypohalite defense factors of the human oral cavity. *ACS Symposium Series (New Biocides Development)* 967:269-290.

Ashby MT, Aneetha H (2004). Reactive sulfur species: aqueous chemistry of sulfenyl thiocyanates. *J Am Chem Soc* 126:10216-10217.

Ashby MT, Nagy P (2006a). On the kinetics and mechanism of the reaction of cysteine and hydrogen peroxide in aqueous solution. Comments. *J Pharm Sci* 95:15-18.

Ashby MT, Nagy P (2006b). Revisiting a proposed kinetic model for the reaction of cysteine and hydrogen peroxide via cysteine sulfenic acid. *Int J Chem Kinet* 39:32-38.

Ashby MT, Carlson AC, Scott MJ (2004). Redox buffering of hypochlorous acid by thiocyanate in physiologic fluids. *J Am Chem Soc* 126:15976-15977.

Aune TM, Thomas EL (1977). Accumulation of hypothiocyanite ion during peroxidase-catalyzed oxidation of thiocyanate ion. *Eur J Biochem* 80:209-214.

Azarpazhooh A, Limeback H (2008). The application of ozone in dentistry: a systematic review of literature. *J Dent* 36:104-116.

Azuma M (2006). Fundamental mechanisms of host immune responses to infection. *J Periodontal Res* 41:361-373.

Baker PJ, Roopenian DC (2002). Genetic susceptibility to chronic periodontal disease. *Microbes Infect* 4:1157-1167.

Barley RDC, Pollock S, Shallow MC, Peters E, Lam EWN (2004). Tobacco-related-compound-induced nitrosative stress injury in the hamster cheek pouch. *J Dent Res* 83:903-908.

Barnard D, Davies J, Figdor D (1996). Susceptibility of *Actinomyces israelii* to antibiotics, sodium hypochlorite and calcium hydroxide. *Int Endod J* 29:320-326.

Barrette WC Jr, Albrich JM, Hurst JK (1987). Hypochlorous acid-promoted loss of metabolic energy in *Escherichia coli*. *Infect Immun* 55:2518-2525.

Barrette WC Jr, Hannum DM, Wheeler WD, Hurst JK (1989). General mechanism for the bacterial toxicity of hypochlorous acid: abolition of ATP production. *Biochemistry* 28:9172-9178.

Benjamin N, McKnight G (1999). Metabolism of nitrate in humans—implications for nitrate intake. *R Soc Chem* 237(Spec Pub):281-288.

Bergström J (2004). Tobacco smoking and chronic destructive periodontal disease. *Odontology* 92:1-8.

Bielski BHJ, Cabelli DE, Arudi RL, Ross AB (1985). Reactivity of perhydroxyl/superoxide radicals in aqueous solution. *J Phys Chem Ref Data* 14:1041-1100.

Bjoerck L, Claesson O (1980). Correlation between concentration of hypothiocyanate and antibacterial effect of the lactoperoxidase system against *Escherichia coli*. *J Dairy Sci* 63:919-922.

Box HC, Dawidzik JB, Budzinski EE (2001). Free radical-induced double lesions in DNA. *Free Radic Biol Med* 31:856-868.

Briseno BM, Wirth R, Hamm G, Standhartinger W (1992). Efficacy of different irrigation methods and concentrations of root canal irrigation solutions on bacteria in the root canal. *Endod Dent Traumatol* 8:6-11.

Bruck TB, Fielding RJ, Symons MCR, Harvey PJ (2001). Mechanism of nitrite-stimulated catalysis by lactoperoxidase. *Eur J Biochem* 268:3214-3222.

Downloaded from jdr.sagepub.com at UNIV OF OKLAHOMA on January 20, 2016 For personal use only. No other uses without permission.

International and American Associations for Dental Research

Buettner GR (1993). The pecking order of free radicals and antioxidants: lipid peroxidation, α-tocopherol, and ascorbate. *Arch Biochem Biophys* 300:535-543.

Burner U, Furtmuller PG, Kettle AJ, Koppenol WH, Obinger C (2000). Mechanism of reaction of myeloperoxidase with nitrite. *J Biol Chem* 275:20597-20601.

Calas P, Rochd T, Druilhet P, Azais JM (1998). In vitro adhesion of two strains of *Prevotella nigrescens* to the dentin of the root canal: the part played by different irrigation solutions. *J Endod* 24:112-115.

Carlsson J (1987). Salivary peroxidase: an important part of our defense against oxygen toxicity. *J Oral Pathol* 16:412-416.

Carlsson J, Iwami Y, Yamada T (1983). Hydrogen peroxide excretion by oral streptococci and effect of lactoperoxidase-thiocyanate-hydrogen peroxide. *Infect Immun* 40:70-80.

Carlsson J, Edlund MB, Hänstroem L (1984). Bactericidal and cytotoxic effects of hypothiocyanite-hydrogen peroxide mixtures. *Infect Immun* 44:581-586.

Carr AC, Hawkins CL, Thomas SR, Stocker R, Frei B (2001). Relative reactivities of N-chloramines and hypochlorous acid with human plasma constituents. *Free Radic Biol Med* 30:526-536.

Carson KR, Goodell GG, McClanahan SB (2005). Comparison of the antimicrobial activity of six irrigants on primary endodontic pathogens. *J Endod* 31:471-473.

Chapple IL, Matthews JB (2007). The role of reactive oxygen and antioxidant species in periodontal tissue destruction. *Periodontol 2000* 43:160-232.

Choudhury T, Sato EF, Inoue M (2007). Nitrite reductase in *Steptoccocus mutans* plays a critical role in the survival of this pathogen in oral cavity. *Oral Microbiol Immunol* 22:384-389.

Clem WH, Klebanof SJ (1966). Inhibitory effect of saliva on glutamic acid accumulation by *Lactobacillus acidophilus* and role of lactoperoxidase-thiocyanate system. *J Bacteriol* 91:1848-1853.

Courtois P, Majerus P, Labbe M, van den Abbeele A, Yourassowsky E, Pourtois M (1992). Susceptibility of anaerobic microorganisms to hypothiocyanite produced by lactoperoxidase. *Acta Stomatol Belg* 89:155-162.

D'Arcangelo C, Varvara G (1998). A comparative in-vitro study of the bactericidal efficacy of sodium hypochlorite and chlorhexidine gluconate plus cetrimide on root canal anaerobic bacterial flora. *Minerva Stomatol* 47:381-386.

D'Arcangelo C, Di Nardo Di Maio F, Varvara G (1998). Alpha-hemolytic Streptococci and root canal irrigants. An evaluation of the bactericidal efficacy of sodium hypochlorite and chlorhexidine gluconate plus cetrimide. *Minerva Stomatol* 47:367-371.

Dahlgren C, Karlsson A (1999). Respiratory burst in human neutrophils. *J Immunol Methods* 232:3-14.

Dalwai F, Spratt DA, Pratten J (2006). Modeling shifts in microbial populations associated with health or disease. *Appl Environ Microbiol* 72:3678-3684.

Davies MJ (2005). The oxidative environment and protein damage. *Biochim Biophys Acta* 1703:93-109.

Dawes C, Weatherell JA (1990). Kinetics of fluoride in the oral fluids. *J Dent Res* 69(Spec Iss):638-644.

Djordjevic VB (2004). Free radicals in cell biology. *Int Rev Cytol* 237:57-89.

Dodds MG, Grobe KJ, Stewart PS (2000). Modeling biofilm antimicrobial resistance. *Biotechnol Bioeng* 68:456-465.

Donko A, Peterfi Z, Sum A, Leto T, Geiszt M (2005). Dual oxidases. *Phil Trans R Soc London B: Biol Sci* 360:2301-2308.

Duncan C, Dougall H, Johnston P, Green S, Brogan R, Leifert C, *et al.* (1995). Chemical generation of nitric oxide in the mouth from the enterosalivary circulation of dietary nitrate. *Nat Med* 1:546-551.

Ebersole JL (2003). Humoral immune responses in gingival crevice fluid: local and systemic implications. *Periodontol 2000* 31:135-166.

Eichholzer M, Gutzwiller F (2003). Dietary nitrates, nitrites and N-nitroso compounds and cancer risk with special emphasis on the epidemiological evidence (abstract). In: Food safety: contaminants and toxins. Zürich, Switzerland: CABI Publishing, pp. 217-234.

Eisenberg AD, Marquis RE (1981). Enhanced transmembrane proton conductance in *Streptococcus mutans* gs-5 due to ionophores and fluoride. *Antimicrob Agents Chemother* 19:807-812.

Eisenbrand G, Spiegelhalder B, Preussmann R (1980). Nitrate and nitrite in saliva. *Oncology* 37:227-231.

Ellen RP, Schwarz-Faulkner S, Grove DA (1988). Coaggregation among periodontal pathogens, emphasizing *Bacteroides gingivalis-Actinomyces viscosus* cohesion on a saliva-coated mineral surface. *Can J Microbiol* 34:299-306.

Eu JP, Liu L, Zeng M, Stamler JS (2000). An apoptotic model for nitrosative stress. *Biochemistry* 39:1040-1047.

Fadel M, Courtois P (1999). Effect of peroxidase-generated hypothiocyanite on the survival rate of *Porphyromonas gingivalis* NCTC 11834. *Med Sci Res* 27:667-669.

Fadel M, Courtois P (2001). Inhibitory effect of lactoperoxidase-generated hypothiocyanite upon black pigmented anaerobe growth. *Int J Mol Med* 8:59-62.

Fahey RC, Brown WC, Adams WB, Worsham MB (1978). Occurrence of glutathione in bacteria. *J Bacteriol* 133:1126-1129.

Fang FC (1997). Mechanisms of nitric oxide-related antimicrobial activity. *J Clin Invest* 99:2818-2825.

Fang M, Chen JH, Xu XL, Yang PH, Hildebrand HF (2006). Antibacterial activities of inorganic agents on six bacteria associated with oral infections by two susceptibility tests. *Int J Antimicrob Agents* 27:513-517.

Featherstone JD (2000). The science and practice of caries prevention. *J Am Dent Assoc* 131:887-899.

Fernandes PN, Mannarino SC, Silva CG, Pereira MD, Panek AD, Eleutherio ECA (2007). Oxidative stress response in eukaryotes: effect of glutathione, superoxide dismutase and catalase on adaptation to peroxide and menadione stresses in *Saccharomyces cerevisiae*. *Redox Rep* 12:236-244.

Ferrari RP, Ghibaudi EM, Traversa S, Laurenti E, De Gioia L, Salmona M (1997). Spectroscopic and binding studies on the interaction of inorganic anions with lactoperoxidase. *J Inorg Biochem* 68:17-26.

Ferreira CM, Bonifacio KC, Froner IC, Ito IY (1999). Evaluation of the antimicrobial activity of three irrigating solutions in teeth with pulpal necrosis. *Braz Dent J* 10:15-21.

Fonteh FA, Grandison AS, Lewis MJ (2005). Factors affecting lactoperoxidase activity. *Int J Dairy Technol* 58:233-236.

Fragoso MA, Fernandez V, Forteza R, Randell SH, Salathe M, Conner GE (2004). Transcellular thiocyanate transport by human airway epithelia. *J Physiol* 561(Pt 1):183-194.

Furtmueller PG, Burner U, Obinger C (1998). Reaction of myeloperoxidase Compound I with chloride, bromide, iodide, and thiocyanate. *Biochemistry* 37:17923-17930.

Gabler WL, Leong PA (1980). Effect of fluoride on polymorphonuclear leukocyte myeloperoxidase. *J Dent Res* 59:135.

Garcia-Graells C, Van Opstal I, Vanmuysen SCM, Michiels CW (2003). The lactoperoxidase system increases efficacy of high-pressure inactivation of foodborne bacteria. *Int J Food Microbiol* 81:211-221.

Gebicka L (1999). Kinetic studies on the oxidation of nitrite by horseradish peroxidase and lactoperoxidase. *Acta Biochim Pol* 46:919-927.

Geiszt M, Witta J, Baffi J, Lekstrom K, Leto TL (2003). Dual oxidases represent novel hydrogen peroxide sources supporting mucosal surface host defense. *FASEB J* 17:1502-1504.

Genco RJ (1996). Current view of risk factors for periodontal diseases. *J Periodontol* 67(10 Suppl):1041S-1049S.

Gerson C, Sabater J, Scuri M, Torbati A, Coffey R, Abraham JW, *et al.* (2000). The lactoperoxidase system functions in bacterial

Downloaded from jdr.sagepub.com at UNIV OF OKLAHOMA on January 20, 2016 For personal use only. No other uses without permission.

International and American Associations for Dental Research

clearance of airways. *Am J Respir Cell Mol Biol* 22:665-671.
Granli T, Dahl R, Brodin P, Bockman OC (1989). Nitrate and nitrite concentrations in human saliva: variations with salivary flow-rate. *Food Chem Toxicol* 27:675-680.
Graves DT, Jiang Y, Genco C (2000). Periodontal disease: bacterial virulence factors, host response and impact on systemic health. *Curr Opin Infect Dis* 13:227-232.
Grisham MB, Jefferson MM, Melton DF, Thomas EL (1984). Chlorination of endogenous amines by isolated neutrophils. Ammonia-dependent bactericidal, cytotoxic, and cytolytic activities of the chloramines. *J Biol Chem* 259:10404-10413.
Hamilton IR (1990). Biochemical effects of fluoride on oral bacteria. *J Dent Res* 69(Spec Iss):660-667.
Hannuksela S, Tenovuo J, Roger V, Lenander-Lumikari M, Ekstrand J (1994). Fluoride inhibits the antimicrobial peroxidase systems in human whole saliva. *Caries Res* 28:429-434.
Hänström L, Johansson A, Carlsson J (1983). Lactoperoxidase and thiocyanate protect cultured mammalian cells against hydrogen peroxide toxicity. *Med Biol* 61:268-274.
Harwood DT, Kettle AJ, Winterbourn CC (2006). Production of glutathione sulfonamide and dehydroglutathione from gsh by myeloperoxidase-derived oxidants and detection using a novel lc-ms/ms method. *Biochem J* 399:161-168.
Hawkins CL, Davies MJ (1998). Hypochlorite-induced damage to proteins: formation of nitrogen-centered radicals from lysine residues and their role in protein fragmentation. *Biochem J* 332(Pt 3):617-625.
Hawkins CL, Davies MJ (2000). Hypochlorite-induced damage to red blood cells. Evidence for the formation of nitrogen-centered radicals. *Redox Rep* 5:57-59.
Hawkins CL, Brown BE, Davies MJ (2001). Hypochlorite- and hypobromite-mediated radical formation and its role in cell lysis. *Arch Biochem Biophys* 395:137-145.
Hawkins CL, Rees MD, Davies MJ (2002). Superoxide radicals can act synergistically with hypochlorite to induce damage to proteins. *FEBS Lett* 510:41-44.
Hawkins CL, Pattison DI, Davies MJ (2003). Hypochlorite-induced oxidation of amino acids, peptides and proteins. *Amino Acids* 25:259-274.
Hidalgo E, Dominguez C (2000). Growth-altering effects of sodium hypochlorite in cultured human dermal fibroblasts. *Life Sci* 67:1331-1344.
Hidalgo E, Bartolome R, Dominguez C (2002). Cytotoxicity mechanisms of sodium hypochlorite in cultured human dermal fibroblasts and its bactericidal effectiveness. *Chem Biol Interact* 139:265-282.
Hirshberg A, Amariglio N, Akrish S, Yahalom R, Rosenbaum H, Okon E, *et al.* (2006). Traumatic ulcerative granuloma with stromal eosinophilia: a reactive lesion of the oral mucosa. *Am J Clin Pathol* 126:522-529.
Hoogendoorn H (1976). Microbial aspects of dental caries. The inhibitory action of the lactoperoxidase system on *Streptococcus mutans* and other microorganisms. *Microb Aspects Dent Caries Proc Workshop* 2:353-357.
Hunt SM, Hamilton MA, Stewart PS (2005). A 3D model of antimicrobial action on biofilms. *Water Sci Technol* 52:143-148.
Huque J, Kota K, Yamaga M, Iwaku M, Hoshino E (1998). Bacterial eradication from root dentine by ultrasonic irrigation with sodium hypochlorite. *Int Endod J* 31:242-250.
Hurst JK, Barrette WC Jr, Michel BR, Rosen H (1991). Hypochlorous acid and myeloperoxidase-catalyzed oxidation of iron-sulfur clusters in bacterial respiratory dehydrogenases. *Eur J Biochem* 202:1275-1282.
Ihalin R, Loimaranta V, Lenander-Lumikari M, Tenovuo J (2001). The sensitivity of *Porphyromonas gingivalis* and *Fusobacterium nucleatum* to different (pseudo)halide-peroxidase combinations compared with mutans streptococci. *J Med Microbiol* 50:42-48.
Ihalin R, Nuutila J, Loimaranta V, Lenander M, Tenovuo J, Lilius EM (2003). Susceptibility of *Fusobacterium nucleatum* to killing by peroxidase-iodide-hydrogen peroxide combination in buffer solution and in human whole saliva. *Anaerobe* 9:23-30.
Ihalin R, Loimaranta V, Tenovuo J (2006). Origin, structure, and biological activities of peroxidases in human saliva. *Arch Biochem Biophys* 445:261-268.
Jenkins GN (1999). Review of fluoride research since 1959. *Arch Oral Biol* 44:985-992.
Jones MN, Hill KJ, Kaszuba M, Creeth JE (1998). Antibacterial reactive liposomes encapsulating coupled enzyme systems. *Int J Pharm* 162:107-117.
Kanofsky JR (1991). Peroxidase-catalyzed generation of singlet oxygen and of free radicals. In: Peroxidases in chemistry and biology. Everse J, Everse KG, Grisham MB, editors. Boston: CRC Publishers, pp. 219-231.
Kettle AJ, Winterbourn CC (2005). Do neutrophils produce ozone? An appraisal of current evidence. *BioFactors* 24:41-45.
Kettle AJ, Clark BM, Winterbourn CC (2004). Superoxide converts indigo carmine to isatin sulfonic acid: implications for the hypothesis that neutrophils produce ozone. *J Biol Chem* 279:18521-18525.
King DA, Hannum DM, Qi JS, Hurst JK (2004). Hocl-mediated cell death and metabolic dysfunction in the yeast *Saccharomyces cerevisiae*. *Arch Biochem Biophys* 423:170-181.
Kirstila V, Lenander-Lumikari M, Tenovuo J (1994). Effects of a lactoperoxidase-system-containing toothpaste on dental plaque and whole saliva in vivo. *Acta Odontol Scand* 52:346-353.
Klebanoff SJ (2005). Myeloperoxidase: friend and foe. *J Leukoc Biol* 77:598-625.
Klein I, Nagler RM, Toffler R, van der Vliet A, Reznick AZ (2003). Effect of cigarette smoke on oral peroxidase activity in human saliva: role of hydrogen cyanide. *Free Radic Biol Med* 35:1448-1452.
Korpela A, Yu X, Loimaranta V, Lenander-Lumikari M, Vacca-Smith A, Wunder D, *et al.* (2002). Lactoperoxidase inhibits glucosyltransferases from *Streptococcus mutans* in vitro. *Caries Res* 36:116-121.
Kou F, Takahama U (1995). Hydrogen peroxide-induced luminescence and evolution of molecular oxygen in human saliva. *Arch Oral Biol* 40:15-21.
Kowolik MJ, Grant M (1983). Myeloperoxidase activity in human gingival crevicular neutrophils. *Arch Oral Biol* 28:293-295.
Lamberts BL, Pruitt KM, Pedersen ED, Golding MP (1984). Comparison of salivary peroxidase system components in caries-free and caries-active naval recruits. *Caries Res* 18:488-494.
Lamster IB, Novak MJ (1992). Host mediators in gingival crevicular fluid—implications for the pathogenesis of periodontal disease. *Crit Rev Oral Biol Med* 3:31-60.
Lappert MF, Pyszora H (1966). Pseudohalides of group IIIb and IVb elements. In: Advances in inorganic chemistry and radiochemistry. Vol. 9. Sharpe AG, editor. USA: Academic Press, pp. 133-184.
Lemma K, Ashby MT (2008). Reactive sulfur species: kinetics and mechanism of the equilibrium between cysteine sulfenyl thiocyanate and cysteine thiosulfinate ester in acidic aqueous solution. *J Org Chem* 73:3017-3023.
Lenander-Lumikari M, Tenovuo J, Mikola H (1993). Effects of a lactoperoxidase system-containing toothpaste on levels of hypothiocyanite and bacteria in saliva. *Caries Res* 27:285-291.
Lenander-Lumikari M, Loimaranta V, Hannuksela S, Tenovuo J, Ekstrand J (1997). Combined inhibitory effect of fluoride and hypothiocyanite on the viability and glucose metabolism of *Streptococcus mutans*, serotype c. *Oral Microbiol Immunol* 12:231-235.
Leung-Toung R, Li W, Tam TF, Karimian K (2002). Thiol-dependent

Downloaded from jdr.sagepub.com at UNIV OF OKLAHOMA on January 20, 2016 For personal use only. No other uses without permission.

International and American Associations for Dental Research

enzymes and their inhibitors: a review. *Curr Med Chem* 9:979-1002.

Li H, Thompson I, Carter P, Whiteley A, Bailey M, Leifert C, *et al.* (2007). Salivary nitrate—an ecological factor in reducing oral acidity. *Oral Microbiol Immunol* 22:67-71.

Lohinai ZM, Szabo C (1998). Role of nitric oxide in physiology and pathophysiology of periodontal tissues. *Med Sci Monit* 4:1089-1095.

Lohinai Z, Stachlewitz R, Virag L, Szekely AD, Hasko G, Szabo C (2001). Evidence for reactive nitrogen species formation in the gingivomucosal tissue. *J Dent Res* 80:470-475.

Lopatin DE, LaBelle D, Lee SW (1991). Measurement of relative avidity of antibodies reactive with *Porphyromonas* (*Bacteroides*) *gingivalis* in the sera of subjects having adult periodontitis. *J Periodontal Res* 26:167-175.

Lumikari M, Soukka T, Nurmio S, Tenovuo J (1991). Inhibition of the growth of *Streptococcus mutans, Streptococcus sobrinus* and *Lactobacillus casei* by oral peroxidase systems in human saliva. *Arch Oral Biol* 36:155-160.

Lundberg JO, Weitzberg E, Cole JA, Benjamin N (2004). Opinion: Nitrate, bacteria and human health. *Nat Rev Microbiol* 2:593-602.

Lundquist P, Rosling H, Sörbo B (1988). The origin of hydrogen cyanide in breath. *Arch Toxicol* 61:270-274.

Luppens SBI, Reij MW, van der Heijden RWL, Rombouts FM, Abee T (2002). Development of a standard test to assess the resistance of *Staphylococcus aureus* biofilm cells to disinfectants. *Appl Environ Microbiol* 68:4194-4200.

Mainnemare A, Megarbane B, Soueidan A, Daniel A, Chapple ILC (2004). Hypochlorous acid and taurine-N-monochloramine in periodontal diseases. *J Dent Res* 83:823-831.

Marnett LJ (2002). Oxy radicals, lipid peroxidation and DNA damage. *Toxicology* 181-182:219-222.

Marquis RE (1995). Antimicrobial actions of fluoride for oral bacteria. *Can J Microbiol* 41:955-964.

Marquis RE, Clock SA, Mota-Meira M (2003). Fluoride and organic weak acids as modulators of microbial physiology. *FEMS Microbiol Rev* 26:493-510.

Marsh PD (2003). Are dental diseases examples of ecological catastrophes? *Microbiology* 149(Pt 2):279-294.

Marshall VME, Reiter B (1980). Comparison of the antibacterial activity of the hypothiocyanite anion towards *Streptococcus lactis* and *Escherichia coli*. *J Gen Microbiol* 120:513-516.

Martinez GR, Di Mascio P, Bonini MG, Augusto O, Briviba K, Sies H, *et al.* (2000). Peroxynitrite does not decompose to singlet oxygen ($^1O_2$) and nitroxyl (NO^-). *Proc Natl Acad Sci USA* 97:10307-10312.

Mattana A, Juliano C, Bennardini F, Mignano M, Marceddu S, Picci V, *et al.* (1992). Hydrogen peroxide-induced cytotoxicity in cultured epithelial cells (WISH): a functional and morphological study. *Toxicol in Vitro* 6:533-541.

McKnight GM, Duncan CW, Leifert C, Golden MH (1999). Dietary nitrate in man: friend or foe? *Br J Nutr* 81:349-358.

Mealey BL, Oates TW (2006). Diabetes mellitus and periodontal diseases. *J Periodontol* 77:1289-1303.

Meister A (1988). Glutathione metabolism and its selective modification. *J Biol Chem* 263:17205-17208.

Mezei MM, Tron VA, Stewart WD, Rivers JK (1995). Eosinophilic ulcer of the oral mucosa. *J Am Acad Dermatol* 33:734-740.

Mikami M, Kato C, Saito K (2003). Characterization of the cell surface substances of *Fusobacterium nucleatum* that induce a chemiluminescence response by human neutrophils. *Nippon Shishubyo Gakkai Kaishi [J Jpn Assoc Periodontol]* 45:215-228 [article in Japanese].

Moller AJ, Fabricius L, Dahlén G, Sundqvist G, Happonen R-P (2004). Apical periodontitis development and bacterial response to endodontic treatment. Experimental root canal infections in monkeys with selected bacterial strains. *Eur J Oral Sci* 112:207-215.

Mombelli A (2003). Periodontitis as an infectious disease: specific features and their implications. *Oral Dis* 9(Suppl 1):6-10.

Morabia A, Bernstein MS, Curtin F, Berode M (2001). Validation of self-reported smoking status by simultaneous measurement of carbon monoxide and salivary thiocyanate. *Prev Med* 32:82-88.

Morita M (1977). Inhibitory effects of some constituents of cigarette smoke on the lactate production of salivary microorganisms and *Streptococcus mutans*. *Shika Kiso Igakkai Zasshi* 19:349-365 [article in Japanese].

Nagayoshi M, Kitamura C, Fukuizumi T, Nishihara T, Terashita M (2004). Antimicrobial effect of ozonated water on bacteria invading dentinal tubules. *J Endod* 30:778-781.

Nagy P, Ashby MT (2007). Reactive sulfur species: kinetics and mechanisms of the oxidation of cysteine by hypohalous acid to give cysteine sulfenic acid. *J Am Chem Soc* 129:14082-14091.

Nagy P, Alguindigue SS, Ashby MT (2006a). Lactoperoxidase-catalyzed oxidation of thiocyanate by hydrogen peroxide: a reinvestigation of hypothiocyanite by nuclear magnetic resonance and optical spectroscopy. *Biochemistry* 45:12610-12616.

Nagy P, Beal JL, Ashby MT (2006b). Thiocyanate is an efficient endogenous scavenger of the phagocytic killing agent hypobromous acid. *Chem Res Toxicol* 19:587-593.

Nagy P, Lemma K, Ashby MT (2007a). Kinetics and mechanism of the comproportionation of hypothiocyanous acid and thiocyanate to give thiocyanogen in acidic aqueous solution. *Inorg Chem* 46:285-292.

Nagy P, Lemma K, Ashby MT (2007b). Reactive sulfur species: kinetics and mechanisms of the reaction of cysteine thiosulfinate ester with cysteine to give cysteine sulfenic acid. *J Org Chem* 72:8838-8846.

Nagy P, Wang X, Lemma K, Ashby MT (2007c). Reactive sulfur species: hydrolysis of hypothiocyanite to give thiocarbamate-S-oxide. *J Am Chem Soc* 129:15756-15757.

Nimmo SLA, Lemma K, Ashby MT (2007). Reactions of cysteine sulfenyl thiocyanate with thiols to give unsymmetrical disulfides. *Heteroatom Chemistry* 18:467-471.

Oliveby A, Lagerlöf F, Ekstrand J, Dawes C (1989). Studies on fluoride excretion in human whole saliva and its relation to flow rate and plasma fluoride levels. *Caries Res* 23:243-246.

Oram JD, Reiter B (1966). The inhibition of streptococci by lactoperoxidase, thiocyanate and hydrogen peroxide. The oxidation of thiocyanate and the nature of the inhibitory compound. *Biochem J* 100:382-388.

Ozok AR, Wu MK, Luppens SBI, Wesselink PR (2007). Comparison of growth and susceptibility to sodium hypochlorite of mono- and dual-species biofilms of *Fusobacterium nucleatum* and *Peptostreptococcus* (*Micromonas*) *micros*. *J Endod* 33:819-822.

Packer L (2002). Superoxide dismutase. In: Methods in enzymology. Vol. 349. San Diego, CA: Academic press, 400 pp.

Palazzolo AM, Suquet C, Konkel ME, Hurst JK (2005). Green fluorescent protein-expressing *Escherichia coli* as a selective probe for HOCl generation within neutrophils. *Biochemistry* 44:6910-6919.

Pannala AS, Mani AR, Spencer JPE, Skinner V, Bruckdorfer KR, Moore KP, *et al.* (2003). The effect of dietary nitrate on salivary, plasma, and urinary nitrate metabolism in humans. *Free Radic Biol Med* 34:576-584.

Pattison DI, Davies MJ (2001). Absolute rate constants for the reaction of hypochlorous acid with protein side chains and peptide bonds. *Chem Res Toxicol* 14:1453-1464.

Pattison DI, Davies MJ (2004). Kinetic analysis of the reactions of hypobromous acid with protein components: implications for cellular damage and use of 3-bromotyrosine as a marker of oxidative stress. *Biochemistry* 43:4799-4809.

Downloaded from jdr.sagepub.com at UNIV OF OKLAHOMA on January 20, 2016 For personal use only. No other uses without permission.

International and American Associations for Dental Research

Peskin AV, Winterbourn CC (2006). Taurine chloramine is more selective than hypochlorous acid at targeting critical cysteines and inactivating creatine kinase and glyceraldehyde-3-phosphate dehydrogenase. *Free Radic Biol Med* 40:45-53.

Phan TN, Kirsch AM, Marquis RE (2001). Selective sensitization of bacteria to peroxide damage associated with fluoride inhibition of catalase and pseudocatalase. *Oral Microbiol Immunol* 16:28-33.

Prousek J (2007). Fenton chemistry in biology and medicine. *Pure Appl Chem* 79:2325-2338.

Pruitt KM, Adamson M, Arnold R (1979). Lactoperoxidase binding to streptococci. *Infect Immun* 25:304-309.

Pruitt KM, Mansson-Rahemtulla B, Baldone DC, Rahemtulla F (1988). Steady-state kinetics of thiocyanate oxidation catalyzed by human salivary peroxidase. *Biochemistry* 27:240-245.

Pullar JM, Vissers MCM, Winterbourn CC (2000). Living with a killer: the effects of hypochlorous acid on mammalian cells. *IUBMB Life* 50:259-266.

Qian M, Eaton JW, Wolff SP (1997). Cyanate-mediated inhibition of neutrophil myeloperoxidase activity. *Biochem J* 326(Pt 1):159-166.

Quinn MT (2005). The neutrophil respiratory burst oxidase. In: Neutrophils. 2nd ed. Gabrilovich DI. editor. London: Imperial College Press, pp. 35-84.

Radcliffe CE, Potouridou L, Qureshi R, Habahbeh N, Qualtrough A, Worthington H, *et al.* (2004). Antimicrobial activity of varying concentrations of sodium hypochlorite on the endodontic microorganisms *Actinomyces israelii, A. naeslundii, Candida albicans* and *Enterococcus faecalis*. *Int Endod J* 37:438-446.

Reszka KJ, Matuszak Z, Chignell CF (1997). Lactoperoxidase-catalyzed oxidation of the anticancer agent mitoxantrone by nitrogen dioxide ($NO_2^{\cdot}$) radicals. *Chem Res Toxicol* 10:1325-1330.

Reszka KJ, Matuszak Z, Chignell CF (1998). Lactoperoxidase-catalyzed oxidation of melanin by reactive nitrogen species derived from nitrite (NO_2^-): an EPR study. *Free Radic Biol Med* 25:208-216.

Reszka KJ, Matuszak Z, Chignell CF, Dillon J (1999). Oxidation of biological electron donors and antioxidants by a reactive lactoperoxidase metabolite from nitrite (NO_2^-): an EPR and spin trapping study. *Free Radic Biol Med* 26:669-678.

Ris-Stalpers C (2006). Physiology and pathophysiology of the duoxes. *Antioxid Redox Signal* 8:1563-1572.

Riva A, Puxeddu P, Del Fiacco M, Testa-Riva F (1978). Ultrastructural localization of endogenous peroxidase in human parotid and submandibular glands. *J Anat* 127(Pt 1):181-191.

Rivera-Hidalgo F (2003). Smoking and periodontal disease. *Periodontol 2000* 32:50-58.

Ryan CS, Kleinberg I (1995). Bacteria in human mouths involved in the production and utilization of hydrogen peroxide. *Arch Oral Biol* 40:753-763.

Sassone LM, Fidel R, Fidel S, Vieira M, Hirata R Jr (2003a). The influence of organic load on the antimicrobial activity of different concentrations of NaOCl and chlorhexidine in vitro. *Int Endod J* 36:848-852.

Sassone LM, Fidel RA, Fidel SR, Dias M, Hirata RJ (2003b). Antimicrobial activity of different concentrations of NaOCl and chlorhexidine using a contact test. *Braz Dent J* 14:99-102.

Sato EF, Choudhury T, Nishikawa T, Inoue M (2008). Dynamic aspect of reactive oxygen and nitric oxide in oral cavity. *J Clin Biochem Nutr* 42:8-13.

Schraufstatter IU, Browne K, Harris A, Hyslop PA, Jackson JH, Quehenberger O, *et al.* (1990). Mechanisms of hypochlorite injury of target cells. *J Clin Invest* 85:554-562.

Segal R, Dunford HB, Morrison M (1968). Kinetics of fluoride binding by lactoperoxidase. *Can J Biochem* 46:1471-1474.

Sena NT, Gomes B, Vianna ME, Berber VB, Zaia AA, Ferraz CCR, *et al.* (2006). In vitro antimicrobial activity of sodium hypochlorite and chlorhexidine against selected single-species biofilms. *Int Endod J* 39:878-885.

Shapira L, Wilensky A, Kinane DF (2005). Effect of genetic variability on the inflammatory response to periodontal infection. *J Clin Periodontol* 32(Suppl 6):72-86.

Shenkin JD, Broffitt B, Levy SM, Warren JJ (2004). The association between environmental tobacco smoke and primary tooth caries. *J Public Health Dent* 64:184-186.

Shin K, Tomita M, Lonnerdal B (2000). Identification of lactoperoxidase in mature human milk. *J Nutr Biochem* 11:94-102.

Silva Mendez LS, Allaker RP, Hardie JM, Benjamin N (1999). Antimicrobial effect of acidified nitrite on cariogenic bacteria. *Oral Microbiol Immunol* 14:391-392.

Singer DL, Chatterjee R, Denepitiya L, Kleinberg I (1983). A comparison of the acid-base metabolisms of pooled human dental plaque and salivary sediment. *Arch Oral Biol* 28:29-35.

Sitte N, Von Zglinicki T (2003). Free radical production and antioxidant defense: a primer. In: Aging at the molecular level (biology of aging and Its modulation). Von Zglinicki T, editor. Dordrecht, the Netherlands: Kluwer Academic Publishers, pp. 1-10.

Smalley JW (1994). Pathogenic mechanisms in periodontal disease. *Adv Dent Res* 8:320-328.

Smirnova GV, Oktyabrsky ON (2005). Glutathione in bacteria. *Biochemistry (Moscow)* 70:1199-1211.

Smith AJ, Benjamin N, Weetman DA, Mackenzie D, MacFarlane TW (1999). The microbial generation of nitric oxide in the human oral cavity. *Microb Ecol Health Dis* 11:23-27.

Smith LL (2004). Oxygen, oxysterols, ouabain, and ozone: a cautionary tale. *Free Radic Biol Med* 37:318-324.

Soszynski M, Zavodnik IB, Zavodnik LB, Zylinska L, Bartosz G, Bryszewska M (2002). Hypochlorous acid inhibits glutathione S-conjugate export from human erythrocytes. *Biochim Biophys Acta* 1564:479-486.

Spahn JD (2007). Asthma biomarkers in sputum. *Immunol Allergy Clin North Am* 27:607-622.

Spratt DA, Pratten J, Wilson M, Gulabivala K (2001). An in vitro evaluation of the antimicrobial efficacy of irrigants on biofilms of root canal isolates. *Int Endod J* 34:300-307.

Stewart PS (1994). Biofilm accumulation model that predicts antibiotic resistance of *Pseudomonas aeruginosa* biofilms. Antimicrob *Agents Chemother* 38:1052-1058.

Stewart PS, Raquepas JB (1995). Implications of reaction-diffusion theory for the disinfection of microbial biofilms by reactive antimicrobial agents. *Chem Eng Sci* 50:3099-3104.

Stewart PS, Rayner J, Roe F, Rees WM (2001). Biofilm penetration and disinfection efficacy of alkaline hypochlorite and chlorosulfamates. *J Appli Microbiol* 91:525-532.

Stewart PS, Mukherjee PK, Ghannoum MA (2004). Biofilm antimicrobial resistance. In: Microbial biofilms. Ghannoum M, O'Toole G, editors. Washington, DC; ASM Press, pp. 250-268.

Sun S, Li X, Zhang G, Ma H, Zhang D, Bao Z (2006). Determination of H_2O_2-dependent generation of singlet oxygen from human saliva with a novel chemiluminescence probe. *Biochim Biophys Acta* 1760:440-444.

Suzuki T, Ohshima H (2003). Modification by fluoride, bromide, iodide, thiocyanate and nitrite anions of reaction of a myeloperoxidase-H_2O_2-Cl system with nucleosides. *Chem Pharm Bull* 51:301-304.

Tahboub YR, Galijasevic S, Diamond MP, Abu-Soud HM (2005). Thiocyanate modulates the catalytic activity of mammalian peroxidases. *J Biol Chem* 280:26129-26136.

Takahama U (1993). Hydrogen peroxide-dependent generation of singlet molecular oxygen by human saliva: its detection by chemiluminescence from a cypridina luciferin analog. *Photochem Photobiol* 57:376-379.

Downloaded from jdr.sagepub.com at UNIV OF OKLAHOMA on January 20, 2016 For personal use only. No other uses without permission.

International and American Associations for Dental Research

Tenovuo J (1976). The variation of salivary peroxidase activities in persons of different oral health. *Acta Odontol Scand* 34:163-168.

Tenovuo J (1979). Formation of the bacterial inhibitor, hypothiocyanite ion, by cell-bound lactoperoxidase. *Caries Res* 13:137-143.

Tenovuo J, Larjava H (1984). The protective effect of peroxidase and thiocyanate against hydrogen peroxide toxicity assessed by the uptake of [3H]-thymidine by human gingival fibroblasts cultured in vitro. *Arch Oral Biol* 29:445-451.

Tenovuo J, Makinen KK (1976). Concentration of thiocyanate and ionizable iodine in saliva of smokers and nonsmokers. *J Dent Res* 55:661-663.

Tenovuo J, Söderling E, Sievers G (1985). The peroxidase system in human tears. In: Protides of the biological fluids. Vol. 32. Peters H, editor. Oxford, UK: Pergamon Press, pp. 107-110.

Thibodeau EA, Bowen WH, Marquis RE (1985). pH-dependent fluoride inhibition of peroxidase activity. *J Dent Res* 64:1211-1213.

Thomas EL, Bates KP, Jefferson MM (1980). Hypothiocyanite ion: detection of the antimicrobial agent in human saliva. *J Dent Res* 59:1466-1472.

Thomas EL, Pera KA, Smith KW, Chwang AK (1983). Inhibition of *Streptococcus mutans* by the lactoperoxidase antimicrobial system. *Infect Immun* 39:767-778.

Thomas EL, Jefferson MM, Joyner RE, Cook GS, King CC (1994a). Leukocyte myeloperoxidase and salivary lactoperoxidase: identification and quantitation in human mixed saliva. *J Dent Res* 73:544-555.

Thomas EL, Milligan TW, Joyner RE, Jefferson MM (1994b). Antibacterial activity of hydrogen peroxide and the lactoperoxidase-hydrogen peroxide-thiocyanate system against oral streptococci. *Infect Immun* 62:529-535.

Thomas EL, Bozeman PM, Jefferson MM, King CC (1995). Oxidation of bromide by the human leukocyte enzymes myeloperoxidase and eosinophil peroxidase. Formation of bromamines. *J Biol Chem* 270:2906-2913.

Thongboonkerd V, Luengpailin J, Cao J, Pierce WM, Cai J, Klein JB, *et al.* (2002). Fluoride exposure attenuates expression of *Streptococcus pyogenes* virulence factors. *J Biol Chem* 277:16599-16605; *erratum* in *J Biol Chem* 277:42417, 2002.

Tipton DA, Braxton SD, Dabbous MK (1995a). Effects of a bleaching agent on human gingival fibroblasts. *J Periodontol* 66:7-13.

Tipton DA, Braxton SD, Dabbous MK (1995b). Role of saliva and salivary components as modulators of bleaching agent toxicity to human gingival fibroblasts in vitro. *J Periodontol* 66:766-774.

Tsuchiya M, Asada A, Kasahara E, Sato EF, Shindo M, Inoue M (2002). Smoking a single cigarette rapidly reduces combined concentrations of nitrate and nitrite and concentrations of antioxidants in plasma. *Circulation* 105:1155-1157.

Tsuge K, Kataoka M, Seto Y (2000). Cyanide and thiocyanate levels in blood and saliva of healthy adult volunteers. *J Health Sci* 46:343-350.

Ueda T, Sakamaki K, Kuroki T, Yano I, Nagata S (1997). Molecular cloning and characterization of the chromosomal gene for human lactoperoxidase. *Eur J Biochem* 243:32-41.

Uehara Y, Agematsu K, Kikuchi K, Matsuzaki S, Imai S, Takamoto M, *et al.* (2006). Secretory IgA, salivary peroxidase, and catalase-mediated microbicidal activity during hydrogen peroxide catabolism in viridans streptococci: pathogen coaggregation. *J Infect Dis* 194:98-107.

van Dalen CJ, Whitehouse MW, Winterbourn CC, Kettle AJ (1997). Thiocyanate and chloride as competing substrates for myeloperoxidase. *Biochem J* 327(Pt 2):487-492.

van den Abbeele A, Pourtois M, Courtois P (1992). Fluoride inhibition of scn- and cl-peroxidase activities in whole saliva and of recombinant myeloperoxidase. Influence of pH and hydrogen peroxide concentration. *J Biol Buccale* 20:219-224.

van den Abbeele A, Courtois P, Pourtois M (1995). The influence of different fluoride salts on fluoride-mediated inhibition of peroxidase activity in human saliva. *Arch Oral Biol* 40:695-698.

van der Hoeven JS, Camp PJ (1993). Mixed continuous cultures of *Streptococcus mutans* with *Streptococcus sanguis* or with *Streptococcus oralis* as a model to study the ecological effects of the lactoperoxidase system. *Caries Res* 27:26-30.

van der Vliet A, Eiserich JP, Halliwell B, Cross CE (1997). Formation of reactive nitrogen species during peroxidase-catalyzed oxidation of nitrite. A potential additional mechanism of nitric oxide-dependent toxicity. *J Biol Chem* 272:7617-7625.

Van Dyke TE (2007). Control of inflammation and periodontitis. *Periodontol 2000* 45:158-166.

Van Haeringen NJ, Ensink FTE, Glasius E (1979). The peroxidase-thiocyanate-hydrogenperoxide system in tear fluid and saliva of different species. *Exp Eye Res* 28:343-347.

Van Loveren C (1990). The antimicrobial action of fluoride and its role in caries inhibition. *J Dent Res* 69(Spec Iss):676-681.

Vannini L, Lanciotti R, Baldi D, Guerzoni ME (2004). Interactions between high pressure homogenization and antimicrobial activity of lysozyme and lactoperoxidase. *Int J Food Microbiol* 94:123-135.

Vianna ME, Gomes B, Berber VB, Zaia AA, Ferraz CCR, de Souza-Filho FJ (2004). In vitro evaluation of the antimicrobial activity of chlorhexidine and sodium hypochlorite. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 97:79-84.

Vile GF, Rothwell LA, Kettle AJ (2000). Initiation of rapid, p53-dependent growth arrest in cultured human skin fibroblasts by reactive chlorine species. *Arch Biochem Biophys* 377:122-128.

Vissers MCM, Pullar JM, Hampton MB (1999). Hypochlorous acid causes caspase activation and apoptosis or growth arrest in human endothelial cells. *Biochem J* 344(Pt 2):443-449.

Waddington RJ, Moseley R, Embery G (2000). Reactive oxygen species: a potential role in the pathogenesis of periodontal diseases. *Oral Dis* 6:138-151.

Wang Y (2008). Bulky DNA lesions induced by reactive oxygen species. *Chem Res Toxicol* 21:276-281.

Wang Z, Nicholls SJ, Rodriguez ER, Kummu O, Hoerkkoe S, Barnard J, *et al.* (2007). Protein carbamylation links inflammation, smoking, uremia and atherogenesis. *Nat Med* 13:1176-1184.

Ward PA (1991). Mechanisms of endothelial cell killing by hydrogen peroxide or products of activated neutrophils. *Am J Med* 91(3C):89-94.

Webb BC, Willcox MDP, Thomas CJ, Harty DWS, Knox KW (1995). The effect of sodium-hypochlorite on potential pathogenic traits of *Candida albicans* and other *Candida* species. *Oral Microbiol Immunol* 10:334-341.

Wentworth AD, Jones LH, Wentworth P Jr, Janda KD, Lerner RA (2000). Antibodies have the intrinsic capacity to destroy antigens. *Proc Natl Acad Sci USA* 97:10930-10935.

Wentworth P Jr, McDunn JE, Wentworth AD, Takeuchi C, Nieva J, Jones T, *et al.* (2002). Evidence for antibody-catalyzed ozone formation in bacterial killing and inflammation. *Science* 298:2195-2199.

Weuffen W, Franzke C, Thürkow B (1984). Alimentary ingestion, analysis and biological significance of thiocyanate. *Nährung* 28:341-355.

Wever R, Bakkenist ARJ (1980). The interaction of myeloperoxidase with ligands as studied by EPR. *Biochim Biophys Acta* 612:178-184.

White JW Jr (1975). Relative significance of dietary sources of nitrate and nitrite. *J Agric Food Chem* 23:886-891.

White WE Jr, Pruitt KM, Mansson-Rahemtulla B (1983). Peroxidase-thiocyanate-peroxide antibacterial system does not damage DNA. *Antimicrob Agents Chemother* 23:267-272.

Winniczuk PP, Parish ME (1997). Minimum inhibitory concentrations

Downloaded from jdr.sagepub.com at UNIV OF OKLAHOMA on January 20, 2016 For personal use only. No other uses without permission.

International and American Associations for Dental Research

of antimicrobials against micro-organisms related to citrus juice. *Food Microbiol* 14:373-381.

Winterbourn CC (1995). Toxicity of iron and hydrogen peroxide: the Fenton reaction. *Toxicol Lett* 82/83:969-974.

Winterbourn CC, Kettle AJ (2000). Biomarkers of myeloperoxidase-derived hypochlorous acid. *Free Radic Biol Med* 29:403-409.

Woldring MG (1955). Free amino acids of human saliva: a chromatographic investigation. *J Dent Res* 34:248-256.

Wong-Chong GM, Ghosh RS, Bushey JT, Ebbs SD, Neuhauser EF (2006). Natural sources of cyanide. In: Cyanide in water and soil. Chemistry, risk & management. Dzombak DA, Ghosh RS, Wong-Chong GM, editors. Boca Raton, FL: CRC Press, pp. 25-40.

Wood JL (1975). Biochemistry of thiocyanates. In: Chemistry and biochemistry of thiocyanic acid & its derivatives. New York: Academic Press Inc., pp. 156-221.

Wunder D, Bowen WH (1999). Action of agents on glucosyltransferases from *Streptococcus mutans* in solution and adsorbed to experimental pellicle. *Arch Oral Biol* 44:203-214.

Yesilsoy C, Whitaker E, Cleveland D, Phillips E, Trope M (1995). Antimicrobial and toxic effects of established and potential root-canal irrigants. *J Endod* 21:513-515.

Yu X, Loimaranta V, Lenander-Lumikari M, Wunder D, Bowen WH, Tenuvuo J (2000). Effect of lactoperoxidase system on glucosyltransferase D of *Streptococcus mutans*. *Chin J Dent Res* 3:61-64.

Zgliczynski JM, Stelmaszynska T, Olszowska E, Krawczyk A, Kwasnowska E, Wrobel JT (1983). Peroxidative oxidation of halides catalyzed by myeloperoxidase. Effect of fluoride on halide oxidation. *Acta Biochim Pol* 30:213-222.

Zhu PC (2007). New biocides development. The combined approach of chemistry and microbiology (ACS Symposium Series). New York: Oxford University Press, p. 967.

Downloaded from jdr.sagepub.com at UNIV OF OKLAHOMA on January 20, 2016 For personal use only. No other uses without permission.

International and American Associations for Dental Research