INFORMATION TO USERS

This was produced from a copy of a document sent to us for microfilming. While the most advanced technological means to photograph and reproduce this document have been used, the quality is heavily dependent upon the quality of the material submitted.

The following explanation of techniques is provided to help you understand markings or notations which may appear on this reproduction.

1. The sign or "target" for pages apparently lacking from the document photographed is "Missing Page(s)". If it was possible to obtain the missing page(s) or section, they are spliced into the film along with adjacent pages. This may have necessitated cutting through an image and duplicating adjacent pages to assure you of complete continuity.

2. When an image on the film is obliterated with a round black mark it is an indication that the film inspector noticed either blurred copy because of movement during exposure, or duplicate copy. Unless we meant to delete copyrighted materials that should not have been filmed, you will find a good image of the page in the adjacent frame.

3. When a map, drawing or chart, etc., is part of the material being photographed the photographer has followed a definite method in "sectioning" the material. It is customary to begin filming at the upper left hand corner of a large sheet and to continue from left to right in equal sections with small overlaps. If necessary, sectioning is continued again—beginning below the first row and continuing on until complete.

4. For any illustrations that cannot be reproduced satisfactorily by xerography, photographic prints can be purchased at additional cost and tipped into your xerographic copy. Requests can be made to our Dissertations Customer Services Department.

5. Some pages in any document may have indistinct print. In all cases we have filmed the best available copy.

University
Microfilms
International

300 N. ZEEB ROAD, ANN ARBOR, MI 48106
18 BEDFORD ROW, LONDON WC1R 4EJ, ENGLAND

Downloaded from SHAREOK. Persistent Handle URL: https://hdl.handle.net/11244/4765

8027513

DORAISWAMY, IDA GRACE PANCHARATNAM

*SEMI *SIMPLICITY OF *MODULES OVER *SIMPLE *RINGS

The University of Oklahoma PH.D. 1980

University Microfilms International 300 N. Zeeb Road, Ann Arbor, MI 48106 18 Bedford Row, London WC1R 4EJ, England

THE UNIVERSITY OF OKLAHOMA

GRADUATE COLLEGE

*SEMI *SIMPLICITY OF *MODULES OVER *SIMPLE *RINGS

A DISSERTATION

SUBMITTED TO THE GRADUATE FACULTY

in partial fulfillment of the requirements for the

degree of

DOCTOR OF PHILOSOPHY

BY

IDA GRACE DORAISWAMY

Norman, Oklahoma

1980

*SEMI *SIMPLICITY OF *MODULES OVER *SIMPLE *RINGS

APPROVED BY

Chairman

Leonard R. Rubin

Harold Hunche

Roger Alperin

Kevin A. Hirsch

DISSERTATION COMMITTEE

ACKNOWLEDGEMENTS

I express my deepest gratitude to Professor Andy R. Magid for accepting me as his student, helping me acquire as much knowledge as I could during this period and, more importantly, for his inexhaustible patience and guidance during the preparation of this thesis.

I express my appreciation to the present members of my dissertation committee: Dr. Harold Huneke, Dr. Leonard Rubin, Dr. Roger Alperin and Dr. Kevin Grasse, and the past members, Dr. Dale Umbach and Dr. Robert Morris. I especially thank Dr. Huneke who made this educational experience possible and Dr. Morris for his encouragement and support when they were needed most.

I thank Trish Abolins for the splendid job of typing.

My gratitude for the silent inspiration of my parents is not among the least.

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
TABLE OF CONTENTS.	iv
INTRODUCTION	v
Chapter	
I. THE CATEGORY OF R-*MODULES	1
II. *SEMI *SIMPLICITY.	22
III. k[SL_n]-*MODULES WITH SL_n-ACTION	39
BIBLIOGRAPHY	

INTRODUCTION

This thesis is a study of the structure of a *simple *ring and its category of *modules.

We state the definitions needed for a description of the thesis.

k is an algebraically closed field. G is a linear algebraic group over k . A *ring is a commutative Noetherian k-algebra which is a rational G-module such that G acts by k-algebra automorphisms, [M]. A *module M over a *ring is an R-module and a rational G-module such that $g(rm) = (gr)(gm)$ for g in G , r in R and m in M , [M]. An *ideal of a *ring is an ideal which is a sub*module of R , [M]. A *ring R is *simple if the only *ideals of R are zero and R , [M].

We define an R-*module M to be *simple if the only G-stable R-sub*modules of M are the zero *module and the *module itself. An R-*module M is *semi*simple if it is a direct sum of *simple R-sub*modules of itself. A *ring R is *semi*simple if it is *semi-*simple as an R-*module.

In the first chapter we establish the category of R-*modules

and R-*module morphisms. In order for an R-module M to be an R-*module, we need to define a rational G-action on M. With an appropriate definition of G-action we prove that the direct sum of a family of R-*modules is an R-*module, the tensor product $M \otimes_R N$ of R-*modules M and N is an R-*module, $\mathrm{Hom}_R(M,N)$ is an R-*module if M is a finitely generated R-*module and M/N is an R-*module if N is an R-sub*module of M. By considering only the morphisms that preserve G-action we establish a subcategory which is an abelian category. Since direct limit exists in this category, every R-*module is the direct limit of its family of finitely generated R-sub*modules. These are useful results for the development of the theory.

The second chapter contains the theory of *semi*simplicity. Three equivalent conditions for defining *semi*simplicity are established. The conditions are similar to those of semisimplicity in the category of R-modules. The techniques for proving the equivalence are also similar except for establishing that every R-*module M contains a *simple sub*module if every sub*module N of M is a direct summand. This is because not every principal submodule of M is a sub*module.

We also examine the properties of *modules over *simple *rings when certain restrictions are imposed on the algebraic group. The main theorem is that when G is a connected linear algebraic group and R is a *simple *ring then every finitely generated R-*module M is R-torsion-free. Thus, a *simple *ring is an integral domain if G is a connected linear algebraic group. We establish this result by proving that every associated prime of M is G-stable and therefore an *ideal.

R being *simple every associated prime reduces to the zero *ideal proving that M is torsion-free. If R is a finitely generated k-algebra that is *simple then M is R-projective. This is shown by proving that M_M is R_M-free for every maximal ideal M of M .

If further G is a connected linearly reductive algebraic group and R a *simple *ring then every finitely generated R-*module is *semi*simple and, therefore, R-projective. Under the same conditions for G every nonzero R-*module is *semi*simple and, therefore, R-projective.

If G is a linear algebraic group then k[G] with appropriate G-action defined on it is a *simple *ring. In Chapter Three, we examine $k[SL_n]$-*modules with SL_n-action for $n \geq 2$ when k is algebraically closed and the characteristic is zero. We prove that every *simple $k[SL_n]$-*module is R-isomorphic to $k[SL_n]$. This isomorphism preserves SL_n-action as well. Consequently, every $k[SL_n]$-*module is R-isomorphic to either $R^{(n)}$, $n < \infty$, or $R^{(\chi)}$, the isomorphism preserving SL_n-action as well. We call such an isomorphism (R-G)-isomorphism. We establish the isomorphism by the following sequence of arguments. It is a fact that for any linear algebraic group G , if $G \to GL(V)$ is a faithful representation of G in V then $k[G] = k[V + V^*]$, V^* being the dual of V , $[M^*]$. If W is a *simple G-module then W is the homomorphic image of $\bigoplus_{i=1}^{n} (V \oplus V^*)^{\otimes_k d_i}$ for $d_i > 0$, the homomorphism being that of G-modules. In particular, if $G = SL_n$ then $k[SL_n] = k[GL_n]/(1 - D)$ where $k[GL_n] = k[x_{11}, x_{12}, \ldots, x_{nn}, 1/D]$ and

$D = D(x_{11},\ldots,x_{nn})$ is the determinant form. We define R_n to be an SL_n-module isomorphic to $\langle x_{11},x_{22},\ldots,x_{n1}\rangle$. Then W is the homomorphic image of $\bigoplus_{i=1}^{m} (R_n \oplus R_n^*)^{\otimes_k d_i}$, the homomorphism being that of SL_n-modules. $R \otimes_k R_n \cong_{RG} R^n$. This leads to the isomorphism $M \cong_{RG} R$ if M is *simple. Therefore every $k[SL_n]$-*module is (R-G)-isomorphic to $R^{(n)}$ or $R^{(\chi)}$. The existence of such an isomorphism follows from a more general theorem [CPM]. But we construct an explicit form of the isomorphism in this thesis.

*SEMI *SIMPLICITY OF *MODULES OVER *SIMPLE *RINGS

CHAPTER I

THE CATEGORY OF R-*MODULES

Starting with the basic definitions of this theory, this chapter establishes the category of R-*modules and R-*module *homomorphisms. Properties of R-*modules and R-*module *homomorphisms needed for the development of this theory are demonstrated. That this category contains a subcategory which is Abelian is also shown.

Throughout this thesis k is an algebraically closed field and G a linear algebraic group over k. For any algebraic set V, $k[V]$ denotes its coordinate ring. If R is a commutative ring then Mod_R is the category of R-modules and R-module homomorphisms. I is an indexing set.

Definition 1.1 A finite dimensional vector space V over k with G-action is a G-module if the induced homomorphism $G \to GL(V)$ is a homomorphism of algebraic groups over k. [M]

That is, if $\varphi : G \to GL(V)$ is the induced homomorphism then φ is a homomorphism of groups and is a k-morphism. k-morphism means that if $f \in k[GL(V)]$ then $f \circ \varphi \in k[G]$. [F]

Definition 1.2 A vector space W over k with G-action is a rational G-module if W is a union of finite dimensional G-modules in the above sense. [M]

Definition 1.3 A *ring R is a commutative Noetherian k-algebra which is a rational G-module such that G acts by k-algebra automorphisms. [M]

G acts rationally on R in the following sense. If $v \in R$ and $S_v = \langle gv \mid g \in G \rangle$, the vector space spanned by gv for all $g \in G$, then S_v is finite dimensional over k and the induced homomorphism $G \to GL(S_v)$ is a homomorphism of algebraic groups and $R = \bigcup_{v \in R} S_v$.

Definition 1.4 (Notation) $\langle gv \mid g \in G \rangle$ denotes the vector space generated over k by gv , for all $g \in G$. $\langle v_1, v_2, \ldots, v_n \rangle$ denotes the vector space generated over k by $v_1, v_2, \ldots, v_n$.

Definition 1.5 A *module M over a *ring R is an R-module and a rational G-module such that $g(rm) = (gr)(gm)$, for all $g \in G$, $r \in R$, $m \in M$. [M]

A *module M over a *ring R is said to be an R-*module.

A *module M over a *ring R is a rational G-module in the following sense. If $m \in M$ and $V_m = \langle gm \mid g \in G \rangle$ then V_m is finite dimensional over k and the induced homomorphism $G \to GL(V_m)$ is a homomorphism of algebraic groups. Moreover $M = \bigcup_{m \in M} V_m$.

Definition 1.6 An (R-G)-*module homomorphism of *modules over a *ring R is an R-module homomorphism preserving G-action.

Definition 1.7 (Notation) If $g \in G$ and M an R-*module then g_M denotes the G-action of g on M .

Definition 1.8 For each pair (M,N) of R-*modules define a set, $^*\mathrm{Hom}(M,N)$, of morphisms of M into N where $^*\mathrm{Hom}(M,N) = \{f \in \mathrm{Hom}_R(M,N) \mid \langle g_N f g_M^{-1} \mid g \in G\rangle$ is finite dimensional over k$\}$. f is said to be a *homomorphism if $f \in {}^*\mathrm{Hom}(M,N)$.

Proposition 1.9 The category whose objects are R-*modules and whose morphisms are *homomorphisms of R-*modules, as defined in 1.8, is a category and denote it by $^*\mathrm{Mod}_R$.

Proof: We first define composition of *homomorphisms, establishing that the composite map so defined is a *homomorphism.

For each triple (M,N,L) of R-*modules, define a map $^*\mathrm{Hom}(M,N) \times {}^*\mathrm{Hom}(N,L) \to {}^*\mathrm{Hom}(M,L)$ by $(u,v) = v \cdot u$ where $u \in {}^*\mathrm{Hom}(M,N)$, $v \in {}^*\mathrm{Hom}(N,L)$ and $\cdot$ is the usual composition of maps. Denote $v \cdot u$ by vu . By the definition of a *homomorphism, $\langle g_N u g_M^{-1} \mid g \in G\rangle$ and $\langle g_L v g_N^{-1} \mid g \in G\rangle$ are finite dimensional vector spaces over k . Let $\langle g_N u g_M^{-1} \mid g \in G\rangle = \langle f_1, f_2, \ldots, f_m \mid f_i \in {}^*\mathrm{Hom}(M,N)\rangle$ and $\langle g_L v g_N^{-1} \mid g \in G\rangle = \langle h_1, h_2, \ldots, h_n \mid h_i \in {}^*\mathrm{Hom}(N,L)\rangle$. Then $\langle g_L vu g_N^{-1} \mid g \in G\rangle \subseteq \langle h_i f_j \mid 1 \le i \le n \text{ , } 1 \le j \le m\rangle$ which is finite dimensional over k . Moreover $vu \in \mathrm{Hom}_R(M,L)$. Therefore $vu \in {}^*\mathrm{Hom}(M,L)$.

(i) Let M_1, M_2, M_3, M_4 be R-*modules and $f_1 \in {}^*\mathrm{Hom}(M_1,M_2)$, $f_2 \in {}^*\mathrm{Hom}(M_2,M_3)$, $f_3 \in {}^*\mathrm{Hom}(M_3,M_4)$. $f_3(f_2f_1) = (f_3f_2)f_1$ as R-module homomorphisms. Moreover, $f_3(f_2f_1), (f_3f_2)f_1 \in {}^*\mathrm{Hom}(M_1,M_4)$.

Thus composition of *homorphisms is associative.

(ii) For any R-*module M, let 1_M be the identity map of M onto M. Then $1_M \in \mathrm{Hom}_R(M,M)$. Moreover, $\langle g 1_M g^{-1}_M \mid g \in G \rangle \cong k$ and therefore finite dimensional over k. Thus, $1_M \in {}^*\mathrm{Hom}(M,M)$. Let $f \in {}^*\mathrm{Hom}(M,N)$ and $h \in {}^*\mathrm{Hom}(N,M)$ for any R-*module N. $f1_M = f$ and $1_M h = h$ as R-module homomorphisms. Also, $f1_M, f \in {}^*\mathrm{Hom}(M,N)$ and $1_M h, h \in {}^*\mathrm{Hom}(N,M)$. Therefore, $f1_M = f$ and $1_M h = h$ as R-*module *homomorphisms. Thus, 1_M is a left identity in ${}^*\mathrm{Hom}(M,N)$ and a right identity in ${}^*\mathrm{Hom}(N,M)$.

(iii) Let the pairs (M_1,N_1) and (M_2,N_2) of R-*modules be distinct. If $f \in {}^*\mathrm{Hom}(M_1,N_1) \cap {}^*\mathrm{Hom}(M_2,N_2)$ with $f \neq 0$, then $f \in \mathrm{Hom}_R(M_1,N_1) \cap \mathrm{Hom}_R(M_2,N_2)$. This implies that $M_1 = M_2$ and $N_1 = N_2$.

(i), (ii) and (iii) establish the proposition.

For an R-module M to be an R-*module, M should be a rational G-module and the G-action on M should satisfy the condition $g(rm) = (gr)(gm)$ for all $g \in G$, $r \in R$ and $m \in M$. We show that these two conditions are satisfied whenever it is necessary to establish that an R-module is an R-*module.

Proposition 1.10 If $\{M_i \mid 1 \leq i \leq n\}$ is a finite family of R-*modules, then $\bigoplus_i M_i$ is an R-*module.

Proof: $M = \bigoplus_i M_i$ is an R-module.

(i) Let G act on M as follows: $g(r((m_i)_i)) = g((rm_i)_i) = (g(rm_i))_i = ((gr)(gm_i))_i = (gr)((gm_i)_i) = (gr)(g((m_i)_i))$ for all $g \in G$, $r \in R$, $m_i \in M_i$, $1 \leq i \leq n$.

(ii) Let $x = (m_i)_i \in M$ with $m_i \in M_i$. For each m_i $\exists$ a G-stable

finite dimensional subspace V_{m_i} of M_i, over k, such that $m_i \in V_{m_i}$ and the homomorphism $\mu_i : G \to GL(V_{m_i})$ is a homomorphism of algebraic groups over k. Let $V_x = \oplus_i V_{m_i}$, $1 \leq i \leq n$. V_x is finite dimensional over k and is G-stable. The induced homomorphism $\mu : G \to GL(V_x)$ is defined by $\mu(g) = (\mu_i(g))_i$ for all $g \in G$. If $g_1, g_2 \in G$ then $\mu(g_1g_2) = (\mu_i(g_1g_2))_i = (\mu_i(g_1)\mu_i(g_2))_i = ((\mu_i(g_1))_i)((\mu_i(g_2))_i) = \mu(g_1)\mu(g_2)$. This proves that μ is a homomorphism of groups.

Now it is sufficient to show that if $\varphi \in k[GL(V_x)]$ then $\varphi \cdot \mu \in k[G]$. The homomorphism $\mu : G \to GL(V_x)$ can be factored as $G \xrightarrow{\lambda} \oplus_j GL(V_{m_j}) \xrightarrow{i} GL(V_x)$ where λ is defined by $\lambda(g) = (\mu_j(g))_j$ for all $g \in G$ and i is the inclusion map. Therefore, $i\lambda = \mu$. $k[\oplus_j GL(V_{m_j})] = \otimes_k k[GL(V_{m_j})]$. If $\varphi \in k[\oplus_j GL(V_{m_j})]$ then $\varphi = \Sigma_\ell (\otimes_j f_{\ell j})$ where $f_{\ell j} \in k[GL(V_{m_j})]$ for all j. But $\lambda : G \to \oplus_j GL(V_{m_j})$ and $\varphi \in \otimes_k k[GL(V_{m_j})]$. Therefore, $\varphi\lambda = \Sigma_\ell(\Sigma_j f_{\ell j}\mu_j) \in k[G]$ since $f_{\ell j}\mu_j \in k[G]$ for all ℓ, j. This proves that $\mu : G \to GL(V_x)$ is a k-morphism and therefore $\mu : G \to GL(V_x)$ is a homomorphism of algebraic groups.

If $x \in M$ then V_x is a G-module and $M = \bigcup_{x \in M} V_x$. That is, M is a rational G-module. (i) and (ii) establish that $M \in {}^*\mathrm{Mod}_R$. This completes the proof of Proposition 1.10.

Proposition 1.11 If $\{M_i \mid i \in I\}$ is an infinite family of R-*modules then $\oplus_{i \in I} M_i$ is an R-*module.

Proof: $\oplus M_i$ is an R-module. If $x = (x_i)_i \in M$ where $M = \oplus_i M_i$ then all but a finite number of x_i terms are zero. A G-action defined on M as in Prop. 1.10 satisfies the required condition for G-action. $V_x = \oplus V_{x_i}$ where all but a finite number of G-modules V_{x_i} are zero modules. This forces V_x to be finite dimensional over k. V_x is also G-stable. Therefore the argument that the homomorphism μ : $G \to GL(V_x)$ is a k-morphism is the same as that in Prop. 1.10. $M = \bigcup_{x \in M} V_x$. Therefore M is an R-*module. This completes the proof of Prop. 1.11.

Proposition 1.12 Let M be a finitely generated R-*module and N an R-*module. If a G-action on $\mathrm{Hom}_R(M,N)$ is defined by $g \circ f = gfg^{-1}$ for all $g \in G$, $f \in \mathrm{Hom}_R(M,N)$ then $\mathrm{Hom}_R(M,N)$ is an R-*module. Moreover, $\mathrm{Hom}_R(M,N) = {}^*\mathrm{Hom}(M,N)$.

Proof: $\mathrm{Hom}_R(M,N)$ is an R-module. If $f \in \mathrm{Hom}_R(M,N)$ let $V_f = \langle gfg^{-1} \mid g \in G \rangle$. It is sufficient to prove that V_f is finite dimensional over k and the homomorphism $G \to GL(V_f)$ is a k-morphism.

Let M be generated by $m_1,\ldots,m_s$, with $m_i \in M$. Then $V_{m_i} = \langle gm_i \mid g \in G \rangle$ is finite dimensional over k. Let $V_{m_i} = \langle m_i, g_{2i}m_i, \ldots, g_{pi}m_i \rangle$ with $g_{ji} \in G$, $1 \le i \le s$ and $2 \le j \le p$. So also let $V_{f(m_i)} = \langle f(m_i), h_{2i}f(m_i), \ldots, h_{qi}f(m_i) \rangle$ with $h_{ji} \in G$, $1 \le i \le s$ and $2 \le j \le q$. $V_{f(g_{ji}m_i)} = \langle f(g_{ji}m_i), d_{2i}f(g_{ji}m_i), \ldots, d_{ri}f(g_{ji}m_i) \rangle$ with $d_{\ell i} \in G$, $1 \le i \le s$, $2 \le j \le p$ and $2 \le \ell \le r$. We now prove that $\langle gfg^{-1} \mid g \in G \rangle$ is contained in the span of f, $h_{n_i}f$, $d_{\ell i}f_{ji}$ with $1 \le i \le s$, $2 \le n \le q$, $2 \le \ell \le r$

and $2 \le j \le p$ and therefore is finite dimensional over k.

If $g \in G$ then $gfg^{-1}(m_i) = \sum_{j=2}^{p} \lambda_{ji}(g^{-1})gf(g_{ji}m_i)$ with $\lambda_{ji} \in k[G]$ since M is a rational G-module. $\sum_{j=2}^{p} \lambda_{ji}(g^{-1})gf(g_{ji}m_i) = \sum_{j=2}^{p} \lambda_{ji}(g^{-1})(\sum_{\ell=2}^{r} \mu_{\ell i}(g)d_{\ell i}f(g_{ji}m_i))$ with $\mu_{\ell i} \in k[G]$ since N is a rational G-module and $f(g_{\ell i}m_i) \in N$ for all i,j. Therefore

$gfg^{-1}(m_i) = \sum_{j=2}^{p} \sum_{\ell=2}^{r} \lambda_{ji}(g^{-1})\mu_{\ell i}(g)d_{\ell i}f(g_{ji}m_i)$ with $\lambda_{ji},\mu_{\ell i}$ $k[G]$, $1 \le i \le s$, $2 \le j \le p$, $2 \le \ell \le r$. Thus, V_f is contained in the span of f, $h_{n_i}f$, $d_{\ell i}fg_{ji}$ for all i,j,ℓ,n and the induced homomorphism $G \to GL(V_f)$ is a k-morphism. Moreover, $Hom_R(M,N) = \bigcup_{f \in Hom_R(M,N)} V_f$ and therefore a rational G-module.

Thus, $Hom_R(M,N)$ is an R-*module. Moreover, $^*Hom(M,N) \subseteq Hom_R(M,N)$. If $f \in Hom_R(M,N)$ then $V_f = \langle gfg^{-1} \mid g \in G \rangle$ is finite dimensional over k. Therefore, $f \in {}^*Hom(M,N)$. That is, $Hom_R(M,N) \subseteq {}^*Hom(M,N)$. Then $^*Hom(M,N) = Hom_R(M,N)$. This completes the proof of Prop. 1.12.

Proposition 1.13 Let M,N be R-*modules. If a G-action on $^*Hom(M,N)$ is defined by $g \circ f = gfg^{-1}$ for all $g \in G$, $f \in {}^*Hom(M,N)$ then $^*Hom(M,N)$ is an R-*module.

Proof: We first prove that $^*Hom(M,N)$ is closed under addition and $R^*Hom(M,N) \subseteq {}^*Hom(M,N)$, thus establishing that $^*Hom(M,N)$ is an R-module.

(i) Let $f_1,f_2 \in {}^*Hom(M,N)$.

$\langle g(f_1 + f_2)g^{-1} \mid g \in G\rangle \subseteq \langle gf_1g^{-1} \mid g \in G\rangle + \langle gf_2g^{-1} \mid g \in G\rangle$ which is finite dimensional over k. Therefore, $f_1 + f_2 \in {}^*\mathrm{Hom}(M,N)$.

(ii) Let $r \in R$, $f \in {}^*\mathrm{Hom}(M,N)$. R is a rational G-module. Therefore, let $\langle gr \mid g \in G\rangle = \langle r_1, r_2, \ldots, r_m\rangle$. $\langle gfg^{-1} \mid g \in G\rangle$ is finite dimensional over k by the definition of ${}^*\mathrm{Hom}(M,N)$. Let $\langle gfg^{-1} \mid g \in G\rangle = \langle f_1, f_2, \ldots, f_m\rangle$. Then $\langle g(rf)g^{-1} \mid g \in G\rangle = \langle (gr)(gfg^{-1}) \mid g \in G\rangle \subseteq \langle r_i f_j \mid 1 \le i \le n,\ 1 \le j \le m\rangle$ which is finite dimensional over k. Therefore, $rf \in {}^*\mathrm{Hom}(M,N)$.

(i) and (ii) imply that ${}^*\mathrm{Hom}(M,N)$ is an R-module.

$V_f = \langle gfg^{-1} \mid g \in G\rangle$ for $f \in {}^*\mathrm{Hom}(M,N)$ is finite dimensional over k. Therefore the argument in Prop. 1.12 can be modified to prove that the induced homomorphism $G \to GL(V_f)$ is a k-morphism. Moreover, ${}^*\mathrm{Hom}(M,N) = \bigcup_{f \in {}^*\mathrm{Hom}(M,N)} V_f$. Then ${}^*\mathrm{Hom}(M,N)$ is a rational G-module and therefore an R-*module. This completes the proof of Proposition 1.13.

Proposition 1.14 Let M and M' be R-*modules. Then $M \otimes_R M'$ is an R-*module.

Proof: $M \otimes_R M'$ is an R-module.

(i) A G-action on $M \otimes_R M'$ is defined by $g(m \otimes m') = (gm) \otimes (gm')$ for all $m \in M$, $m' \in M'$ and $g \in G$. If $r \in R$, $m \in M$, $g \in G$ then $g(r(m \otimes m')) = g(rm \otimes m') = g(rm \otimes m') = g(rm) \otimes gm' = ((gr)(gm)) \otimes gm' = (gr)(gm \otimes gm') = (gr)g(m \otimes m')$. This satisfies the requirement for G-action.

(ii) Any element in $M \otimes_k M'$ is of the form $\sum_{i=1}^{n} m_i \otimes m_i'$ with $m_i \in M$ and $m_i' \in M'$ for all i. Let $V_{m_i} = \langle gm_i \mid g \in G \rangle$ and $V_{m_i'} = \langle gm_i' \mid g \in G \rangle$. Since M and M' are R-*modules, V_{m_i} and $V_{m_i'}$ are G-modules. Let $\{\alpha_1,\ldots,\alpha_n\}$ and $\{\beta_1,\beta_2,\ldots,\beta_m\}$ be k-bases for V_{m_i} and $V_{m_i'}$, respectively. Then $\{\alpha_i \otimes \beta_j \mid 1 \le i \le n, 1 \le j \le m\}$ is a k-basis for $V_{m_i} \otimes_k V_{m_i'}$. If $g \in G$, then $g\alpha_i = \sum_{j=1}^{n} a_{ij}(g)\alpha_j$ with $a_{ij} \in k[G]$ for all j. $g\beta_j = \sum_{\ell=1}^{m} b_{j\ell}(g)\beta_\ell$ with $b_{j\ell} \in k[G]$ for all ℓ. $g(\alpha_i \otimes \beta_j) = (g\alpha_i) \otimes (g\beta_j) = (\sum_{j=1}^{n} a_{ij}(g)\alpha_j) \otimes (\sum_{\ell=1}^{m} b_{j\ell}(g)\beta_\ell) =$

$$\sum_{j=1}^{n}\sum_{\ell=1}^{m} (a_{ij}(g)\alpha_j \otimes b_{j\ell}(g)\beta_\ell) = \sum_{j=1}^{n}\sum_{\ell=1}^{m} (a_{ij}(g) \otimes b_{j\ell}(g))(\alpha_j \otimes \beta_\ell)$$ with

$a_{ij} \otimes b_{j\ell} \in k[G] \otimes_k k[G] = k[G \times G]$. This establishes that the induced homomorphism $G \to GL(V_{m_i} \otimes_k V_{m_i'})$ is a k-morphism. Therefore, $V_{m_i} \otimes_k V_{m_i'}$ is a rational G-module. So also is $\sum_i V_{m_i} \otimes_k V_{m_i'}$. Let $S_x = \sum_i V_{m_i} \otimes_k V_{m_i'}$. Then $M \otimes_k M' = \bigcup_{x \in M \otimes_k M'} S_x$ is a rational G-module. $M \otimes_k M' \to M \otimes_R M'$ is a G-module surjection. So by Lemma 1.15, $M \otimes_R M'$ is a rational G-module.

(i) and (ii) imply that $M \otimes_R M'$ is an R-*module. This completes the proof of Prop. 1.14.

Lemma 1.15 Let W and V be finite dimensional G-modules such that $W \subseteq V$. Then V/W is a G-module.

Proof: Let $x_1, x_2, \ldots, x_m$ be a basis of W and $x_1, x_2, \ldots, x_m, x_{m+1}, \ldots, x_n$ a basis of V. The homomorphism $\mu : G \to GL(V)$ induced by the G-action on V is a k-morphism. That is, if $g \in G$ then

$$gx_i = \sum_{j=1}^{n} \alpha_{ji}(g)x_j \quad \text{with} \quad \alpha_{ji} \in k[G] .$$

(i) V/W is a finite dimensional vector space over k with $x_{m+1} + W, \ldots, x_n + W$ as basis.

(ii) Let $y_i = x_i + W$, $m + 1 \leq i \leq n$. A G-action on V/W is defined as $gy_i = gx_i + W$ for $m + 1 \leq i \leq n$ and $gx_i + W = \sum_{j=1}^{n} \alpha_{ji}(g)y_j$. This implies that the homomorphism $\mu : G \to GL(V/W)$ is a k-morphism.

(i) and (ii) establish that V/W is a G-module. This completes the proof of Lemma 1.15.

Proposition 1.16 If N is an R-sub*module of an R-*module M, then M/N is an R-*module.

Proof: M/N is an R-module.

(i) A G-action on M/N is defined as follows. If $g \in G$ and $\bar{x}$ is the canonical image of $x \in M$ in M/N then let $g\bar{x} = \overline{gx}$. If $r \in R$ then $g(r\bar{x}) = g(\overline{rx}) = \overline{g(rx)} = \overline{(gr)(gx)} = (gr)\overline{gx} = (gr)g\bar{x}$.

(ii) If $x \in M$ then let $V_x = \langle gx \mid g \in G \rangle$ and $V_{\bar{x}} = (V_x + N)/N = V_x/V_x \cap N \subseteq M/N$. The induced homomorphism $\mu : G \to GL(V_x)$ is a k-morphism and $V_x \cap N$ is a G-submodule of V_x. By Lemma 1.15, $V_{\bar{x}}$ is a G-module. $M/N = \bigcup_{\bar{x} \in M/N} V_{\bar{x}}$ is a rational G-module.

(i) and (ii) establish that M/N is an R-*module. This completes the proof of Prop. 1.16.

Next we show that the category whose objects are R-*modules and whose morphisms are (R-G)-*module homomorphisms as defined in 1.6 (written as (R-G)-homomorphism) is a category. It is a subcategory of $^*\mathrm{Mod}_R$. We denote it by Mod(R-G). We also establish some properties of Mod(R-G) that make Mod(R-G) an abelian category. We prove that the direct limit of a direct system of R-*modules over a directed set exist in this category. Finally we prove that every R-*module is the direct limit of its family of finitely generated sub*modules.

Definition 1.17 If M and M' are R-*modules then let $\mathrm{Hom}_{RG}(M,M')$ denote the set of (R-G)-homomorphisms from M to M'.

Proposition 1.18 The category whose objects are R-*modules and whose morphisms are (R-G)-homomorphisms is a subcategory of $^*\mathrm{Mod}_R$.

Proof: (R-G)-homomorphisms of R-*modules are R-*module *homomorphisms. Therefore for every pair of objects (M,N) of $^*\mathrm{Mod}_R$, $\mathrm{Hom}_{RG}(M,N) \subseteq {}^*\mathrm{Hom}(M,N)$. Moreover, for any R-*module M, the identity morphism 1_M, defined in Prop. 1.9, preserves G-action and, therefore, is an (R-G)-homomorphism. If M,N,L are R-*modules and $f : M \to N$, $g : N \to L$ are (R-G)-homomorphisms then the composite map gf preserves G-action and therefore is an (R-G)-homomorphism. Thus the category whose objects are R-*modules and whose morphisms are (R-G)-homomorphisms is a subcategory of $^*\mathrm{Mod}_R$. This completes the proof of Prop. 1.18.

Proposition 1.19 If $A \in \mathrm{Mod}(R\text{-}G)$ then $^*\mathrm{Hom}(A,_)$ and $A \underset{R}{\otimes} __$ are functors from Mod(R-G) to Mod(R-G). The ordered pair $^*\mathrm{Hom}(A,_)$, $A \underset{R}{\otimes} __$ is an adjoint pair. That is, if $B,C \in \mathrm{Mod}(R\text{-}G)$ there exists an isomorphism $\Phi : \mathrm{Hom}_{RG}(B, {}^*\mathrm{Hom}(A,C)) \to \mathrm{Hom}_{RG}(A \underset{R}{\otimes} B, C)$ which is

natural in each variable.

Proof: (i) Let $B \in \text{Mod}(R\text{-}G)$. Then ${}^*\text{Hom}(A,B) \in \text{Mod}(R\text{-}G)$ by Prop. 1.13.

(ii) Let $f : B \to C$ be in $\text{Mod}(R\text{-}G)$. Then ${}^*\text{Hom}(A,_)(f) : {}^*\text{Hom}(A,B) \to {}^*\text{Hom}(A,C)$ is defined by $\varphi \mapsto f \circ \varphi$ for all $\varphi \in {}^*\text{Hom}(A,B)$. $f \in \text{Hom}_{RG}(B,C) \subseteq {}^*\text{Hom}(B,C)$ and $\varphi \in {}^*\text{Hom}(A,B)$. Therefore, $f \circ \varphi \in {}^*\text{Hom}(A,C)$. If $r \in R$ then $r\varphi \mapsto f \circ r\varphi$. $f \circ r\varphi(x) = f(r\varphi(x)) = rf(\varphi(x)) = rf \circ \varphi(x)$ for all $x \in A$. Therefore ${}^*\text{Hom}(A,_)(f) \in \text{Hom}_R({}^*\text{Hom}(A,B), {}^*\text{Hom}(A,C))$. If $g \in G$ then $g \circ \varphi \mapsto f \circ (g \circ \varphi) = f \circ (g\varphi g^{-1}) = gf\varphi g^{-1}$ since $f \in \text{Hom}_{RG}(B,C)$. Therefore, $f \circ (g \circ \varphi) = g \circ ({}^*\text{Hom}(A,_)(f)(\varphi))$. That is, ${}^*\text{Hom}(A,_)(f) : {}^*\text{Hom}(A,B) \to {}^*\text{Hom}(A,C)$ is in $\text{Mod}(R\text{-}G)$.

(iii) Let $M \xrightarrow{h} N \xrightarrow{\varphi} P$ be in $\text{Mod}(R\text{-}G)$. Then ${}^*\text{Hom}(A,_)(\varphi h) : {}^*\text{Hom}(A,M) \to {}^*\text{Hom}(A,P)$ is defined by $f \mapsto \varphi h f$ and $\varphi h f : ({}^*\text{Hom}(A,_)(\varphi) \circ {}^*\text{Hom}(A,_)(h))(f)$.

(iv) ${}^*\text{Hom}(A,_)(1_A) : {}^*\text{Hom}(A,A) \to {}^*\text{Hom}(A,A)$ is defined by $\varphi \mapsto 1_A\varphi = \varphi$. This implies that ${}^*\text{Hom}(A,_)(1_A) = 1_{{}^*\text{Hom}(A,A)}$.

(i), (ii), (iii), (iv) prove that ${}^*\text{Hom}(A,_)$ is a functor from $\text{Mod}(R\text{-}G)$ to $\text{Mod}(R\text{-}G)$.

Next we establish that $A \underset{R}{\otimes} _$ is a functor from $\text{Mod}(R\text{-}G)$ to $\text{Mod}(R\text{-}G)$.

$A \underset{R}{\otimes} _$ is a functor from $\text{Mod}(R\text{-}G)$ to the category of abelian groups. Therefore it is sufficient to prove that if $B \in \text{Mod}(R\text{-}G)$ then $A \underset{R}{\otimes} B$ is in $\text{Mod}(R\text{-}G)$ and if $f : B \to C$ is in $\text{Mod}(R\text{-}G)$ then $(A \underset{R}{\otimes} _)(f) : A \underset{R}{\otimes} B \to A \underset{R}{\otimes} C$ defined by $\sum_i (x_i \otimes y_i) \mapsto \sum_i (x_i \otimes f(y_i))$

is in $\mathrm{Mod}(R\text{-}G)$.

(i) By Prop. 1.14, if $B \in \mathrm{Mod}(R\text{-}G)$ then $A \underset{R}{\otimes} B$ is in $\mathrm{Mod}(R\text{-}G)$.

(ii) $(A \underset{R}{\otimes} _)(f)(r(\Sigma_i x_i \otimes y_i)) = \Sigma_i (rx_i \otimes f(y_i)) = r(\Sigma_i x_i \otimes f(y_i)) =$

$r(A \otimes _)(f)(\Sigma_i (x_i \otimes y_i))$. Therefore $(A \underset{R}{\otimes} _)(f) \in \mathrm{Hom}_R(A \underset{R}{\otimes} B , A \underset{R}{\otimes} C)$.

If $g \in G$, then $g(\Sigma_i (x_i \otimes y_i)) = \Sigma_i (gx_i \otimes gy_i)$.

$(A \underset{R}{\otimes} _)(f)(\Sigma_i (gx_i \otimes gy_i)) = \Sigma_i (gx_i \otimes f(gy_i)) = \Sigma_i (gx_i \otimes gf(y_i))$ since

$f \in \mathrm{Hom}_{RG}(B,C)$. Then $(A \underset{R}{\otimes} _)(f)(\Sigma_i (gx_i \otimes gy_i)) = \Sigma_i g(x_i \otimes f(y_i))$.

This implies $(A \underset{R}{\otimes} _)(f) \in \mathrm{Hom}_{RG}(A \underset{R}{\otimes} B , A \underset{R}{\otimes} C)$. Therefore

$(A \otimes _)(f) : A \underset{R}{\otimes} B \to A \underset{R}{\otimes} C$ is in $\mathrm{Mod}(R\text{-}G)$.

(i) and (ii) imply that $A \underset{R}{\otimes} _$ is a functor from $\mathrm{Mod}(R\text{-}G)$ to

$\mathrm{Mod}(R\text{-}G)$.

Now we prove the second assertion. Define Φ :

$\mathrm{Hom}_{RG}(B, {}^*\mathrm{Hom}(A,C)) \to \mathrm{Hom}_{RG}(A \underset{R}{\otimes} B, C)$ as follows. If $f \in$

$\mathrm{Hom}_{RG}(B, \mathrm{Hom}^*(A,C))$ then let f_b denote $f(b)$ for $b \in B$. Then

$f_b : A \to C$ is an R-*module *homomorphism. Define $\Phi(f) : A \underset{R}{\otimes} B \to C$

by $\Sigma_i a_i \otimes b_i \to \Sigma_i f_{b_i}(a_i)$ for $a_i \in A$, $b_i \in B$. This map is the same

as the one constructed for proving the adjointness of the pair

$\mathrm{Hom}_R(A,C)$, $A \underset{R}{\otimes} _$ in Mod_R . Therefore, $\Phi(f) \in \mathrm{Hom}_R(A \underset{R}{\otimes} B, C)$.

It is sufficient to prove that $\Phi(f)$ preserves G-action. Suppose

$g \in G$. Then $\Phi(f)(g(a \otimes b)) = \Phi(f)(ga \otimes gb) = f_{gb}(ga)$. $f \in$

$\mathrm{Hom}_{RG}(B, {}^*\mathrm{Hom}_R(A,C))$ and therefore f preserves G-action. Therefore

$f(gb) = g \circ f(b) = g \circ f_b$. $g \circ f_b = gf_bg^{-1}$ and

$f_{gb}(ga) = gf_b g^{-1}(ga) = gf_b(a) = g\Phi(f)(a \otimes b)$.

Now define $\Psi : \mathrm{Hom}_{RG}(A \underset{R}{\otimes} B, C) \to \mathrm{Hom}_{RG}(B, {}^*\mathrm{Hom}(A,C))$ as follows. If $(f : A \underset{R}{\otimes} B \to C) \in \mathrm{Hom}_{RG}(A \underset{R}{\otimes} B, C)$ then let $\Psi(f) : B \to {}^*\mathrm{Hom}(A,C)$ be defined by $b \mapsto (f_b : A \to C)$ where $f_b(a) = f(a \otimes b)$ for all $a \in A$, $b \in B$. This is essentially the same map as the one for proving the adjointness of the pair $\mathrm{Hom}_R(A,C)$, $A \underset{R}{\otimes} _$ in Mod_R . Therefore, it is sufficient to prove that (i) $\Psi(f)(b) \in {}^*\mathrm{Hom}(A,C)$ for all $b \in B$ and (ii) $\Psi(f)$ preserves G-action.

(i) $b \in B$ and B is an R-*module. Therefore, let $\langle gbg^{-1} \mid g \in G\rangle = \langle b_1, b_2, \ldots, b_n \mid b_i \in B\rangle$. Then $\Psi(f)(b) = f_b \cdot gf_b g^{-1} : A \to C$ is defined by $a \mapsto gf_b(g^{-1}a) = gf(g^{-1}a \otimes b) = f(a \otimes gb) = \sum_i \lambda_i f(a \otimes b_i) = \sum_i \lambda_i f_{b_i}(a)$ with $a \in A$, $\lambda_i \in k$ for all i . That is, $\langle gf_b g^{-1} \mid g \in G\rangle \subseteq \langle f_{b_1}, f_{b_2}, \ldots, f_{b_n}\rangle$ and therefore finite dimensional over k .

(ii) $\Psi : \mathrm{Hom}_{RG}(A \underset{R}{\otimes} B, C) \to \mathrm{Hom}_{RG}(B, {}^*\mathrm{Hom}(A,C))$. If $f \in \mathrm{Hom}_{RG}(A \underset{R}{\otimes} B, C)$ and $g \in G$ then $\Psi(g \circ f) = \Psi(gfg^{-1}) = \Psi(gg^{-1}f)$ since f preserves G-action. Therefore, $\Psi(g \circ f) = \Psi(f)$. On the other hand, $g \circ \Psi(f) = g\Psi(f)g^{-1}$ and $g\Psi(f)g^{-1} : B \to {}^*\mathrm{Hom}(A,C)$ where $b \mapsto (g \circ f_{g^{-1}b} : A \to C)$ But $g \circ f_{g^{-1}b}(a) = gf_{g^{-1}b}(g^{-1}a) = gfg^{-1}(a \otimes b) = a \otimes b$. Therefore, $\Psi(g \circ f) = g \circ \Psi(f)$.

That Φ and Ψ are inverse to each other and the isomorphism $\mathrm{Hom}_{RG}(B, {}^*\mathrm{Hom}(A,C)) \cong \mathrm{Hom}_{RG}(A \underset{R}{\otimes} B, C)$ is natural in each variable follows from the isomorphism $\mathrm{Hom}_R(B, \mathrm{Hom}(A,C)) \cong \mathrm{Hom}_R(A \underset{R}{\otimes} B, C)$ and the fact that it is natural in each variable. This completes the proof of

Prop. 1.19.

Remark 1.20 The left exactness of ${}^*\mathrm{Hom}$ follows from that of Hom . So also the right exactness of $A \underset{R}{\otimes} -$.

Proposition 1.21 If M and M' are R-*modules then $\mathrm{Hom}_{RG}(M,M')$ is an abelian group under the usual addition of morphisms.

Proof: It is sufficient to prove that $\mathrm{Hom}_{RG}(M,M')$ is a subgroup of the abelian group $\mathrm{Hom}_R(M,M')$.

Suppose $\varphi \in \mathrm{Hom}_{RG}(M,M')$. $(-\varphi)$ preserves G-action. If $\varphi,\psi \in \mathrm{Hom}_{RG}(M,M')$ then $(\psi - \varphi)$ preserves G-action. Thus, $(-\varphi)$, $(\varphi - \psi)$ are in $\mathrm{Hom}_{RG}(M,M')$. This completes the proof of Prop. 1.21.

Proposition 1.22 Composition of morphisms is bilinear in Mod(R-G) . That is, given R-*modules M,N,P and (R-G)-homomorphisms $M \underset{f'}{\overset{f}{\rightrightarrows}} N$, $N \underset{g'}{\overset{g}{\rightrightarrows}} P$, the distributive laws $(g + g') \circ f = g \circ f + g' \circ f$ and $g \circ (f + f') = g \circ f + g \circ f'$ are satisfied.

Proof: The distributive laws are satisfied in Mod_R . But $(g + g')f$, gf , gf' , $gf + gf'$, $g(f + f')$ are (R-G)-homomorphisms. Therefore, the above equalities are true in Mod(R-G) also.

Proposition 1.23 Mod(R-G) has a zero object such that for each object $A \in$ Mod(R-G) there is a unique homomorphism $0 \to A$ and a unique morphism $A \to 0$.

Proof: The zero object 0 of Mod_R is an object of Mod(R-G) since the zero module is an R-*module. If M is an R-*module then each set $\mathrm{Hom}_{RG}(0,M)$ and $\mathrm{Hom}_{RG}(M,0)$ has exactly one element, the inclusion map and the zero map, respectively, for if they have more than one element then 0 cannot be the zero object in Mod_R . This completes the proof

of Prop. 1.23.

Proposition 1.24 For every pair of objects M,N in $\mathrm{Mod}(R\text{-}G)$ there is a diagram in the category $M \underset{i_1}{\overset{p_1}{\leftrightarrows}} C \underset{i_2}{\overset{p_2}{\leftrightarrows}} N$ with $p_1 \circ i_1 = 1_M$, $p_2 \circ i_2 = 1_N$ and $(i_1 \circ p_1) + (i_2 \circ p_2) = 1_C$.

Proof: M,N are in Mod_R. Therefore there is a diagram in Mod_R, $M \underset{i_1}{\overset{p_1}{\leftrightarrows}} C \underset{i_2}{\overset{p_2}{\leftrightarrows}} N$, where $C = M \oplus N$, p_1 and p_2 are projection maps, i_1 and i_2 are inclusion maps satisfying the above equalities. But $M \oplus N$ is an $R\text{-}^*$module (by 1.10). Projection and inclusion maps preserve G-action. Therefore, $M \underset{i_1}{\overset{p_1}{\leftrightarrows}} M \oplus N \underset{i_2}{\overset{p_2}{\leftrightarrows}} N$ is the required diagram in $\mathrm{Mod}(R\text{-}G)$. This completes the proof of Prop. 1.24.

Propositions 1.21, 1.22, 1.23 and 1.24 establish that $\mathrm{Mod}(R\text{-}G)$ is an additive category.

Proposition 1.25 If M and M' are $R\text{-}^*$modules and $f : M \to M'$ is and$(R\text{-}G)$-homomorphism then the kernel object and the cokernel object in Mod_R are $R\text{-}^*$modules. We denote them by $\ker f$ and $\mathrm{coker}\ f$, respectively.

Proof: $\ker f = \{m \in M \mid f(m) = 0\}$ is an R-module

(i) G-action on $\ker f$ is the same as that on M. Therefore, $g(rm) = (gr)(gm)$ for all $g \in G$, $r \in R$, $m \in \ker f$.

(ii) Let $m \in \ker f$ and $V_m = \langle gm \mid g \in G \rangle$. $f(gm) = gf(m) = 0$ for all $g \in G$. Therefore, $V_m \subseteq \ker f$ and $\ker f = \bigcup_{m \in \ker f} V_m$. V_m is a rational G-module since M is an $R\text{-}^*$module. Thus, $\ker f$ is a rational G-module and therefore an $R\text{-}^*$module.

coker $f = M'/f(M)$ is an R-module.

(i) G-action on $f(M)$ is the same as that on M'.

(ii) Any element in $f(M)$ is of the form $f(m)$ with $m \in M$. $V_{f(m)} = \langle gf(m) \mid g \in G \rangle \subseteq f(M)$. Thus, $f(M) = \bigcup_{m \in M} V_{f(m)}$. $V_{f(m)}$ is a rational G-module since M' is an R-*module. Therefore, $f(M)$ is a rational G-module.

(i) and (ii) establish that $f(M)$ is an R-sub*module of M'. By Prop. 1.16, coker f is an R-*module. This completes the proof of Prop. 1.25.

Proposition 1.26 If M and M' are R-*modules then every (R-G)-homomorphism $f : M \to M'$ has a kernel and a cokernel.

Proof: By Prop. 1.25, ker f and coker f are R-*modules.

(i) Consider the (R-G)-homomorphism $i : \ker f \to M$ which is the inclusion map. Then $f \circ i$ is the zero map. If $h : P \to M$ is any (R-G)-homomorphism such that $f \circ h$ is the zero map then there is a unique R-module homomorphism $h' : P \to \ker f$ with $h = i \circ h'$. That is, we have the following commutative diagram in Mod_R.



If $g \in G$, $x \in P$ then $i \circ h'(gx) = h(gx) = gh(x) = gh'(x)$ since i is the inclusion map. Therefore, $h'(gx) = gh'(x)$. That is, h' is an (R-G)-homomorphism. (h' is unique in Mod(R-G) since it is unique in Mod_R.) Therefore, $\ker f \xrightarrow{i} M$ is the kernel of $f : M \to M'$ in Mod(R-G).

(ii) Consider the canonical map $M \xrightarrow{\eta} M/\ker f$. η preserves G-action. $\eta \circ f$ is the zero map. If $h : M \to P$ is any (R-G)-homomorphism then there is a unique R-module homomorphism $h' : M/\ker f \to P$ with $h = h' \circ \eta$. That is, we have the following commutative diagram in Mod_R .



If $g \in G$ and $\bar{m} \in M/\ker f$ where $\bar{m} = \eta(m)$ with $m \in M$ then $h'(\bar{m}) = h(m)$. $h'(g\bar{m}) = h'(\overline{gm}) = h(gm) = gh(m) = gh'(\bar{m})$. h' preserves G-action. h' is unique in Mod(R-G) since it is unique in Mod_R . Therefore, $\eta : M \to M/\ker f$ is the cokernel of $f : M \to M'$ in Mod(R-G) . This completes the proof of Prop. 1.26.

Remark 1.27 The above proposition implies that if $f : M \to M'$ is an (R-G)-homomorphism then the kernel of f in Mod(R-G) is the kernel of f in Mod_R and the cokernel of f in Mod(R-G) is the cokernel of f in Mod_R .

Proposition 1.28 If $f : M \to M'$ is an (R-G)-homomorphism whose kernel is 0 , then f is the kernel of its cokernel. If $f : M \to M'$ is an (R-G)-homomorphism whose cokernel is 0 , then f is the cokernel of its kernel. An (R-G)-homomorphism whose kernel and cokernel are 0 is an (R-G)-isomorphism.

Proof: Mod_R is an abelian category. Therefore the above statement is true in Mod_R . By Remark 1.27, the proposition is true in Mod(R-G) . This completes the proof of Prop. 1.28.

Propositions 1.25, 1.26, 1.28 establish that Mod(R-G) is an abelian category.

A direct system $\{M,\pi\}$ over the directed set I, in Mod(R-G), means that to each $i \in I$, there is an R-*Module M_i and to each pair $(i,j) \in I \times I$ with $i < j$, there is an (R-G)-homomorphism $\pi_i^j : M_i \to M_j$ such that for all $i \in I$, π_i^i is the identity map and for $i < j < \ell$ in I, $\pi_j^\ell \circ \pi_i^j = \pi_i^\ell$.

Proposition 1.29 Let $\{M,\pi\}$ be a direct system over I in Mod(R-G). For each pair $(j,\ell) \in I \times I$ with $j < \ell$ and each $m_j \in M_j$, the element $\pi_j^\ell m_j - m_j$ is an element of $\oplus_i M_i$. Then there exists a smallest R-sub*module N of $\oplus_i M_i$ containing such elements as $\pi_j^\ell m_j - m_j$ for all pairs $(j,\ell) \in I \times I$ with $j < \ell$. Moreover the quotient module $\oplus_i M_i/N$ is an R-*module.

Proof: If N is an R-sub*module of $\oplus_i M_i$ then $\oplus_i M_i/N$ is an R-*module by Prop. 1.16. Therefore it is sufficient to prove that such an N exists. We prove this by using Zorn's Lemma. Let A be the collection of all R-sub*modules of $\oplus_i M_i$ containing the set of elements described in the proposition. This collection is nonempty since $\oplus_i M_i \in A$. A can be partially ordered by $\supseteq$. If $\{N_i\}$ is any chain in A then $\cap N_i$ is an R-*module in A and is minimal for $\{N_i\}$. Therefore, A has a minimal element N. This completes the proof of Proposition 1.29.

We now establish that the above quotient $\oplus_i M_i/N$ is the appropriate categorical definition of direct limit in the category

Mod(R-G) .

Proposition 1.30 Let $\{M,\pi\}$ be a direct system over the directed set I as in Prop. 1.29. Let $\oplus_i M_i/N$ be the R-*module described in Prop. 1.29. If P is an R-*module and for each $i \in I$ there are (R-G)-homomorphisms $f_i : M_i \to P$ such that $f_j \pi_i^j = f_i$ for all $i < j$, then there is a unique (R-G)-homomorphism $\Phi : \oplus_i M_i/N \to P$ such that $\Phi(x + N) = f_i(x)$ if $x \in M_i \subseteq \oplus_i M_i$.

Proof: $\eta : \oplus_i M_i \to \oplus_i M_i/N$ is the cokernel of $N \to \oplus_i M_i$. The set $\{f_i \mid i \in I\}$ induces $f : \oplus_i M_i \to P$ with $f(N) = 0$. Therefore there exists a $\Phi : \oplus_i M_i/N \to P$ satisfying the above conditions. This completes the proof of Prop. 1.30.

Proposition 1.31 Let P be an R-*module. Then P is the direct limit of the family of finitely generated R-sub*modules of P .

Proof: Let $A = \{M_i \mid i \in I\}$ be the collection of finitely generated R-sub*modules of P . $A \neq \phi$. For if $p \in P$, $p \neq 0$, then $RV_p \in A$. Let $M_i, M_j \in A$ and let $M_i = \langle m_1,\ldots,m_s\rangle$ and $M_j = \langle n_1,\ldots,n_r\rangle$ as R-*modules. $V_{m_i} = \langle gm_i \mid g \in G\rangle$ is a finitely generated vector space over k . Let $V_{m_i} = \langle m_{i_1}, m_{i_2},\ldots,m_{i_{\ell_i}}\rangle$, $1 \le i \le s$ and $V_{n_i} = \langle n_{i_1},\ldots,n_{i_{k_i}}\rangle$, $1 \le i \le r$ as k-vector spaces. Let $V = \langle m_{11},\ldots,m_{1s_1},\ldots,m_{s\ell_s},n_{11},\ldots,n_{rk_r}\rangle$ be a finitely generated vector space over k , which is G-stable. This implies that RV is a finitely generated R-*module. Moreover, $M_i + M_j \subseteq RV \in A$. That is, there exists a $k_0 \in I$ $\ni$ $RV = M_{k_0}$ and $M_i + M_j \subseteq M_{k_0}$. Define $i \le j$ if $M_i \subseteq M_j$ and let

$\mu_{ij} : M_i \to M_j$ be the embedding of M_i in M_j. By this definition I is a directed set. Then $\varinjlim M_i = \sum_i M_i = \bigcup_i M_i \subseteq P$. If $p \in P$ then $RV_p \in A$. This implies that $P \subseteq \bigcup M_i$. Therefore $P = \varinjlim M_i$. This completes the proof of Proposition 1.31.

CHAPTER II

*SEMI*SIMPLICITY

Starting with the definitions of a *simple *ring and a *simple R-*module, we define a *semi*simple R-*module. We show that if G is a connected linear algebraic group and R is a *simple *ring then every finitely generated R-*module is R-torsion free and R-projective. Thus a *simple *ring is an integral domain if G is a connected linear algebraic group. If further G is a connected, linearly reductive algebraic group and R is a *simple *ring then every R-*module is *semi*simple and, therefore, R-projective.

Notation: $\cong_G$, $\cong_R$, $\cong_{RG}$ denote G-module isomorphism, R-module isomorphism and R-G isomorphism, respectively.

Definition 2.1 An *ideal of a *ring R is an ideal which is a sub-*module of R . [M]

Definition 2.2 A *ring R is *simple if the only *ideals of R are the zero *ideal and R . [M]

Definition 2.3 An R-*module M is *simple if the only sub*modules of M are the zero *module and M .

Example of a *Simple *Ring 2.4

If G is a linear algebraic group over k then $k[G]$ is a finitely generated k-algebra. This can be made into a *ring by defining a G-action on $k[G]$. If $g \in G$, $f \in k[G]$ then let $g \circ f(h) = f(g^{-1}h)$ for all $h \in G$. If $\Delta : k[G] \to k[G] \otimes_k k[G]$ be comultiplication, then $\Delta(f) = \sum_1^n a_i \otimes b_i$ if $f(xy) = \sum_i a_i(x)b_i(y)$ for all $x,y \in G$. Then $(g \circ f)(x) = f(g^{-1}x) = \sum_i a_i(g^{-1})b_i(x)$. So $g \circ f = \sum_i a_i(g^{-1})b_i$ and $V_f = \langle gf \mid g \in G \rangle \subseteq \langle b_1, b_2, \ldots, b_n \rangle$. Moreover, $G \to GL(V_f)$ is known to be a k-morphism. Suppose I is a nonzero *ideal in $k[G]$. $I \subset k[G]$ implies $V(I) \subset G$, $V(I) \neq 0$ and $V(I)$ is closed in G. If $x \in V(I)$ with $x \neq 0$, $f \in I$ with $f \neq 0$ and $g \in G$ then $f(x) = 0$ and $g \circ f(x) = 0$. Therefore, $f(g^{-1}x) = 0$. This implies that $g^{-1}x \in V(I)$. That is, $Gx \subseteq V(I) \subset G$. But $Gx = G$. Therefore, $V(I) = G$. This implies that $I = 0$, which is a contradiction. Therefore the only *ideals of $k[G]$ are the zero *ideal and $k[G]$.

Remark 2.5 The following proposition leads to the definition of *semi-*simple R-*modules.

Proposition 2.6 Let G be a linear algebraic group over k, R a *ring and M a nonzero R-*module. Then the following conditions on M are equivalent.

(i) M is the sum of a family of *simple sub*modules of M.

(ii) M is the direct sum of a family of *simple sub*modules of M.

(iii) Every sub*module N of M is a direct summand. That is, there exists a sub*module N' of M such that $M = N \oplus N'$.

Proof: (i) $\Rightarrow$ (ii). Let $M = \sum_{i\in I} M_i$ be a sum (not necessarily direct) of *simple sub*modules where I is the indexing set. Let J be a maximal subset of I such that $M' = \sum_{i\in J} M_i$ is a direct sum. If $j \in I$, $M_j \cap M'$ is a sub*module of M_j. But M_j is *simple. Therefore $M_j \cap M' = M_j$ or the zero *module 0. If $M_j \cap M'$ is 0 then $M_j + M'$ is a direct sum contradicting the maximality of J. Therefore, $M_j \subseteq M'$ for all $j \in I$. This implies that $M = \sum_{j\in J} M_j$ which is a direct sum.

(ii) $\Rightarrow$ (iii). Let N be a sub*module of M where $M = \sum_{i\in I} M_i$ is a direct sum of *simple sub*modules of M. Let J be the maximal subset of I such that $M' = N + \sum_{i\in J} M_i$ is a direct sum. By repeating the same argument as above, $M' = M$. That is, N is a direct summand of M.

(iii) $\Rightarrow$ (i). It is sufficient to prove that M contains a *simple sub*module. If M is not *simple, let m be a nonzero element of M. Let $V_m = \langle gm \mid g \in G\rangle$. Then RV_m is a sub*module of M. If RV_m is not *simple, then replace M by RV_m. Now we prove that RV_m contains a *simple sub*module. For this, we use Zorn's Lemma. Let $A = \{M' \subset M \mid m \notin M', M' \text{ is a sub*module of } M\}$. Since M is not *simple, let N be a proper sub*module of M. Either $m \in N$ or $m \notin N$. Suppose $m \in N$. By (iii) there exists a nonzero sub*module N' of M such that $M = N \oplus N'$. Then $m \notin N'$ and, therefore, $N' \in A$. This proves that $A \neq \phi$.

Let $\{M_i\}$ be a chain in A. $\cup M_i$ is a sub*module of M and

$m \notin \cup M_i$. Therefore, $\cup M_i \in A$. By Zorn's Lemma A has a maximal element. Let M_1 be the maximal element in A . That is, if M' is a sub*module of M such that M' does not contain m and $M' \supset M_1$, then $M' = M_1$. Let $\bar{N}$ be a nonzero sub*module of M/M_1 where $\bar{N}$ is the canonical image of $N \subseteq M$ in M/M_1 . Since $\bar{N} \neq 0$, $N \neq M$, so $m \in N$. Since $M = RV_m$, $M = N$. Then $\bar{N} = M/M_1$. Therefore, M/M_1 is *simple. By (iii), there exists a sub*module M_2 of M such that $M = M_1 \oplus M_2$. The canonical map $\eta : M \twoheadrightarrow M/M_1$ induces an R-module isomorphism $f : M_2 \twoheadrightarrow M/M_1$. If $g \in G$, $m_2 \in M_2$ then $f(gm_2) = gm_2 + M_1 = g(m_2 + M_1) = gf(m_2)$. That is, f preserves G-action. Therefore, M_2 and M/M_1 are isomorphic as rational G-modules. Therefore, M_2 is *simple. Thus, M contains a *simple sub*module M_2 . This completes the proof of Prop. 2.6.

Definition 2.7 An R-*module satisfying the above three conditions is said to be a *semi*simple R-*module.

Definition 2.8 A *ring R is said to be *semi*simple if R is *semi-*simple as a *module over itself.

Proposition 2.9 Every sub*module and every factor *module of a *semi-*simple *module is *semi*simple.

Proof: (i) Let N be a sub*module of an R-*module M where M is *semi*simple. Let N' be the sum of all *simple sub*modules of N . Since M is *semi*simple, $M = N' \oplus M'$ where M' is a sub*module of M . If $x \in N$, $x \neq 0$ then $x = n' + m'$, $n' \in N'$ and $m' \in M'$. Therefore, $m' = x - n' \in N$ and $N = N' \oplus M' \cap N$, a direct sum. This contradicts the maximality of N' since $M' \cap N$ is a sub*module of M and therefore is either *simple or contains a *simple sub*module. Thus,

$N = N'$.

(ii) Let N be a sub*module of M . M/N is an R-*module. M is *semi*simple. Therefore, $M = N \oplus N'$. N' is the direct sum of *simple sub*modules of M , by (i). The canonical map $\eta : M \twoheadrightarrow M/N$ induces an R-module isomorphism $f : N' \twoheadrightarrow M/N$. If $g \in G$, $n' \in N'$ then $f(gn') = g(n' + N) = gf(n')$. That is, f preserves G-action. Therefore, N' and M/N are isomorphic as rational G-modules. Therefore M/N is the direct sum of *simple sub*modules of M since N' is. This completes the proof of Corollary 2.9.

Lemma 2.10 Let G be a connected linear algebraic group. If R is a *simple *ring and M a finitely generated nonzero R-*module then M is a torsion free R-module and R is an integral domain.

Proof: R is Noetherian. Therefore there are only finitely many associated primes of M . Let $\{M_1, M_2, \ldots, M_r\} = \text{ass}(M)$. Let each M_i be the annihilator, $\text{ann}(a_i)$, of $a_i \in M$, $a_i \neq 0$. The zero divisors $Z(R)$ of M is $\bigcup_i M_i$. We prove that each M_i is G-stable and, in fact, if $g \in G$ then $gM_i = M_i$ for all i .

(i) Let $xy \in g(M_i)$, $x \neq 0$, $y \neq 0$. Then $xy = ga$ for some $a \in M_i$. Therefore, $(g^{-1}x)(g^{-1}y) = a$ and is in M_i . Since M_i is a prime ideal either $g^{-1}x \in M_i$ or $g^{-1}y \in M_i$. That is, either $x \in g(M_i)$ or $y \in g(M_i)$ proving that $g(M_i)$ is a prime ideal for all $g \in G$ and for all i .

(ii) $M_i = \text{ann}(a_i)$, $a_i \in M$, $a_i \neq 0$. If $x \in g(M_i)$, $x \neq 0$, $g \in G$ then $x = ga$ for some $a \in M_i$, $a \neq 0$. Then $aa_i = 0$; $\Rightarrow g(aa_i) = 0$; $\Rightarrow (ga)(ga_i) = 0$; $\Rightarrow ga \in \text{ann}(ga_i)$; $\Rightarrow g(M_i) \subseteq \text{ann}(ga_i)$. Conversely,

if $r \in \text{ann}(ga_i)$ then $r(ga_i) = 0$; $\Rightarrow (g^{-1}r)a_i = 0$; $\Rightarrow g^{-1}r \in M_i$; $\Rightarrow r \in g(M_i) \Rightarrow \text{ann}(ga_i) \subseteq g(M_i)$; $\Rightarrow g(M_i) = \text{ann}(ga_i)$ for all i, where $ga_i \neq 0$. (i) and (ii) imply that $g(M_i) = M_j$ for some $1 \leq j \leq r$.

(iii) But G is connected and G permutes the finite number of elements $M_1, M_2, \ldots, M_r$. This implies that $g(M_i) = M_i$ for all i. Therefore, M_i is an *ideal of R for all i. But R is *simple. Therefore, $M_i = 0$ for all i. Then M is R-torsion-free.

R is a finitely generated R-*module. Therefore by the above result R is an integral domain.

This completes the proof of Lemma 2.10.

Proposition 2.11 Let G be a connected linear algebraic group. If R is a finitely generated k-algebra that is *simple and M a nonzero finitely generated R-*module, then M is R-projective.

Proof: We first establish that if S is a multiplicatively closed subset of R and $S^{-1}M$ is a free $S^{-1}R$-module generated by $\frac{m_1}{1}, \frac{m_2}{1}, \ldots, \frac{m_n}{1}$ with $m_i \in M$, for all i, then there exists an $\alpha_0 \in S$ such that

(a) $F = \sum_i Rm_i$ is a free R-module.

(b) If $S_1 = \{1, \alpha_0, \alpha_0^2, \ldots\}$ then $S_1^{-1}F = S_1^{-1}M$ and are free as $S_1^{-1}R$-modules with $\frac{m_1}{1}, \frac{m_2}{1}, \ldots, \frac{m_n}{1}$ as basis.

(i) R is an integral domain. Let K be its quotient field. An R-module homomorphism $\varphi : M \to K \otimes_R M = (R-0)^{-1}M$ defined as $m \mapsto 1 \otimes m$

for all $m \in M$ is injective. $(R-0)^{-1}M$ is a K-vector space and M is a finitely generated R-module. Therefore, $(R-0)^{-1}M$ is finite dimensional over K. Let $\frac{m_1}{1}, \ldots, \frac{m_n}{1}$ be the K-basis with $m_i \in M$ for all i.

(ii) Let $F = \sum Rm_i \subseteq M$. Suppose there exist $r_i \in R$, $1 \leq i \leq n$ such that $\sum_i r_i m_i = 0$. Then, $\sum_i r_i \frac{m_i}{1} = 0$. But $\sum_i r_i \frac{m_i}{1} \in (R-0)^{-1}M$, a K-vector space with $\frac{m_1}{1}, \ldots, \frac{m_n}{1}$ as basis. Therefore $r_i = 0$, $1 \leq i \leq n$. Therefore, F is a free R-module with $m_1, m_2, \ldots, m_n$ as basis.

Moreover, $F \subseteq M$ implies $(R-0)^{-1}F \subseteq (R-0)^{-1}M$. If $x \in (R-0)^{-1}M$, $x \neq 0$, then $x = \sum_i \alpha_i \frac{m_i}{1}$ with $\alpha_i \in K$ and $\sum_i \alpha_i \frac{m_i}{1} = \sum_i \frac{\alpha_i}{1} m_i \in (R-0)^{-1}F$. Thus, $(R-0)^{-1}F = (R-0)^{-1}M$. $(R-0)^{-1}M$ has a K-basis $\frac{m_1}{1}, \ldots, \frac{m_n}{1}$. Therefore, $(R-0)^{-1}F$ has $\frac{m_1}{1}, \ldots, \frac{m_n}{1}$ as K-basis. Moreover, $(R-0)^{-1}M/F = 0$. This implies that $\exists\ d_0 \in R-0$ such that $S_0^{-1}M/F = 0$ where $S_0 = \{1, d_0, d_0^2, \ldots\}$. That is, $S_0^{-1}M = S_0^{-1}F$.

(iii) Now we will prove that $S_0^{-1}F$ is a free $S_0^{-1}R$ module.

If $x \in S_0^{-1}F$ then $x = \sum_i \frac{r_i m_i}{d_0^{\alpha}}$ with $r_i \in R$, $\alpha \in Z^+$ for all i. Therefore, $S_0^{-1}F$ is generated by $\frac{m_1}{1}, \ldots, \frac{m_n}{1}$ over $S_0^{-1}R$.

Suppose there exist $r_1, \ldots, r_n \in R$ such that $\frac{r_1}{d_0^{\alpha_1}} \cdot \frac{m_1}{1} + \cdots + \frac{r_n}{d_0^{\alpha_n}} \cdot \frac{m_n}{1} = 0$ with $\alpha_i \in Z^+$ for all i. This implies $\sum_i \frac{r_i' m_i}{d_0^{\alpha}} = \frac{0}{d_0^{\beta}}$

for $\alpha,\beta \in Z^+$, $r_i' \in R$ for all i . Then there exists $d_0^{\gamma} \in S_0$ such that $d_0^{\gamma}d_0^{\beta}(\sum_i r_i'm_i) = 0$. But F is a free R-*module. Therefore, $r_i'd_0^{\gamma+\beta} = 0$ for all i . But R is an integral domain and $d_0^{\gamma+\beta} \neq 0$. Therefore, $r_i' = 0$ for all i . That is, $\frac{r_i}{d_0^{\alpha_i}} = 0$ for all i .

Thus, $S_0^{-1}F$ is free over $S_0^{-1}R$. Since $S_0^{-1}F = S_0^{-1}M$, each is generated by $\frac{m_1}{1}, \frac{m_2}{1}, \ldots, \frac{m_n}{1}$ over $S_0^{-1}R$.

Suppose S is a multiplicatively closed subset of R and $S^{-1}M$ is a free $S^{-1}R$-module generated by $\frac{m_1}{1}, \frac{m_2}{1}, \ldots, \frac{m_\ell}{1}$ with $m_i \in M$, for all i . Then we can replace K by $S^{-1}R$ in (i), (ii) and (iii) thus establishing (a) and (b).

(iv) Let M be a maximal ideal of R such that M_M is a free R_M-module with $\frac{m_1}{1}, \ldots, \frac{m_r}{1}$ as basis where $m_i \in M$ for all i . By what we have established above there exists a $d \in R - M$ such that if $F = \sum_i Rm_i$, then F is R-free and $S^{-1}F = S^{-1}M$ as $S^{-1}R$-modules where $S = \{1,d,d^2,\ldots\}$. Moreover, $S^{-1}F$ and $S^{-1}M$ are free $S^{-1}R$-modules with $\frac{m_1}{1}, \ldots, \frac{m_r}{1}$ as basis.

Let M' be any maximal ideal of R such that $d \notin M'$. This implies that $S \subseteq R - M'$. $S^{-1}F = S^{-1}M$ implies $(S^{-1}F)_{R-M'} = (S^{-1}M)_{R-M'}$. Then $(R - M')^{-1}F = (R - M')^{-1}M$ as $(R - M')^{-1}R$-modules. That is, $F_{M'} = M_{M'}$ as $R_{M'}$-modules. By an argument similar to that in (iii) we can establish that $F_{M'}$ is a free $R_{M'}$-module.

(v) Now we will establish that there exists a maximal ideal M_0 of R such that M_{M_0} is a free R_{M_0}-module.

R is a Hilbert ring and is a domain. Therefore the Jacobson radical $J = 0$. This implies that there is a maximal ideal M_0 of R such that $d_0 \notin M_0$ (d_0 defined in (ii)). For otherwise, $d_0 \in J$ which is a contradiction. Then $S_0 \subseteq R - M_0$. But by (iii), $S_0^{-1}F = S_0^{-1}M$. Therefore, by localizing at $R - M_0$, we have $F_{M_0} = M_{M_0}$. By an argument similar to that in (iii) we can establish that F_{M_0} is R_{M_0}-free. Therefore, M_{M_0} is R_{M_0}-free.

(vi) Let $\max(R)$ be the collection of maximal ideals of R. Let $X = \{M \in \max(R) \mid M_M$ is a free R_M-module$\}$. By (v), X is nonempty. If $M \in X$ then, by (iv), there exists a $d \notin M$, such that if M' is a maximal ideal of R not containing d then $M_{M'}$ is a free $R_{M'}$-module.

If $U_s = \{M \in \max(R) \mid s \notin M\}$ then U_s is a basic open set in $\max(R)$ under the Zariski topology on $\max(R)$. Therefore, $M \in X$ implies that $M \in U_d \subseteq X$ for some $d \in R - M$. Thus, X is open in $\max(R)$.

Now we will prove that X is G-stable. If $M \in X$ then M_M is a free R_M-module. Suppose for $g \in G$, $g(M) \subsetneq M'$ where M' is a maximal ideal of R. Then $M \subsetneq g^{-1}(M')$, which is a contradiction. Therefore, $g(M)$ is maximal for all $g \in G$. Now it is sufficient to prove that $M_{g(M)}$ is a free $R_{g(M)}$-module.

Let $\frac{m_1}{1}, \ldots, \frac{m_n}{1}$ be the basis of M_M as free R_M-module

with $m_i \in M$ for all i. If $g \in G$ then let N be the $R_{g(M)}$-module generated by $\frac{g(m_1)}{1}, \frac{g(m_2)}{1}, \ldots, \frac{g(m_n)}{1}$. If there exists $\frac{r_i}{s_i} \in (R - g(M))^{-1}R$ for all i such that $\sum_{i=1}^{n} \frac{r_i}{s_i}(g(m_i)) = \frac{0}{s}$ with $s \in R - g(M)$ then $g(\sum_{i=1}^{s} g^{-1}(r_i)g^{-1}(s_i')m_i) = 0$ with $g^{-1}s_i' \in R - M$. This can be written as $\sum_{i=1}^{s} \frac{g^{-1}(r_i)}{q_i}\frac{m_i}{1} = \frac{0}{q}$ with $q_i, q \in R - M$. But M_M is R_M-free. Therefore, $\frac{g^{-1}(r_i)}{q_i} = \frac{0}{p_i}$ with $q_i, p_i \in R - M$ for all i. Then, $p_i g^{-1}(r_i) = 0$ for all i. That is, $g(p_i)r_i = 0$ for all i. But $g(p_i) \neq 0$. Therefore, $r_i = 0$ for all i since R is an integral domain. Thus, N is $R_{g(M)}$-free. It remains to be proved that $M_{g(M)} \subseteq N$.

If $y \in M$, $y \neq 0$ then $y = g(g^{-1}y)$. $g^{-1}(y) = (r_1m_1 + r_2m_2 + \cdots + r_nm_n)$ for some $r_i \in R$ for all i. Therefore, $y = (gr_1)(gm_1) + \cdots + (gr_n)(gm_n)$ with $gr_i \in R$ for all i. Then $\frac{y}{1} \in N$. Thus $M_{g(M)} \subseteq N$ proving $M_{g(M)} = N$. $M_{g(M)}$ is $R_{g(M)}$-free for all $g \in G$. Therefore X is G-stable.

(vii) We now prove that M_M is R_M-free for every $M \in \max(R)$.

By (vi), X is open in $\max(R)$. Therefore, $\max(R) - X$ is closed in $\max(R)$ under Zariski topology and is G-stable. Suppose $\max(R) - X \neq \phi$. $(\max(R), R)$ is an affine algebraic set and $\max(R) - X$ is closed in $\max(R)$. This implies that there exists an ideal I of R such that $\max(R) - X = \{M \in \max(R) \mid M \supseteq I\}$. $\max(R) - X$ is G-stable since X is. Therefore, I is G-stable. This implies that I

is an *ideal. But R is *simple. Therefore, either $I = 0$ or $I = R$. But $I \neq R$. Therefore, $I = 0$. Then $\max(R) - X = \max(R)$. This implies that $X = \phi$ which is a contradiction. Therefore, $\max(R) - X = \phi$. That is, $\max(R) = X$. Therefore, M_M is R_M-free for every $M \in \max(R)$.

(viii) R is Noetherian. M is a finitely generated R-module, therefore is of finite presentation. Moreover, M_M is R_M-free for all maximal ideals M of R. Therefore, M is R-projective. [(K), 3.3.7] This completes the proof of Proposition 2.11.

Corollary 2.11.1 Let G be a connected linear algebraic group. If R is a finitely generated k-algebra that is *simple and M a nonzero R-*module then M is R-flat.

Proof: By Prop. 1.31, M is the direct limit of the family of finitely generated R-sub*modules of M. Every nonzero finitely generated R-*module is R-projective and the direct limit of a family of R-projective modules is R-flat. This completes the proof of Cor. 2.11.1.

We quote some definitions and results from Fogarty's Invariant Theory needed for further development of this theory.

Definition 2.12 If G is an affine group, we say that G is linearly reductive if every rational G-module is completely reducible. [F, 4.6]

Notation: If M is a rational G-module then $M^G = \{m \in M \mid gm = m, \forall\ g \in G\}$.

Definition 2.13 If M is a rational G-module, then M is said to be G-ergodic if $M^G = (0)$.

From now on we assume that G is a linearly reductive algebraic group.

Lemma 2.14 Any rational G-module M contains a unique G-ergodic

submodule M_G . Moreover, $M = M^G \oplus M_G$ and M_G is the unique G-complement of M^G in M . [F - 5.2]

Definition 2.15 Let M be a rational G-module. We denote by P_M the projection of M onto M^G whose kernel is M_G . P_M is called the Reynolds operator of M . [F]

Remark 2.16 From the uniqueness of the Reynolds operator, it follows that if $\eta : M \to M'$ is a G-homomorphism of rational G-modules, then $\eta \circ P_M = P_{M'} \circ \eta$. [F]

Remark 2.17 If M and N are rational G-modules and $\eta : M \to N$ is a G-homomorphism that is onto then the restriction of η to M^G , that is, $\eta : M^G \to N^G$, is onto.

Proof: By the property of Reynolds operator (2.16) we have the following commutative diagram of rational G-modules and G-homomorphisms.

$$\begin{array}{ccc} M & \xrightarrow{P_M} & M^G \\ {\scriptstyle\eta}\downarrow & & \downarrow{\scriptstyle\eta} \\ N & \xrightarrow[P_N]{} & N^G \end{array}$$

$\eta : M \to N$ is onto. Therefore, $P_N \circ \eta$ is onto. This implies that $\eta \circ P_M$ is onto. Thus, $\eta : M^G \to N^G$ is onto.

Lemma 2.18 If R is a *ring, N a finitely generated R-*module and M an R-*module then $\mathrm{Hom}_R(N,M)$ is an R-*module with G-action defined by $g \circ f = gfg^{-1}$ for all $g \in G$, $f \in \mathrm{Hom}_R(N,M)$. (Prop. 1.12) If $\mathrm{Hom}_{RG}(N,M)$ is the collection of (R-G)-homomorphisms of N into M then $\mathrm{Hom}_{RG}(N,M) = \mathrm{Hom}_R(N,M)^G$.

Proof: (i) If $f \in \mathrm{Hom}_R(N,M)^G$ and $g \in G$ then $g \circ f = f$. Therefore, $g \circ f(n) = gf(g^{-1}n) = f(n)$. That is, $f(g^{-1}n) = g^{-1}f(n)$.

Therefore, f preserves G-action which implies that $f \in \mathrm{Hom}_{RG}(N,M)$.

(ii) Conversely, if $f \in \mathrm{Hom}_{RG}(N,M)$ then $g \circ f(n) = gf(g^{-1}n) = gg^{-1}f(n) = f(n)$ for all $n \in N$, $g \in G$. That is, $f \in \mathrm{Hom}_R(N,M)^G$.

(i) and (ii) imply that $\mathrm{Hom}_{RG}(N,M) = \mathrm{Hom}_R(N,M)^G$. This completes the proof of Lemma 2.18.

Proposition 2.19 Let G be a linearly reductive algebraic group, R a finitely generated k-algebra that is *simple, N a finitely generated R-*module and M an R-*module. Let $p \in \mathrm{Hom}_R(N,M)$ and $f \in \mathrm{Hom}_{RG}(M,N)$ such that $f \circ p = I_N$, the identity map on N . Then there exists an (R-G)-homomorphism $h : N \to M$ such that $f \circ h = I_N$.

Proof: By Prop. 1.12, $\mathrm{Hom}_R(N,M)$ and $\mathrm{Hom}_R(N,N)$ are R-*modules with appropriate G-action.

(i) Consider the R-module homomorphism $\Phi : \mathrm{Hom}_R(N,M) \to \mathrm{Hom}_R(N,N)$, $\varphi \xrightarrow{\Phi} f \circ \varphi$ where $\varphi \in \mathrm{Hom}_R(N,M)$. If $g \in G$, then $\Phi(g \circ \varphi) = f \circ (g \circ \varphi) = f \circ (g\varphi g^{-1}) = g(f \circ \varphi)g^{-1} = g \circ \Phi(\varphi)$. Therefore, Φ is a G-homomorphism. Let $\psi \in \mathrm{Hom}_R(N,N)$ then $p \circ \psi \in \mathrm{Hom}_R(N,M)$ and $f \circ (p \circ \psi) = (f \circ p) \circ \psi = \psi$ since $f \circ p = I_N$. That is, $\Phi(p \circ \psi) = \psi$. Therefore, Φ is onto.

(ii) G is a linearly reductive algebraic group. Therefore, $\mathrm{Hom}_R(N,M) = \mathrm{Hom}_R(N,M)^G \oplus \mathrm{Hom}_R(N,M)_G$ and $\mathrm{Hom}_R(N,N) = \mathrm{Hom}_R(N,N)^G \oplus \mathrm{Hom}_R(N,N)_G$. (Lemma 2.14). Moreover, by Remark 2.17, the G-homomorphism $\Phi : \mathrm{Hom}_R(N,M)^G \to \mathrm{Hom}_R(N,N)^G$ is onto. But $I_N \in \mathrm{Hom}_R(N,N)^G$. This implies that there exists $h \in \mathrm{Hom}_R(N,M)^G$ such that $\Phi(h) = f \circ h = I_N$. It was proved in Lemma 2.18 that $\mathrm{Hom}_{RG}(N,M) = \mathrm{Hom}_R(N,M)^G$. Therefore, $h \in \mathrm{Hom}_{RG}(N,M)$ as required.

This completes the proof of Proposition 2.19.

Corollary 2.19.1 Let G be a connected linearly reductive algebraic group, R a finitely generated k-algebra that is *simple, M a finitely generated R-*module and N an R-*module. If $\varphi : M \to N$ is an $(R\text{-}G)$-homomorphism which is onto, then there exists an $(R\text{-}G)$-homomorphism $\pi : N \to M$ such that $\varphi \circ \pi = I_N$. Moreover, $M = \ker \varphi \oplus \pi(N)$, direct sum of R-*modules.

Proof: $\varphi : M \to N$ is an R-module homomorphism that is onto. M is a finitely generated R-module. Therefore, N is a finitely generated R-*module. By Prop. 2.11, N is a projective R-module. Therefore, φ splits. That is, there exists an R-module homomorphism $f : N \to M$ such that $\varphi \circ f = I_N$. But φ preserves G-action. Therefore, by Prop. 2.19, there exists an $(R\text{-}G)$-homomorphism π in $\mathrm{Hom}_{RG}(N,M)$ such that $\varphi \circ \pi = I_N$. Then $\pi : N \to M$ is the required $(R\text{-}G)$-homomorphism. Thus, $\ker \varphi \hookrightarrow M \underset{\pi}{\overset{\varphi}{\rightleftarrows}} N$ is a split exact sequence of R-*modules and $(R\text{-}G)$-homomorphisms. Therefore, $M = \ker \varphi \oplus \pi(N)$. This completes the proof of Corollary 2.19.1.

Proposition 2.20 Let G be a connected linearly reductive algebraic group and R a finitely generated k-algebra that is *simple. Then every finitely generated nonzero R-*module M is a direct sum of finitely generated *simple R-sub*modules of M. That is, M is *semi*simple.

Proof: R is a Noetherian ring. M is a finitely generated R-module. Therefore, M is a Noetherian R-*module. Therefore, by Prop. 2.6, it is sufficient to prove that every nonzero sub*module N of M is a direct summand of M.

Consider the exact sequence of R-*modules and $(R\text{-}G)$-homomorphisms

$N \xrightarrow{i} M \xrightarrow{\eta} M/N$ where i is the inclusion map and η is the canonical map $M \twoheadrightarrow M/N$. i and η are (R-G)-homomorphisms. M is a finitely generated R-*module. Therefore, by Corollary 2.19.1, $M = N \oplus \pi(M/N)$ where π is a (R-G)-homomorphism $\pi : M/N \to M$ such that $\eta \circ \pi = I_{M/N}$. Thus, N is a direct summand of M. This completes the proof of Proposition 2.20.

Corollary 2.20.1 If G is a connected linearly reductive algebraic group, R a finitely generated k-algebra that is *simple and V a finite dimensional rational G-module, then $R \otimes_k V$ is a finitely generated R-*module and, therefore, *semi*simple.

Proposition 2.21 Let G be a connected linearly reductive algebraic group and R a finitely generated k-algebra that is *simple. Then every nonzero R-*module M is the direct sum of finitely generated *simple R-sub*modules of M and, therefore, *semi*simple.

Proof: Let $A = \{N \subseteq M \mid N$ is a direct sum of finitely generated *simple sub*modules of $M\}$. M is a nonzero R-*module. Therefore, M contains a nonzero element m. Let $V_m = \langle gm \mid g \in G \rangle$. Then RV_m is a finitely generated R-*module. Therefore by Prop. 2.20, RV_m is a direct sum of finitely generated *simple sub*modules of RV_m and, therefore, of M. $RV_m \in A$, thus A is nonempty.

Let $\{M_i \mid i \in I\}$ be a chain in A. $\cup M_i \in A$ and contains each M_i. Therefore by Zorn's Lemma A has a maximal element. Let it be M'. If $M' \neq M$, then let $x \in M - M'$. RV_x is a finitely generated sub*module of M and $RV_x \not\subseteq M'$. Let $RV_x = N$ and $M'' = M' \oplus N$. $M' \neq M''$. Therefore, M''/M' is a nonzero R-*module and the natural map $\eta : M'' \to M''/M'$ is an (R-G)-homomorphism that is onto.

M''/M' is a finitely generated R-*module. Therefore by Prop. 2.11, M''/M' is R-projective. Therefore there exists an R-module homomorphism $\pi : M''/M' \to M''$ such that $\eta \circ \pi = I_{M''/M'}$. But by Prop. 2.19, there exists an (R-G)-homomorphism $\pi' : M''/M' \to M''$ such that $\eta\circ\pi' = I_{M''/M'}$. Thus we have a split short exact sequence of R-*modules and (R-G)-homomorphisms $M' \xhookrightarrow{i} M'' \underset{\pi'}{\overset{\eta}{\twoheadrightarrow}} M''/M'$ where i is the inclusion map.

Therefore, $M'' = M' \oplus \pi'(M''/M')$. M''/M' is a finitely generated R-*module. π' is an (R-G)-homomorphism. Therefore, $\pi'(M''/M')$ is a finitely generated sub*module of M and therefore, *semi*simple. Let $\pi'(M''/M') = \bigoplus_i N_i$ where each N_i is a finitely generated *simple sub-*module of M. Then $M'' = M' \oplus (\bigoplus_i N_i)$. This contradicts the maximality of M'. Therefore, $M = M'$. This completes the proof of Proposition 2.21.

Corollary 2.21.1 Let G be a connected linearly reductive algebraic group. If R is a finitely generated k-algebra that is *simple, then every nonzero R-*module M is R-projective.

Proof: Every nonzero R-*module M is the direct sum of finitely generated R-sub*modules of M. Every finitely generated R-*module is R-projective by 2.11. Therefore, M is R-projective. This completes the proof of Corollary 2.21.1.

Proposition 2.22 Let G be a connected linearly reductive algebraic group. If R is a finitely generated k-algebra that is *simple and M is a nonzero sub*module of $R^{(n)}$, $n < \infty$, then $M = R^{(m)}$ for some m, $1 \le m \le n$.

Proof: Let $\pi_i : R^{(n)} \twoheadrightarrow R$ be the projection map on the i-th

coordinate. Consider the sequence of R-*modules and (R-G)-homomorphisms given by $M \xhookrightarrow{\mu} R^{(n)} \xrightarrow{\pi_i} R$ where μ is the inclusion map. Then $\pi_i \circ \mu$ is an (R-G)-homomorphism of M into R. $\pi_i \circ \mu(M)$ is a sub-*module of R. But R is *simple. Therefore, $\pi_i \circ \mu(M) = 0$ or R.

Case (i) If M is *simple, then $\pi_i \circ \mu : M \to R$ is either the zero map or an (R-G)-isomorphism. But M is nonzero and therefore, $\pi_i \circ \mu$ is nonzero for some i. Then, $M \underset{RG}{\cong} R$.

Case (ii) If M is not *simple, then M is *semi*simple. Therefore, $M = \bigoplus_j M_j$ where M_j is a *simple sub*module of $R^{(n)}$, for all j. By Case (i), $M_j \underset{RG}{\cong} R$ for all j. Therefore, $M \underset{RG}{\cong} R^{(m)}$ for some $m \in Z^+$, $1 \le m \le n$ since M is an R-submodule of $R^{(n)}$. This completes the proof of Proposition 2.22.

CHAPTER III

$k[SL_n]$-*MODULES WITH SL_n-ACTION

$SL_n(k)$ is a connected linearly reductive algebraic group if $n \geq 2$ and the characteristic of k is zero. We denote $SL_n(k)$ by either SL_n or G and $k[SL_n]$ by R. If SL_n-action on R is defined by $g \circ f(h) = f(g^{-1}h)$ for all $g,h \in G$, $f \in k[SL_n]$ then $k[SL_n]$ is a *simple *ring by 2.4.

In this chapter we establish that every *simple R-*module is $(R\text{-}G)$-isomorphic to R. Consequently, every R-*module is $(R\text{-}G)$-isomorphic to either $R^{(n)}$, $n > 0$, or $R^{(X)}$.

The existence of, but not the explicit form of, the isomorphism follows from a general theorem of Cline, Parshall and Scott [CPS]. But in this chapter we give the explicit form of the isomorphism for $k[SL_n]$-*modules.

First, we introduce some notations and state the definitions and facts needed for the sequence of results that lead to the final statement. $R = k[SL_n] = k[x_{11}, x_{12}, \ldots, x_{nn}]$ with

$\Gamma_{11}x_{11} + \Gamma_{21}x_{21} + \cdots + \Gamma_{n1}x_{n1} = 1$ where Γ_{i1} is the cofactor of

x_{i1} in the determinant $\begin{vmatrix} x_{11} & x_{12} & \cdots & x_{1n} \\ x_{21} & x_{22} & \cdots & x_{2n} \\ \vdots & & & \\ x_{n1} & x_{n2} & \cdots & x_{nn} \end{vmatrix}$ and x_{ij} is the

coordinate function for all i,j .

Definition 3.1 Definition of G-action on R . If $g \in SL_n$ then let

$g = \begin{bmatrix} \alpha_{11} & \alpha_{12} & \cdots & \alpha_{1n} \\ \vdots & & & \\ \alpha_{n1} & \alpha_{n2} & \cdots & \alpha_{nn} \end{bmatrix}$ with $\alpha_{ij} \in k$ for all i,j . Then

$g^{-1} = \begin{bmatrix} A_{11} & A_{21} & \cdots & A_{n1} \\ \vdots & & & \\ A_{1n} & A_{2n} & \cdots & A_{nn} \end{bmatrix}$ where A_{ij} is the cofactor of α_{ij}

for all i,j .

SL_n-action is defined by $g \circ f(h) = f(g^{-1}h)$ for all $g,h \in SL_n$ and

$f \in k[SL_n]$. Suppose $h = \begin{bmatrix} a_{11} & \cdots & a_{1n} \\ a_{21} & \cdots & a_{2n} \\ \vdots & & \\ a_{n1} & \cdots & a_{nn} \end{bmatrix}$ with $a_{ij} \in k$ for all

i,j . Then $g \circ x_{ij}(h) = x_{ij}(g^{-1}h)$ where

$$g^{-1}h = \begin{bmatrix} A_{11} & \cdots & A_{n1} \\ \vdots & & \\ A_{1n} & \cdots & A_{nn} \end{bmatrix} \begin{bmatrix} a_{11} & \cdots & a_{1n} \\ a_{21} & \cdots & a_{2n} \\ \vdots & & \\ a_{n1} & \cdots & a_{nn} \end{bmatrix} .$$

$x_{ij}(g^{-1}h) = A_{1i}a_{1j} + A_{2i}a_{2j} + \cdots + A_{ni}a_{nj}$. Therefore, $g \circ x_{ij} = A_{1i}x_{1j} + A_{2i}x_{2j} + \cdots + A_{ni}x_{nj}$ for all i,j .

Some facts about G and R . 3.2

1. $\sum_{j=1}^{n} \Gamma_{j1}x_{j1} = 1$. Therefore, $\sum_{j=1}^{n} \Gamma_{j1}x_{j1}$ is a G-invariant element in R .
2. R is a *simple *ring. Therefore every finitely generated R-*module is projective, by 2.11.
3. $V = \langle x_{1j}, x_{2j}, \ldots, x_{nj} \rangle$ is a simple G-module for all $1 \leq j \leq n$.

   Proof: $V \cong k^{(n)}$ as k-vector spaces. Suppose $u,w \in k^{(n)}$ then there exists a $\sigma \in GL_n$ such that $\sigma u = w$. This implies that there is a $\sigma \in SL_n$ such that $\sigma u = w$. Then $\sigma(\langle u \rangle) = \sigma(\langle w \rangle)$. Suppose W is a G-submodule of V . Then W contains every line in W . Then, W = V . That is, V is G-simple.
4. $RV \subseteq R$ and RV is an R-*module. But R is *simple. Therefore, RV = R .

Definition 3.3 Let $R_n = \langle e_1, e_2, \ldots, e_n \rangle$ with G-action defined by $g^{-1} \circ e_i = \alpha_{i1}e_1 + \alpha_{i2}e_2 + \cdots + \alpha_{in}e_n$ for all i and g is as in 3.1. Then $g \circ e_i = A_{1i}e_1 + A_{2i}e_2 + \cdots + A_{ni}e_n$ for all i .

Remark 3.4 If $V = \langle x_{11}, x_{21}, \ldots, x_{n1} \rangle$ then $V \underset{G}{\cong} R_n$ by the G-module

homomorphism $f : V \to R_n$ defined by $x_{i1} \to e_i$ for all i. Thus, we have $R \otimes_k V \cong_{RG} R \otimes_k R_n$. Next we establish that $R \otimes_k V \cong_{RG} R^{(n)}$ and therefore $R \otimes_k V \cong_{RG} R \otimes_k R_n \cong_{RG} R^{(n)}$.

Isomorphism 3.5 $\quad R \otimes_k R_n \cong_{RG} R^{(n)}$.

Proof: By 3.4, it is sufficient to prove $R \otimes_k V \cong_{RG} R^{(n)}$.

(i) Consider the exact sequence of R-*modules and $(R\text{-}G)$-homomorphisms, $K \overset{i}{\rightarrowtail} R \otimes_k V \overset{\varphi}{\twoheadrightarrow} RV = R$ where $K = \ker \varphi$, i is the injection map and φ is defined as $1 \otimes v \mapsto v$ for all $v \in V$. $\sum_{i=1}^{n} \Gamma_{i1} \otimes x_{i1} \in R \otimes_k V$. Then $\varphi(\sum_{i=1}^{n} \Gamma_{i1} \otimes x_{i1}) = \sum_{i=1}^{n} \Gamma_{i1} x_{i1} = 1$. Define an R-homomorphism $\Pi : R \to R \otimes_k V$ by $1 \overset{\pi}{\longmapsto} \sum_{i=1}^{n} \Gamma_{i1} \otimes x_{i1}$. With G-action as defined in 3.1, for $g \in G$, we have

$$
\begin{aligned}
&g(\Gamma_{11} \otimes x_{11} + \Gamma_{21} \otimes x_{21} + \cdots + \Gamma_{n1} \otimes x_{n1}) \\
&\quad = \big((A_{11}\operatorname{cof} A_{n1} + A_{12}\operatorname{cof} A_{n2} + \cdots + A_{1n}\operatorname{cof} A_{nn})(\Gamma_{n1} \otimes x_{11}) \\
&\qquad + (A_{11}\operatorname{cof} A_{n-1,1} + A_{12}\operatorname{cof} A_{n-1,2} + \cdots + A_{1n}\operatorname{cof} A_{n-1,n})(\Gamma_{n-1,1} \otimes x_{11}) \\
&\qquad + \cdots + (A_{11}\operatorname{cof} A_{11} + A_{12}\operatorname{cof} A_{12} + \cdots + A_{1n}\operatorname{cof} A_{1n})(\Gamma_{11} \otimes x_{11})\big) \\
&\qquad + \cdots + \big((A_{n1}\operatorname{cof} A_{n1} + A_{n2}\operatorname{cof} A_{n2} + \cdots + A_{nn}\operatorname{cof} A_{nn})(\Gamma_{n1} \otimes x_{n1}) \\
&\qquad + (A_{n1}\operatorname{cof} A_{n-1,1} + A_{n2}\operatorname{cof} A_{n-1,2} + \cdots + A_{nn}\operatorname{cof} A_{n-1,n})(\Gamma_{n-1,1} \otimes x_{n1}) \\
&\qquad + \cdots + (A_{n1}\operatorname{cof} A_{11} + A_{n2}\operatorname{cof} A_{12} + \cdots + A_{nn}\operatorname{cof} A_{1n})(\Gamma_{11} \otimes x_{n1})\big) \\
&\quad = \Gamma_{11} \otimes x_{11} + \Gamma_{21} \otimes x_{21} + \cdots + \Gamma_{n1} \otimes x_{n1},
\end{aligned}
$$

since $\sum_{j=1}^{n} A_{ij} \operatorname{cof} A_{kj} = 0$ if $i \neq k$ and 1 if $i = k$. That is,

$\sum_{i=1}^{n} \Gamma_{il} \otimes x_{il}$ is G-invariant. Moreover, $r \xmapsto{\pi} r(\sum_{i=1}^{n} \Gamma_{il} \otimes x_{il})$.

Thus, φ and π are (R-G)-homomorphisms such that $\varphi \circ \pi = I_R$.

Therefore the above short exact sequence splits. Therefore,

$$R \otimes_k V \cong K \oplus R .$$
$$RG$$

(ii) Characterization of the elements of K. If $x \in R \otimes_k V$ then

$x - \pi \circ \varphi(x) \in K$. That is, $x - \varphi(x)\pi(1) \in K$. On the other hand, if

$x \in K$, then $x = \sum_{i=1}^{n} r_i \otimes x_{il}$ such that $\varphi(x) = \sum_i r_i x_{il} = 0$. Then

$\varphi(x)\pi(1) = 0$. That is, $x = x - \varphi(x)\pi(1)$. Therefore, $K =$

$\{x - \varphi(x)\pi(1) \mid x \in R \otimes_k V\}$. Let $x = b_1 \otimes x_{11} + b_2 \otimes x_{21} + \cdots +$

$b_n \otimes x_{n1}$ with $b_i \in R$ for all i.

$$x - \varphi(x)\pi(1) = b_1 \otimes x_{11} + b_2 \otimes x_{21} + \cdots + b_n \otimes x_{n1}$$
$$- \sum_i b_i x_{il}(\Gamma_{11} \otimes x_{11} + \cdots + \Gamma_{n1} \otimes x_{n1})$$
$$= \sum_{i=1}^{n} (-b_i\Gamma_{11}x_{il} \otimes x_{11} - b_i\Gamma_{21}x_{il} \otimes x_{21} - \cdots$$
$$+ b_i(1 - \Gamma_{il})x_{il} \otimes x_{il} - \cdots - b_i\Gamma_{n1}x_{il} \otimes x_{n1}) .$$

Coefficient of $b_i = -\Gamma_{11}x_{il} \otimes x_{11} - \Gamma_{21}x_{il} \otimes x_{21} - \cdots +$

$(1 - \Gamma_{il})x_{il} \otimes x_{il} - \cdots - \Gamma_{n1}x_{il} \otimes x_{n1}$. But from the determinantal

properties of $\det(x_{ij})$, $\Gamma_{j1}x_{il} + \Gamma_{j2}x_{i2} + \cdots + \Gamma_{jn}x_{in} = 0$ for

all $j \neq i$ and $\Gamma_{il}x_{il} + \Gamma_{i2}x_{i2} + \cdots + \Gamma_{in}x_{in} = 1$. Therefore, the

coefficient of b_i can be written as $\sum_{j=2}^{n} x_{ij}(\Gamma_{1j} \otimes x_{11} + \Gamma_{2j} \otimes x_{21} +$

$\cdots + \Gamma_{nj} \otimes x_{n1})$. Let $\alpha_j = \Gamma_{1j} \otimes x_{11} + \Gamma_{2j} \otimes x_{21} + \cdots + \Gamma_{nj} \otimes x_{n1}$

for all $2 \leq j \leq n$. α_j is G-invariant for all j.

$\varphi(\alpha_j) = \sum_{i=1}^{n} \Gamma_{ij}x_{i1} = 0$, $2 \le j \le n$, by the property of a determinant.

Therefore, $\alpha_j \in K$ for all j . On the other hand, if $y \in K$, then y can be written as $\sum_{i=1}^{n} b_i(x_{i2}\alpha_2 + x_{i3}\alpha_3 + \cdots + x_{in}\alpha_n)$ with $b_i \in R$, $1 \le i \le n$.

(iii) Now define an (R-G)-homomorphism $\Phi : R^{(n-1)} \to K$ as $(0,\ldots,1,0,\ldots,0) \mapsto \alpha_{i+1}$ where 1 is the i-th coordinate in $(0,\ldots,0,1,0,\ldots,0)$. Then $(r_i)_i \mapsto \sum_{i=1}^{n-1} r_i\alpha_{i+1}$ where $(r_i)_i \in R^{(n-1)}$. We will prove that Φ is an (R-G)-isomorphism.

If $(b_i)_i,(b_i')_i \in R^{(n-1)}$ then $(b_i)_i = (b_i')_i \Leftrightarrow b_i = b_i'$ for all i . Then $\sum_{i=1}^{n-1} b_i\alpha_{i+1} = \sum_{i=1}^{n-1} b_i'\alpha_{i+1}$. That is, $\Phi((b_i)) = \Phi((b_i'))$. Thus, Φ is well-defined. Let $y \in K$, then $y = \sum_{i=1}^{n} b_i(x_{i2}\alpha_2 + x_{i3}\alpha_3 + \cdots + x_{in}\alpha_n)$ with $b_i \in R$ for all i . Therefore, $\sum_{i=1}^{n} b_ix_{ij} \xrightarrow{\Phi}$ $\sum_{2\le j\le n} (\sum_{i=1}^{n} b_ix_{ij}\alpha_j)$ proving that Φ is onto. If $(b_i),(b_i') \in R^{(n-1)}$ $\Phi((b_i)) = \Phi((b_i'))$ then $\sum_{i=1}^{n} b_i\alpha_{i+1} = \sum_{i=1}^{n} b_i'\alpha_{i+1}$. That is, $\sum_{j=1}^{n} (\sum_{i=1}^{n-1} (b_i - b_i')\Gamma_{j,i+1}) \otimes x_{j1} = 0$. But $K \subseteq R \otimes_k V$ which is a free R-module generated by $\{1 \otimes x_{j1} \mid 1 \le j \le n\}$. Therefore, $\sum_{i=1}^{n-1} (b_i - b_i')\Gamma_{j,i+1} = 0$ for all $1 \le j \le n$. Multiplying the n equations successively by x_{j2} , $1 \le j \le n$ and adding, we get $\sum_{i=1}^{n-1} (\sum_{j=2}^{n} (b_i - b_i')x_{j2}\Gamma_{ji}) = 0$. Again by the properties of determinants

$(b_1 - b_1')1 = 0$. That is, $b_1 = b_1'$. Similarly, $b_i = b_i'$, $1 \le i \le n-1$. Thus, $(b_i)_i = (b_i')_i$ proving Φ is 1-1.

α_i is G-invariant for all i . Therefore, Φ is an (R-G)-homomorphism. Thus, $K \underset{RG}{\cong} R^{(n-1)}$. Therefore, $R \underset{k}{\otimes} V \underset{RG}{\cong} R^{(n)}$ by combining (i) and (ii). That is, $R \underset{k}{\otimes} R_n \underset{RG}{\cong} R^{(n)}$.

Isomorphism 3.6 If A is an R-*module then $A \underset{R}{\otimes} (R \underset{k}{\otimes} R_n) \underset{RG}{\cong} (A \underset{R}{\otimes} R) \underset{k}{\otimes} R_n$ and therefore, $A \underset{R}{\otimes} (R \underset{k}{\otimes} R_n) \underset{RG}{\cong} A \underset{k}{\otimes} R_n$.

Proof: $A \underset{R}{\otimes} (R \underset{k}{\otimes} R_n) \underset{Z}{\cong} (A \underset{R}{\otimes} R) \underset{k}{\otimes} R_n$. The Z-homomorphism is defined by $a \underset{R}{\otimes} (r \underset{k}{\otimes} r_n) \xmapsto{\varphi} (a \underset{R}{\otimes} r) \underset{k}{\otimes} r_n$ for $a \in A$, $r \in R$ and $r_n \in R_n$. It is sufficient to prove that this is an (R-G)-homomorphism. Let $\alpha \in R$. Then $\alpha(a \underset{R}{\otimes} (r \underset{k}{\otimes} r_n)) = \alpha a \underset{R}{\otimes} (r \underset{k}{\otimes} r_n)$ and $\alpha a \underset{R}{\otimes} (r \underset{k}{\otimes} r_n) \xmapsto{\varphi}$ $(\alpha a \underset{R}{\otimes} r) \underset{k}{\otimes} r_n = \alpha((a \underset{R}{\otimes} r) \underset{k}{\otimes} r_n) = \alpha\varphi(a \underset{R}{\otimes} (r \underset{k}{\otimes} r_n))$. Thus, φ is an R-homomorphism. If $g \in G$ then $g \circ (a \underset{R}{\otimes} (r \underset{k}{\otimes} r_n)) = ga \underset{R}{\otimes} g(r \underset{k}{\otimes} r_n) =$ $ga \underset{R}{\otimes} (gr \underset{k}{\otimes} gr_n)$ and $ga \underset{R}{\otimes} (gr \underset{k}{\otimes} gr_n) \xmapsto{\varphi} (ga \underset{R}{\otimes} gr) \underset{k}{\otimes} gr_n$ and $(ga \underset{R}{\otimes} gr) \underset{k}{\otimes} gr_n = g(a \underset{R}{\otimes} r) \underset{k}{\otimes} gr_n = g((a \underset{R}{\otimes} r) \underset{k}{\otimes} r_n) = g\varphi(a \underset{R}{\otimes} (r \underset{k}{\otimes} r_n))$. Thus, φ preserves G-action.

Remark 3.7 $\underset{R}{\otimes}$ can be replaced by $\underset{k}{\otimes}$ and we get the isomorphism $A \underset{k}{\otimes} (R \underset{k}{\otimes} R_n) \underset{RG}{\cong} (A \underset{k}{\otimes} R) \underset{k}{\otimes} R_n$.

Isomorphism 3.8 $R \underset{k}{\otimes} R_n^{\underset{k}{\otimes} d} \underset{RG}{\cong} R^{(n^d)}$, $d \ge 0$.

Proof: The proof is by induction on d .

(i) If $d = 1$ then $R \underset{k}{\otimes} R_n \underset{RG}{\cong} R^{(n)}$ by 3.5.

(ii) Induction Hypothesis: Let $R \underset{k}{\otimes} R_n^{\overset{\otimes}{k}\, d-1} \underset{RG}{\cong} R^{(n^{d-1})}$, $d > 1$.

In 3.6, replace A by $R \underset{k}{\otimes} R_n^{\overset{\otimes}{k}\, d-1}$. Then $\left[R \underset{k}{\otimes} R_n^{\overset{\otimes}{k}\, d-1}\right] \underset{R}{\otimes} (R \underset{k}{\otimes} R_n) \underset{RG}{\cong} \left[R \underset{k}{\otimes} R_n^{\overset{\otimes}{k}\, d-1}\right] \underset{k}{\otimes} R_n$. By 3.7, $\left[R \underset{k}{\otimes} R_n^{\overset{\otimes}{k}\, d-1}\right] \underset{k}{\otimes} R_n \underset{RG}{\cong} R \underset{k}{\otimes} R_n^{(d)}$. But

$(R \underset{k}{\otimes} R_n)^{\overset{\otimes}{k}\, d-1} \underset{R}{\otimes} (R \underset{k}{\otimes} R_n) \underset{RG}{\cong} R^{(n^{d-1})} \underset{R}{\otimes} R^{(n)}$ by induction hypothesis and

by 3.5. Thus, $R \underset{k}{\otimes} R_n^{(d)} \underset{RG}{\cong} R^{(n^{d-1})} \underset{R}{\otimes} R^{(n)} \underset{RG}{\cong} R^{(n^d)}$.

*Remark 3.9 $\rho : SL_n \to R_n$ is a faithful representation. Therefore $k[SL_n] = k[R_n + R_n^*]$ where $k[R_n + R_n^*]$ is the k-algebra generated by by $R_n + R_n^*$ over k. R_n^* is the dual of R_n. $R_n = \langle e_1, e_2, \ldots, e_n \rangle$ (Def. 3.3). $R_n^* = \langle e_1^*, e_2^*, \ldots, e_n^* \rangle$ (dual basis). R_n^* is a G-module G-action being defined by $g \circ e_n^*(x) = e_n^*(g^{-1}x)$ for all $g \in G$, $x \in R_n$.

Lemma 3.10 If W is any nonzero SL_n-module that is simple, then there exist $i > 0$, $1 \leq i < \infty$ and a SL_n-module homomorphism

$$\Phi : \bigoplus_{i=1}^{m} (R_n \oplus R_n^*)^{\overset{\otimes}{k}\, d_i} \to W \text{ that is onto.}$$

Proof: Choose $\varphi \in W^*$. For each $x \in W$, define $f_x \in k[SL_n]$ by

---

* We refer to "Representative functions on discrete groups and solvable arithmetic subgroups" by G. D. Mostow, American Journal, 1970 for the result.

$f_x(h) = \varphi(h^{-1}x)$ for all $h \in SL_n$. Then a map $\hat{\varphi} : W \to k[SL_n]$ can be defined by $x \longmapsto f_x$. Let $g \in SL_n$. Then $gx \xmapsto{\hat{\varphi}} f_{gx}$ and $f_{gx}(h) = \varphi(h^{-1}gx)$ and $g \circ f_x(h) = f_x(g^{-1}h) = \varphi(h^{-1}gx)$. Therefore $\hat{\varphi}(gx) = g \circ \hat{\varphi}(x)$. That $\hat{\varphi}$ is an SL_n-module homomorphism is verified easily. Since $\hat{\varphi}$ preserves SL_n-action and W is SL_n-simple, $\hat{\varphi}$ is SL_n-module injection.

Thus it is sufficient to prove the lemma for SL_n-submodules of $k[SL_n]$. Since $k[SL_n] = k[R_n + R_n^*]$, any SL_n-submodule W of $k[SL_n]$ satisfies $W \subseteq \sum_{i=1}^{\ell} (R_n + R_n^*)^{d_i}$ for some $\ell < \infty$. Since SL_n-modules are semisimple, W is a direct summand, hence, a homomorphic image of $\sum_{i=1}^{\ell} (R_n + R_n^*)^{d_i}$. That is, there is $f : \sum_{i=1}^{\ell} (R_n + R_n^*)^{d_i} \to W$ that is onto. Then we have $\bigoplus_{i=1}^{\ell} (R_n \oplus R_n^*)^{\otimes_k d_i} \xrightarrow{\text{onto}} \sum_{i=1}^{\ell} (R_n + R_n^*)^{d_i} \xrightarrow{f} W$. This completes the proof of Lemma 3.10.

Lemma 3.11 $R \otimes_k R_n^* \cong_{RG} R^{(n)}$ implies $R \otimes_k R_n^* \cong_{RG} R^{(W)}$.

Proof: $R_n^* \cong \mathrm{Hom}_k(R_n, k)$ as k-vector spaces. $\mathrm{Hom}_k(R_n, k)$ is a G-module, G-action being defined by $g \circ f(x) = f(g^{-1}x)$ for all $f \in \mathrm{Hom}_k(R_n, k)$. Then $R_n^* \cong_G \mathrm{Hom}_k(R_n, k)$. Thus $R \otimes_k R_n^* \cong_{RG} R \otimes_k \mathrm{Hom}_k(R_n, k)$. $R \otimes_k \mathrm{Hom}_k(R_n, k)$ is a free R-module with basis $1 \otimes e_i^*$, $1 \le i \le n$. $R \otimes_k R_n$ is a free R-module with basis $1 \otimes e_i$, $1 \le i \le n$. Therefore we can define $\Phi : R \otimes_k \mathrm{Hom}_k(R_n, k) \to \mathrm{Hom}_R(R \otimes_k R_n, R)$ by $1 \otimes e_i^* \mapsto \widehat{1 \otimes e_i^*}$ where $\widehat{1 \otimes e_i^*} : R \otimes_k R_n \to R$ is defined by $1 \otimes e_j \mapsto e_i^*(e_j)$ for $1 \le j \le n$. This completely defines Φ. That Φ is an R-module homomorphism follows from the fact

$\sum_{i=1}^{n} a_i \otimes e_i^* \to \widehat{\sum_i a_i \otimes e_i^*}$ where $\widehat{\sum a_i \otimes e_i^*} : R \otimes_k R_n \to R$ is defined by $1 \otimes e_j \mapsto a_j$ for $a_j \in R$ and $1 \le j \le n$. If $\widehat{\sum a_i \otimes e_i^*} = \widehat{\sum r_i \otimes e_i^*}$ for $a_i, r_i \in R$, $1 \le i \le n$, then $a_i = r_i$ for all i. Then $\sum_i a_i \otimes e_i^* = \sum_i r_i \otimes e_i^*$. Thus Φ is one-to-one. Let $h \in \mathrm{Hom}_R(R \otimes_k R_n, R)$. Let $h(1 \otimes e_i) = r_i$, $1 \le i \le n$. Then $\widehat{\sum r_i \otimes e_i^*} = h$. Thus Φ is onto. Let $g \in G$. Then $g \circ (1 \otimes e_i^*) = 1 \otimes g \circ e_i^*$, $\widehat{g \circ (1 \otimes e_i^*)} = \widehat{1 \otimes g \circ e_i^*}$ where $\widehat{1 \otimes g \circ e_i^*}$ is defined by $1 \otimes e_j \mapsto g \circ e_i^*(e_j) = e_i^*(g^{-1} e_j)$. On the other hand, $g \circ \widehat{1 \otimes e_i^*}(1 \otimes e_j) = \widehat{1 \otimes e_i^*}(1 \otimes g^{-1} e_j) = e_i^*(g^{-1} e_j)$. Thus Φ preserves G-action. Therefore $R \otimes_k \mathrm{Hom}_k(R_n, k) \cong_{RG} \mathrm{Hom}_R(R \otimes_k R_n, R)$. But $R \otimes_k R_n \cong_{RG} R^{(n)}$. Let $\alpha : R \otimes_k R_n \to R^{(n)}$ be the isomorphism. Then $\mathrm{Hom}_R(R \otimes_k R_n, R) \cong_R \mathrm{Hom}_R(R^{(n)}, R)$ and the isomorphism $\Phi : \mathrm{Hom}_R(R \otimes_k R_n, R) \to \mathrm{Hom}_R(R^{(n)}, R)$ is defined by $h \mapsto h\alpha^{-1}$ for all $h \in \mathrm{Hom}_R(R \otimes_k R_n, R)$. Let $g \in G$, then $g \circ h \mapsto (g \circ h)\alpha^{-1}$ where $(g \circ h)\alpha^{-1}(x) = gh(g^{-1}\alpha^{-1}x) = gh\alpha^{-1}(g^{-1}x)$ for all $x \in R^{(n)}$, since α preserves G-action. But $g \circ (h\alpha^{-1})(x) = gh\alpha^{-1}(g^{-1}x)$. That is, Φ preserves G-action. Thus, $\mathrm{Hom}_R(R \otimes_k R_n, R) \cong_{RG} \mathrm{Hom}_R(R^{(n)}, R)$. But $\mathrm{Hom}_R(R^{(n)}, R) \cong_R \prod_{i=1}^{n} \mathrm{Hom}_R(R, R) = \bigoplus_{i=1}^{n} \mathrm{Hom}_R(R, R)$ and the isomorphism is defined as follows. Let $\mu_i : R \to R^{(n)}$ be the R-module injection into the i-th coordinate for all $1 \le i \le n$. Then if $h \in \mathrm{Hom}_R(R^{(n)}, R)$,

$h \mapsto \bigoplus_{i=1}^{n} h \circ \mu_i$ under the above isomorphism. Let $g \in G$, then $g \circ h \mapsto \bigoplus_{i=1}^{n} (g \circ h) \circ \mu_i = g(\bigoplus_{i=1}^{n} h \circ \mu_i)$. Thus, $\mathrm{Hom}_R(R^{(n)},R) \underset{RG}{\cong} \bigoplus_{i=1}^{n} \mathrm{Hom}_R(R,R)$. $\mathrm{Hom}_R(R,R) \underset{R}{\cong} R$ and the isomorphism $\lambda : \mathrm{Hom}_R(R,R) \to R$ is defined by $f \mapsto f(1)$. If $g \in G$, then $g \circ f \mapsto (g \circ f)(1) = f(g^{-1}1) = f(1)$. On the other hand, $g \circ (f(1)) = f(1)$ since $f(1) \in k$. That is, λ preserves G-action. Therefore, $\mathrm{Hom}_R(R,R) \underset{RG}{\cong} R$. Thus, $R \underset{k}{\otimes} R_n^* \underset{RG}{\cong} R^{(n)}$. This completes the proof of Lemma 3.11.

Proposition 3.12 Every nonzero *simple R-*module is (R-G)-isomorphic to R.

Proof: Let M be a nonzero *simple R-*module. If $m \in M$ and $m \neq 0$, then let $V_m = \langle gm \mid g \in G \rangle$. Since G is linearly reductive, $V_m = \bigoplus_i V_i$ where each V_i is G-simple. Since M is *simple, $M = RV_i$ for each i. We choose one such V_i and let $V_i = V$. Then, $M = RV$ where V is a simple G-submodule of M. The map $\mu : R \underset{k}{\otimes} V \to RV$ defined by $r \otimes v \mapsto rv$ where $r \in R$, $v \in V$ is an (R-G)-homomorphism that is onto. $R \underset{k}{\otimes} V$ is a finitely generated R-*module. Therefore by Cor. 2.20.1, $R \underset{k}{\otimes} V \underset{RG}{\cong} RV + \ker \mu$. V is a simple G-module. By Lemma 3.10, there is a G-module homomorphism $\varphi : \bigoplus_{i=1}^{m} (R_n \oplus R_n^*)^{\underset{k}{\otimes} d_i} \twoheadrightarrow V$ with $m < \infty$, $d_i < \infty$. Then $R \underset{k}{\otimes} \left[\bigoplus_{i=1}^{m} (R_n \oplus R_n^*)^{\underset{k}{\otimes} d_i} \right] \xrightarrow{1 \otimes \varphi} R \underset{k}{\otimes} V$

is an (R-G)-homomorphism that is onto. $R \otimes_k \left[\sum_{i=1}^{m} (R_n \oplus R_n^*)^{\otimes_k d-1} \right]$ is a finitely generated R-*module. Therefore, by Cor. 2.20.1 again,

$R \otimes_k \left[\sum_{i=1}^{m} (R_n \oplus R_n^*)^{\otimes_k d_i} \right] \underset{RG}{\cong} \ker(1 \otimes \varphi) \oplus R \otimes_k V$. But

$R \otimes_k R_n^{\otimes_k d} \underset{RG}{\cong} R^{(n^d)}$ and $(R \otimes_k R_n^*)^{\otimes_k d} \underset{RG}{\cong} R^{(n^d)}$. Therefore, by Prop. 2.22, $R \otimes_k V \underset{RG}{\cong} R^{(m_0)}$ for some $m_0 > 0$. The same proposition applied to the (R-G)-isomorphism $R \otimes_k V \underset{RG}{\cong} RV \oplus \ker \mu$ gives $RV \underset{RG}{\cong} R^{(p)}$ for some $p > 0$. But $RV = M$ and M is *simple. Therefore, $M \underset{RG}{\cong} R$.

This completes the proof of Proposition 3.12.

Corollary 3.12.1 Every nonzero R-*module M is (R-G)-isomorphic to either $R^{(n)}$, $n < \infty$, or $R^{(\chi)}$.

Proof: By Prop. 2.21, M is the direct sum of *simple R-sub*modules of M . Therefore, by Prop. 3.12, $M \underset{RG}{\cong} R^{(n)}$, $n < \infty$, or $M \underset{RG}{\cong} R^{(\chi)}$.

This completes the proof of Cor. 3.12.1.

BIBLIOGRAPHY

[A] Alain, Robert. *Introduction to Algebraic Geometry Through Affine Algebraic Groups*, Queen's Papers in Pure and Applied Mathematics, No. 44, 1976.

[B] Bourbaki, N. *Commutative Algebra*, Hermann, Paris XV, France, 1972.

[CPS] Cline, Parshall and Scott. *An Imprimitivity Theorem for Algebraic Groups*, to appear.

[F] Fogarty. *Invariant Theory*, W. A. Benjamin Inc., New York, 1969.

[H] Humphreys, James E. *Linear Algebraic Groups*, Springer-Verlag, New York, 1975.

[HS] Hilton, P. J. and Stammbach, U. *A Course in Homological Algebra*, Springer-Verlag, New York, 1971.

[K] Knight, J. T. *Commutative Algebra*, Cambridge University Press, 1971.

[M] Magid, A. *Dimension in Rings with Solvable Algebraic Group Action*, to appear.

[M*] Mostow, G. D. *Representative Functions on Discrete Groups and Solvable Arithmetic Subgroups*, American Journal, 1971.

[MS] Maclane, Saunders. *Categories for the Working Mathematician*, Springer-Verlag, New York, 1971.

[S] Springer, T. A. S. *Invariant Theory*, Lecture Notes in Mathematics, Springer-Verlag, New York, 1977.