NK1 OF NONABELIAN GROUPS (NIL, WHITEHEAD)
Harmon, Dennis Ray
ProQuest Dissertations and Theses; 1984; Dissertations & Theses @ University of Oklahoma
pg. n/a



Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.
Downloaded from SHAREOK. Persistent Handle URL: https://hdl.handle.net/11244/5207

INFORMATION TO USERS

This reproduction was made from a copy of a document sent to us for microfilming. While the most advanced technology has been used to photograph and reproduce this document, the quality of the reproduction is heavily dependent upon the quality of the material submitted.

The following explanation of techniques is provided to help clarify markings or notations which may appear on this reproduction.

1. The sign or "target" for pages apparently lacking from the document photographed is "Missing Page(s)". If it was possible to obtain the missing page(s) or section, they are spliced into the film along with adjacent pages. This may have necessitated cutting through an image and duplicating adjacent pages to assure complete continuity.

2. When an image on the film is obliterated with a round black mark, it is an indication of either blurred copy because of movement during exposure, duplicate copy, or copyrighted materials that should not have been filmed. For blurred pages, a good image of the page can be found in the adjacent frame. If copyrighted materials were deleted, a target note will appear listing the pages in the adjacent frame.

3. When a map, drawing or chart, etc., is part of the material being photographed, a definite method of "sectioning" the material has been followed. It is customary to begin filming at the upper left hand corner of a large sheet and to continue from left to right in equal sections with small overlaps. If necessary, sectioning is continued again—beginning below the first row and continuing on until complete.

4. For illustrations that cannot be satisfactorily reproduced by xerographic means, photographic prints can be purchased at additional cost and inserted into your xerographic copy. These prints are available upon request from the Dissertations Customer Services Department.

5. Some pages in any document may have indistinct print. In all cases the best available copy has been filmed.

University
Microfilms
International
300 N. Zeeb Road
Ann Arbor, MI 48106

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

8413975

Harmon, Dennis Ray

NK1 OF NONABELIAN GROUPS

The University of Oklahoma PH.D. 1984

University
Microfilms
International 300 N. Zeeb Road, Ann Arbor, MI 48106

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

THE UNIVERSITY OF OKLAHOMA

GRADUATE COLLEGE

NK_1 OF NONABELIAN GROUPS

A DISSERTATION

SUBMITTED TO THE GRADUATE FACULTY

in partial fulfillment of the requirements for the

degree of

DOCTOR OF PHILOSOPHY

By

DENNIS RAY HARMON

Norman, Oklahoma

1984

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

NK_1 OF NONABELIAN GROUPS

A DISSERTATION

APPROVED FOR THE DEPARTMENT OF MATHEMATICS

By

Roger C. Alperin

Darryl McCullough

Kevin A. Hare

Gregory A. Parker

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

TABLE OF CONTENTS

		Page
INTRODUCTION	. . . . .	iv
Chapter		
I.	PRELIMINARIES	1
II.	ANOTHER FROBENIUS MODULE	4
III.	A VANISHING RESULT	8
IV.	A NONVANISHING RESULT	13
V.	NK_1 OF FREE PRODUCTS	17
BIBLIOGRAPHY	. . . . .	21

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

INTRODUCTION

Let Rng denote the category of associative rings with identity and let Z-mod denote the category of abelian groups. For a functor F: Rng $\to$ Z-mod the functors NF, LF: Rng $\to$ Z-mod are defined as follows: $NF(A) = \ker (F(A[t]) \to F(A))$ induced by the augmentation $A[t] \to A$ sending t to 0, and $LF(A) = \operatorname{coker} (F(A[t]) \oplus F(A[t^{-1}]) \to F(A[t,t^{-1}]))$ induced by the obvious inclusions. For a contracted functor F such as K_0, K_1 or SK_1 there is an isomorphism [1, Cor. 7.3, p. 663]

$$F(A[t_1,\ldots,t_n,t_1^{-1},\ldots,t_n^{-1}]) \cong (I + 2N + L)^n F(A).$$

In particular, for $F = K_1$ and $A = Z\pi$ the integral group ring,

$$K_1Z[Z \times \pi] \cong K_1Z\pi \oplus NK_1Z\pi \oplus NK_1Z\pi \oplus K_0Z\pi.$$

Martin [8, Thm. 2.2] showed that $NK_1Z\pi = 0$ for π a finite abelian group of square-free order.

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

This thesis extends Martin's results to the case of an arbitrary finite group. The method is outlined as follows:

We first reduce to the case where π is hyper-elementary by showing that $NK_1Z\pi$ is a Frobenius module over $G_0Z\pi$. Cartesian squares then give rise to natural Mayer-Vietoris exact sequences of K-groups [9] and hence exact sequences of NK-groups. We then use the fact that $NK_iA = 0$ for a regular ring A, $i = 0,1$ [3] and $i = 2$ [10].

Symbols used throughout this thesis are: Z = the integers, Q = the rationals, and C_m = the cyclic group of order m.

At this time I would like to extend my thanks to my thesis advisor, Bruce Magurn for his invaluable guidance and support.

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

NK_1 OF NONABELIAN GROUPS

CHAPTER I

PRELIMINARIES

We begin by stating some definitions due to Lam [6, p. 106]. Let Sg denote the category whose objects are finite groups and whose arrows are inclusions $i: \rho \to \pi$ whenever $\rho \prec \pi$. A Frobenius functor is a contravariant functor $F: Sg \to$ Comm. Ring with arrow map $(-)^*$, which is also a covariant functor $Sg \to Z$-mod under an arrow map $(-)_*$, such that $i_*(i^*(x)\cdot y) = x\cdot i_*(y)$ for $\rho \to \pi$ an arrow in Sg, $x \in F(\pi)$, $y \in F(\rho)$.

Let F be a Frobenius functor. An F-module is an object map and two arrow maps $M(-)$, $(-)^*$, $(-)_*: Sg \to Z$-mod such that $M(-)$, $(-)^*$ is a contravariant functor, $M(-)$, $(-)_*$ is a covariant functor, $M(\pi)$ is an $F(\pi)$-module for each finite group π, and for $i: \rho \to \pi$ in Sg,

$$i^*(x\cdot y) = i^*(x)\cdot i^*(y) \quad \text{for } x \in F(\pi),\ y \in M(\pi)$$
$$i_*(i^*(x)\cdot y) = x\cdot i_*(y) \quad \text{for } x \in F(\pi),\ y \in M(\rho)$$
$$i_*(x\cdot i^*(y)) = i_*(x)\cdot y \quad \text{for } x \in F(\rho),\ y \in M(\pi).$$

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

A morphism of Frobenius functors $F \to G$ is a transformation $F \to G$ natural with respect to both arrow maps. A morphism of F-modules is a transformation $f: M \to N$ natural with respect to both arrow maps, such that $f: M(\pi) \to N(\pi)$ is $F(\pi)$-linear for each finite group π.

For the remainder of this chapter, let R be a commutative ring with identity. For π a finite group and $R\pi$ the group ring of π over R, let $\mathcal{L}^R$ denote the category whose objects are the finitely generated left $R\pi$-modules which are projective as R-modules and let $G_0^R R\pi$ denote $K_0(\mathcal{L}^R)$, its Grothendieck group. $G_0^R R[-]$ is a Frobenius functor. For M,N objects in $\mathcal{L}^R$ the multiplication in $G_0^R R\pi$ is $[M]\cdot[N] = [M \overline{\otimes}_R N]$ where the bar " $-$ " denotes the diagonal action of π on $M \overline{\otimes}_R N$. An arrow $i: \rho \to \pi$ in Sg induces the homomorphisms $i_*: G_0^R R\rho \to G_0^R R\pi$ defined by $i_*[M] = [R\pi \otimes_{R\rho} M]$ for $[M] \in G_0^R R\rho$ and $i^*: G_0^R R\pi \to G_0^R R\rho$ defined by $i^*[N] = [N]$ for $[N] \in G_0^R R\pi$.

According to Lam [6, p. 114], $K_0R[-]$ and $K_1R[-]$ are $G_0^R R[-]$-modules. For $[M] \in G_0^R R\pi$, $[N] \in K_0 R\pi$ the action is $[M]\cdot[N] = [M \overline{\otimes}_R N]$, and for $[N,\alpha] \in K_1 R\pi$ the action is $[M]\cdot[N,\alpha] = [M \overline{\otimes}_R N, 1_M \otimes \alpha]$. An arrow $i: \rho \to \pi$ extends to a homomorphism $i: R\rho \to R\pi$ inducing homomor-

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

phisms on K_0 and K_1 by the usual restriction and extension of scalars.

Let $\mathcal{f}(R\pi)$ denote the category of finitely generated left $R\pi$-modules and let $G_0R\pi$ denote its Grothendieck group $K_0(\mathcal{f}(R\pi))$. For a regular ring R, $G_0R\pi \cong G_0^RR\pi$ [13, Thm. 1.2] giving us a ring structure on $G_0R\pi$ even when $R\pi$ is not commutative, i.e. when π is not abelian.

Let C_m denote the cyclic group of order m. A finite group π is hyperelementary if $C_m \lhd \pi$ and $[\pi : C_m] = p^n$ for some prime p, $p \nmid m$, and some integer $n \geq 0$. This is equivalent to $\pi \cong C_m \rtimes B$ with $|B| = p^n$. Let $\mathcal{H}$ be the collection of hyperelementary groups. By the work of Lam [6, p. 123],

$$M(\pi) = \sum_{\substack{\rho \in \mathcal{H} \\ \rho < \pi}} i_*M(\rho)$$

for any $G_0^RR[-]$-module $M[-]$. This technique for computing $M(\pi)$ is referred to as "hyperelementary induction" and will be used in Chapter III.

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

CHAPTER II

ANOTHER FROBENIUS MODULE

Let A,B be commutative rings with identity, $f: A \to B$ a ring homomorphism, π a finite group and $f_*: K_1A\pi \to K_1B\pi$ the induced homomorphism. For $[M] \in G_0^A A\pi$, $[N,\alpha] \in K_1A\pi$,

$$f_*([M]\cdot[N,\alpha]) = f_*([M \underset{A}{\bar{\otimes}} N,\ 1_M \otimes \alpha])$$

$$= [B\pi \underset{A\pi}{\otimes} (M \underset{A}{\bar{\otimes}} N),\ 1_{B\pi} \otimes (1_M \otimes \alpha)]$$

$$[M]\cdot f_*([N,\alpha]) = [B \underset{A}{\bar{\otimes}} M]\cdot[B\pi \underset{A\pi}{\otimes} N,\ 1_{B\pi} \otimes \alpha]$$

$$= [(B \underset{A}{\bar{\otimes}} M) \underset{B}{\bar{\otimes}} (B\pi \underset{A\pi}{\otimes} N),\ (1_B \otimes 1_M) \otimes (1_{B\pi} \otimes \alpha)]$$

Lemma 2.1: $B\pi \underset{A\pi}{\otimes} (M \underset{A}{\bar{\otimes}} N) \cong (B \underset{A}{\bar{\otimes}} M) \underset{B}{\bar{\otimes}} (B\pi \underset{A\pi}{\otimes} N)$ as $B\pi$-modules.

Proof: Define $\phi: B\pi \underset{A\pi}{\otimes} (M \underset{A}{\bar{\otimes}} N) \to (B \underset{A}{\bar{\otimes}} M) \underset{B}{\bar{\otimes}} (B\pi \underset{A\pi}{\otimes} N)$ by $(bg\otimes(m\otimes n)) = (b\otimes gm)\otimes(1\otimes gn)$ for $bg \in B\pi$, $m \in M$, $n \in N$. Then ϕ is π-linear:

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

for $h \in \pi$,

$$\begin{aligned}\phi(h\cdot(bg\otimes(m\otimes n))) &= \phi(bhg\otimes(m\otimes n))\\ &= (b\otimes hgm)\otimes(1\otimes hgn)\\ &= h\cdot((b\otimes gm)\otimes(1\otimes gn))\end{aligned}$$

by the diagonal π-actions and the trivial action of π on B. Thus ϕ is B-linear.

Define $\psi: (B \overset{}{\otimes}_A M) \overset{}{\otimes}_B (B\pi \otimes_{A\pi} N) \to B\pi \otimes_{A\pi} (M \overset{}{\otimes}_A N)$

by $\psi((b\otimes m)\otimes(rg\otimes n)) = rbg\otimes(g^{-1}m\otimes n)$ for $b \in B$, $rg \in B\pi$, $m \in M$, and $n \in N$. Then ψ is π-linear:

for $h \in \pi$,

$$\begin{aligned}\psi(h\cdot((b\otimes m)\otimes(rg\otimes n))) &= \psi((b\otimes hm)\otimes(rhg\otimes n))\\ &= rbhg\otimes(r(hg)^{-1}hm\otimes n)\\ &= rbhg\otimes(rg^{-1}m\otimes n)\\ &= h\cdot(rbg\otimes(g^{-1}m\otimes n))\end{aligned}$$

So ψ is $B\pi$-linear as well. Checking composites,

$$\begin{aligned}\phi\psi((b\otimes m)\otimes(rg\otimes n)) &= \phi(rbg\otimes(g^{-1}m\otimes n))\\ &= (rb\otimes g(g^{-1}m))\otimes(1\otimes gn)\\ &= (rb\otimes m)\otimes(1\otimes gn)\\ &= (b\otimes m)\otimes(rg\otimes n)\end{aligned}$$

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

$$\begin{aligned}\psi\phi(bg\otimes(m\otimes n) &= \psi((b\otimes gm)\otimes(1\otimes gn))\\ &= b\otimes(gm\otimes gn)\\ &= bg\otimes(m\otimes n).\end{aligned}$$

Thus. ϕ and ψ are inverses and the isomorphism has been proved.

By the commutative square



and Lemma 2.1, $f_*([M]\cdot[N,\alpha]) = [M]\cdot f_*([N,\alpha])$ in $K_1B\pi$. For $i: \rho \to \pi$ in Sg, the $B\pi$-linear isomorphisms

$$B\pi \otimes_{B\rho} (B\rho \otimes_{A\rho} N) \cong B\pi \otimes_{A\rho} N \cong B\pi \otimes_{A\pi} (A\pi \otimes_{A\rho} N)$$

yield $i_*f_* = f_*i_*$. Also, $i^*f_* = f_*i^*$ so f is natural with respect to both arrow maps. Thus we have proved,

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

Theorem 2.2: For a homomorphism $f: A \to B$ of commutative rings, the induced homomorphism $f_*: K_1A[-] \to K_1B[-]$ is a morphism of $G_0^AA[-]$-modules.

Corollary 2.3: $NK_1Z[-]$ is a $G_0Z[-]$-module.

Proof: The category of $G_0^AA[-]$-modules is an abelian category [6, p. 108], so $\ker f_*$ is a $G_0^AA[-]$-module. The ring homomorphism $Z \to A$ induces a morphism $G_0Z[-] \cong G_0^ZZ[-] \to G_0^AA[-]$, making $\ker f_*$ a $G_0Z[-]$-module. Now apply Theorem 2.2 with $A = Z[t]$, $B = Z$, and f = augmentation.

Corollary 2.4: For any finite group π,

$$NK_1Z\pi = \sum_{\substack{\rho \in \mathcal{H} \\ \rho < \pi}} i_*NK_1Z\rho$$

where $\mathcal{H}$ is the class of hyperelementary groups.

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

CHAPTER III

A VANISHING RESULT

We are now in a position to prove the main result of this thesis.

Theorem 3.1: If π is a finite group of square-free order then $NK_1Z\pi = 0$.

Proof: By Corollary 2.4 we can assume that π is hyper-elementary. Let $\pi = C_m \rtimes B$ with $|B| = p$, p a prime and $p \nmid m$. Let $C_m = \langle a \rangle$, $C_p = B = \langle b \rangle$. Then π has the presentation

$$\pi = \langle a,b: a^m, b^p, bab^{-1}a^{-\alpha} \rangle$$

where $\alpha^p \equiv 1 \pmod{m}$. If $\alpha \equiv 1 \pmod{m}$ then π is abelian and thus $NK_1Z\pi = 0$ [8, Thm. 2.2].

For a divisor d of m, let ζ_d denote a complex primitive d^{th} root of unity. The twisted group algebra $Q(\zeta_d)^{\circ}B$ has the additive structure of a free right $Q(\zeta_d)$-module based on B and multiplicative structure determined

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

by that in $Q(\zeta_d)$, by that in B, and the rule $ba = a^{\alpha}b$.

For any collection M of positive divisors of m, let $\mathcal{O}(M)$ denote the image of $Z\pi$ under the composite

$$Z\pi \hookrightarrow Q\pi \xrightarrow{\cong} \bigoplus_{d|m} Q(\zeta_d)\circ B \longrightarrow \bigoplus_{d\in M} Q(\zeta_d)\circ B$$

For each $d \in M$ there is a Cartesian square

$$\begin{array}{ccc} \mathcal{O}(M) & \xrightarrow{\phi} & \mathcal{O}(M-\{d\}) \\ \downarrow{\scriptstyle\psi} & & \downarrow \\ \mathcal{O}(d) & \longrightarrow & \mathcal{O}(d)/J \end{array}$$

where ϕ,ψ are projections and $J = \psi(\ker \phi)$ is the ideal of $\mathcal{O}(d)$ generated by $\prod_{e\in M-\{d\}} \Phi_e(\zeta_d)$, and $\Phi_e(x)$ is the minimal polynomial of ζ_e over Q [7, pp. 403-4]. Then $\Phi_e(\zeta_d)$ is a unit of $Z[\zeta_d]$ if neither d/e nor e/d is a power of a prime, and is associate in $Z[\zeta_d]$ to a prime q if $e/d = q^r$ for some $r > 0$ [7, Lemma 9.3]. Choosing d minimal in M and using the fact that m is square-free we obtain

$$\prod_{e\in M-\{d\}} \Phi_e(\zeta_d) \sim \prod_{\substack{p_i \text{ prime} \\ p_i d\in M}} \Phi_{p_i d}(\zeta_d) \sim \prod_{\substack{p_i \text{ prime} \\ p_i d\in M}} p_i = \gamma$$

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

where " $\sim$ " means associate to. So for a minimal divisor $d \in M$, J is the ideal generated by γ.

If d is such that $B \to \mathrm{Aut}(Q(\zeta_d))$ has a non-trivial kernel then $\mathcal{O}(d)$ is in fact the group ring $Z[\zeta_d]B$ and $NK_1 Z[\zeta_d]B = 0$ [8, Thm. 2.2]. This occurs exactly when $d \mid (m, \alpha-1)$.

Otherwise, B acts faithfully on $Q(\zeta_d)$. Let $F = Q(\zeta_d)^B$, the subfield of $Q(\zeta_d)$ left fixed by the action of B. The cyclic algebra $Q(\zeta_d) \circ B = (Q(\zeta_d)/F, b, 1)$ [11, p. 259] is a crossed-product algebra with trivial factor set and hence isomorphic to $M_p(F)$ [11, Cor. 29.8]. Then $\dim_F Q(\zeta_d) = p$ and by choosing the integral basis $\{1, \zeta_d, \ldots, \zeta_d^{p-1}\}$ for $Q(\zeta_d)/F$, the isomorphism $Q(\zeta_d) \circ B \to M_p(F)$ above restricts to $Z[\zeta_d] \circ B = \mathcal{O}(d) \to M_p(R)$ where R is the ring of integers in $Q(\zeta_d)^B$.

For a nonzero prime $P \lhd R$, $P \cdot Z[\zeta_d] = P_1^e \cdots P_g^e$ where $P_1, \ldots, P_g$ are distinct maximal ideals in the Dedekind domain $Z[\zeta_d]$ and e is the ramification index. Since $e \mid p$, $e = 1$ or $e = p$. If $e = 1$ then P is unramified in $Z[\zeta_d]$. Otherwise $e = p$; but the only primes $P \lhd R$ that ramify are those for which $q \mid d$ where $Z \cap P = qZ$, and $p \nmid d$ since $p \nmid m$. Hence $e \neq 0$ in Z/qZ and P is tamely ramified in $Z[\zeta_d]$. So $Z[\zeta_d] \circ B$ is hereditary [11, Thm. 40.15] and hence regular. Thus $NK_i Z[\zeta_d] \circ B = 0$ for $i = 0, 1, 2$.

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

$M_p(R)$ is a maximal order containing $\mathcal{O}(d)$ and $d \cdot M_p(R) \subset \mathcal{O}(d)$, so $\mathcal{O}(d)/(\gamma) \cong M_p(R/(\gamma))$ [7, Prop. 10.2]. For each pair of primes p_i, p_j in the factorization of γ, $p_i Z + p_j Z = Z$ and so $p_i R + p_j R = R$. By the Chinese Remainder Theorem $R/(\gamma) \cong \bigoplus_i R/(p_i)$ and thus $M_p(R/(\gamma)) \cong \bigoplus_i M_p(R/(p_i))$. Since $p_i \nmid d$, (p_i) is unramified in $Z[\zeta_d]$, hence in R as well. So $(p_i) = \prod_j P_{ij}$ where $\{P_{ij}\}$ are distinct maximal ideals in R. Thus $\bigoplus_i M_p(R/(p_i)) \cong \bigoplus_{i,j} M_p(R/P_{ij})$, a direct sum of matrix rings over fields, hence regular. So $NK_i(\mathcal{O}(d)/(\gamma)) = 0$ for $i = 0,1,2$.

The Mayer-Vietoris exact sequence resulting from the Cartesian square is

$$NK_2\mathcal{O}(d)/(\gamma) \to NK_1\mathcal{O}(M) \to NK_1\mathcal{O}(M-\{d\}) \oplus NK_1\mathcal{O}(d) \to NK_1\mathcal{O}(d)/(\gamma)$$

and thus $NK_1\mathcal{O}(M) \cong NK_1\mathcal{O}(M-\{d\})$.

By iterating this procedure, starting with M the set of all divisors of m and peeling off the minimal divisor $d \in M$, we obtain $NK_1 Z\pi \cong NK_1\mathcal{O}(M) = 0$, proving the theorem.

Higher N's are defined recursively by $N^{j+1}K_1 = N(N^jK_1)$ for $j = 1,2,\ldots$ Using Theorem 2.2 with

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

$A = Z[s,t]$, $B = Z[t]$ and $f: s \to 0$ we can use hyperelementary induction to compute $N^2K_1Z\pi$. For π a hyperelementary group of square-free order, we tensor the Cartesian squares in the proof of Theorem 3.1 with $Z[t]$, producing new Cartesian squares and preserving regularity [2, Thm. 9.5]. Thus $NK_1Z\pi[t] = 0$. Continuing inductively we obtain

Corollary 3.2: $N^jK_1Z\pi = 0$ for π a finite group of square-free order, $j = 1,2,...$

Theorem 2.2 remains valid when K_1 is replaced by K_0 and so $NK_0Z[-]$ is also a $G_0Z[-]$-module. The Mayer-Vietoris sequences in the proof of Theorem 3.1 hold for NK_0 and $NK_0A = 0$ for A regular. Thus we have

Corollary 3.3: $N^jK_0Z\pi = 0$ for π a finite group of square-free order, $j = 1,2,...$

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

CHAPTER IV

A NONVANISHING RESULT

Theorem 4.1: Let R be the ring of integers in a number field F, π a finite group and $\alpha: \pi \to \rho$ a surjective homomorphism with ρ abelian. Then the induced homomorphism $\alpha_*: NK_1R\pi \to NK_1R\rho$ is surjective.

Proof: Let $e = (1/n) \cdot \sum_{x \in K} x$ where $K = \ker \alpha$ and $n = |K|$. Then e is a central idempotent in $F\pi$, and from the idempotents e and $1-e$ we obtain a Cartesian square

$$\begin{array}{ccc} R\pi & \xrightarrow{\beta} & \Lambda \\ \downarrow{\scriptstyle \alpha} & & \downarrow{\scriptstyle \delta} \\ R\rho & \xrightarrow[\gamma]{} & (R/nR)\rho \end{array}$$

where $\Lambda = R\pi/(\sum_{x \in K} x)$. We get a Mayer-Vietoris sequence

$$NK_1R\pi \xrightarrow{(\alpha_*, \beta_*)} NK_1R\rho \oplus NK_1\Lambda \xrightarrow{\gamma_* - \delta_*} NK_1(R/nR)\rho \ .$$

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

By inspection of the sequence it suffices to show $\gamma_*(NK_1R\rho) \subset \delta_*(NK_1\Lambda)$.

For any commutative ring A, the decomposition $K_1A \cong SK_1A \oplus U(A)$ gives $NK_1A \cong NSK_1A \oplus NU(A)$ where U is the functor $A \to$ units of A. If $f: A \to B$ is a homomorphism of commutative rings then the induced homomorphism $f_*: NK_1A \to NK_1B$ takes NSK_1A to NSK_1B and $NU(A)$ to $NU(B)$. Furthermore, $NU(A) \cong 1 + t \cdot Nil(A)[t]$ where $Nil(A)$ is the nilradical of A.

Since $R\rho$ is contained in the semisimple group algebra $(Q \otimes R)\rho$, $R\rho$ is reduced and so $NU(R\rho) = 0$. $(R/nR)\rho$ is a commutative Artinian ring and hence $NSK_1(R/nR)\rho = 0$ [1, Prop. 10.1, p. 685]. Therefore, $\gamma_*: NK_1R\rho \to NK_1(R/nR)\rho$ is the zero homomorphism and the assertion follows.

Corollary 4.2: Let π be a finite group and π^{ab} its abelianization. Then $NK_1Z\pi \to NK_1Z\pi^{ab}$ is surjective.

Martin [8, Thm. 3.12] showed that $NK_1ZC_{p^r} \neq 0$ for p an odd prime, $r \geq 2$ or $p = 2$, $r \geq 3$. So if π is a finite group with quotient C_{p^r} as above, then $NK_1Z\pi \neq 0$.

If $NK_1Z[C_2 \times C_2] \neq 0$ then for n even, $NK_1ZD_n \neq 0$ since $D_n^{ab} = C_2 \times C_2$. Here D_n denotes the

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

dihedral group of order $2n$. Whether $NK_1Z[C_2 \times C_2]$ vanishes or not is unknown at present. Knowledge of NK_2ZC_n would help in answering questions of this type.

We now observe that $NK_1Z[-]$ is in fact a Green module over $G_0Z[-]$:

For A a commutative ring and π a finite group, $K_1A[t]\pi$ has the Mackey subgroup property: for any groups $\rho, \rho' < \pi$ the composite

$$K_1A[t]\rho' \xrightarrow{i_*} K_1A[t]\pi \xrightarrow{i^*} K_1A[t]\rho$$

is equal to the sum, over all double cosets $\rho g \rho' \subset \pi$ of the composite

$$K_1A[t]\rho' \xrightarrow{i^*} K_1A[t](g^{-1}\rho g \cap \rho') \xrightarrow{\text{conj.}} K_1A[t](\rho \cap g\rho' g^{-1}) \xrightarrow{i_*} K_1A[t]\rho$$

This follows by the extension of the Mackey Subgroup Theorem to the case of an arbitrary commutative ring [4, p. 237]. Since $NK_1A\pi$ is a summand of $K_1A[t]\pi$ it inherits the above property, and with the $G_0^AA\pi$-module structure becomes a module over the Green functor $G_0^AA[-]$. By [5, Prop. 1.1' and 1.2],

$$NK_1Z\pi \cong \varinjlim_{\substack{\rho < \pi \\ \rho \in \mathcal{H}}} NK_1Z\rho \cong \varprojlim_{\substack{\rho < \pi \\ \rho \in \mathcal{H}}} NK_1Z\rho$$

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

where $\mathcal{H}$ is the collection of hyperelementary subgroups and the limits are taken with respect to inclusion and conjugation among subgroups in $\mathcal{H}$.

It is worth noting that the above observations hold for NK_0 as well.

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

CHAPTER V

NK_1 OF FREE PRODUCTS

Let Λ be an R-ring, i.e. R is a subring of Λ and there is a ring homomorphism $\varepsilon: \Lambda \to R$ such that $\varepsilon(r) = r$ for $r \in R$. The homomorphism $K_1R \to K_1\Lambda$ is injective, and denote the cokernel by $K_1(\Lambda;R)$.

Note that $\Lambda[t]$ is an $R[t]$-ring. The augmentation $t \to 0$ induces homomorphisms $K_1\Lambda[t] \to K_1\Lambda$ and $K_1R[t] \to K_1R$, thus inducing a homomorphism $K_1(\Lambda[t];R[t]) \to K_1(\Lambda;R)$. Denote the kernel by $NK_1(\Lambda;R)$.

We consider the case $R = Z$ and $\Lambda = Z\pi$ for any group π. Consider the diagram

$$
\begin{array}{ccccccccc}
 & & 0 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \to & NK_1Z & \to & K_1Z[t] & \to & K_1Z & \to & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \to & NK_1Z\pi & \to & K_1Z[t]\pi & \to & K_1Z\pi & \to & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \to & NK_1(Z\pi;Z) & \to & K_1(Z[t]\pi;Z[t]) & \to & K_1(Z\pi;Z) & \to & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 0 & &
\end{array}
$$

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

All rows and columns are exact, and $NK_1Z = 0$ since Z is regular. Thus $NK_1(Z\pi;Z) \cong NK_1Z\pi$.

Let ρ,η be arbitrary groups and let $\rho * \eta$ denote their free product. The integral group ring $Z[\rho * \eta]$ is isomorphic to the free product of rings $Z\rho *_Z Z\eta$. For a complete discussion of free products see [12, p. 355] or perhaps [1, p. 198].

$$(Z[\rho * \eta])[t] \cong (Z\rho *_Z Z\eta)[t]$$

$$\cong (Z\rho *_Z Z\eta) \otimes_Z Z[t]$$

$$\cong (Z\rho \otimes_Z Z[t]) *_{Z[t]} (Z\eta \otimes_Z Z[t]) \quad [1, p. 202]$$

$$\cong Z\rho[t] *_{Z[t]} Z\eta[t]$$

We state a result of Stallings [13, Cor. 6.2.1]:

Lemma 5.1: If R is a regular commutative ring, and if A and B are kernels of retractions of the R-rings Λ and Γ onto R, and if $A \otimes_R B$ is a flat R-module, then $K_1(\Lambda * \Gamma;R) \cong K_1(\Lambda;R) \oplus K_1(\Gamma;R)$.

We shall apply this result to the cases $R = Z, Z[t]$, $\Lambda = Z\rho, Z\rho[t]$ and $\Gamma = Z\eta, Z\eta[t]$.

Consider the diagram,

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

$$
\begin{array}{ccccccccc}
0 & \to & NK_1(Z[\rho * \eta];Z) & \to & K_1(Z[\rho * \eta][t];Z[t]) & \to & K_1(Z[\rho * \eta];Z) & \to & 0 \\
 & & \uparrow = & & \uparrow \cong & & \uparrow = & & \\
0 & \to & NK_1(Z[\rho * \eta];Z) & \to & K_1(Z\rho[t] *_{Z[t]} Z\eta[t];Z[t]) & \to & K_1(Z[\rho * \eta];Z) & \to & 0 \\
 & & \uparrow \psi \text{ induced} & & \uparrow \cong & & \uparrow \cong \text{ induced by } \rho\to\rho*\eta\leftarrow\eta & & \\
0 & \to & NK_1(Z\rho;Z) \oplus NK_1(Z\eta;Z) & \to & K_1(Z\rho[t];Z[t]) \oplus K_1(Z\eta[t];Z[t]) & \to & K_1(Z\rho;Z) \oplus K_1(Z\eta;Z) & \to & 0
\end{array}
$$

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

All rows are exact by definition. The two bottom-right vertical maps are isomorphisms by Lemma 5.1, and commute with the horizontal maps by the naturality of the isomorphisms of Bass and Stallings. Thus ψ is an isomorphism and we have proved,

Theorem 5.2: Let ρ,η be arbitrary groups. Then $NK_1Z[\rho * \eta] \cong NK_1Z\rho \oplus NK_1Z\pi$.

Corollary 5.3: $NK_1Z[F] = 0$ for any free group F of finite rank.

Proof:
$$\begin{aligned} NK_1Z[F] &\cong NK_1Z[Z * \cdots * Z] \\ &\cong NK_1Z[Z] \oplus \cdots \oplus NK_1Z[Z] \\ &\cong NK_1Z[t,t^{-1}] \oplus \cdots \oplus NK_1Z[t,t^{-1}] \\ &= 0 \end{aligned}$$

by the regularity of $Z[t,t^{-1}]$ [2, Cor. 9.7].

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

BIBLIOGRAPHY

1. Bass, H., Algebraic K-Theory, Benjamin, N.Y. (1968).

2. Bass, H., "Introduction to some methods of algebraic K-theory", CBSM Regional Conference, A.M.S. 20 (1974).

3. Bass, H., Heller, A., Swan, R., "The Whitehead group of a polynomial extension", Publ. Math. I.H.E.S. 22 (1964), 61-80.

4. Curtis, C., Reiner, I., Methods of Representation Theory, Vol. 1, John Wiley and Sons (1981).

5. Dress, A., "Induction and structure theorems for orthogonal representations of finite groups", Annals of Math. 102 (1975), 291-325.

6. Lam, T.-Y., "Induction theorems for Grothendieck groups and Whitehead groups of finite groups", Ann. Sci. Ecole Norm. Sup. 1 (1968), 91-148.

7. Magurn, B., "SK_1 of dihedral groups", J. Algebra 51 (1978), 399-415.

8. Martin, R., Thesis, Columbia University, 1975.

9. Milnor, J., Introduction to Algebraic K-Theory, Annals of Math. Studies 72, Princeton University Press, Princeton, N.J. (1971).

10. Quillen, D., "Higher algebraic K-theory I", Lecture Notes in Math., 341, Springer-Verlag (1973), 85-147.

11. Reiner, I., Maximal Orders, Academic Press, London (1975).

12. Stallings, J., "Whitehead torsion of free products", Annals of Math. 82 (1965), 354-363.

13. Swan, R., K-Theory of Finite Groups and Orders, Lecture Notes in Math., 149, Springer-Verlag (1970).

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.